

# Guidelines for Environmental Assurance in Australian Horticulture

### Developed by:

### HORTICULTURE FOR TOMORROW

Managed by Horticulture Australia Limited

### Second Edition – 2014

#### Disclaime

This guide has been produced by Horticulture Australia Ltd (HAL) to provide general information on environmental management in horticulture. The information has been provided in good faith, and the views expressed do not necessarily represent the views of HAL. HAL does not accept responsibility for the accuracy of the information provided and no responsibility or liability will be accepted for any use or reliance on the material contained in the document.

Laws governing environmental protection and management by horticultural producers can be complex, and may be include numerous Federal, State and Local Acts and Regulations. In addition, changes to these laws may be made from time to time. The user of this guide should seek expert advice from an appropriate professional or the relevant government agency to ensure the precise effect of current laws is fully understood before implementing any course of action referred to in this guide.





© 2014 – Horticulture Australia Limited. Parts of this document may be reproduced provided they are not sold for profit and the copyright owner is clearly identified on any reproduction.

#### Where to start?

How you use these guidelines is a matter of personal choice. Some people may like to start at the front cover and read to the last page – but you don't have to. The Guidelines are designed so you can get a quick overview of the issues and then prioritise the areas on which you want to focus for your business.

These guidelines have three main components:

- Introductory sections that provide background information about the issues and the guidelines;
- The main working sections to help you assess and manage your enterprise. These sections cover risk assessment, suggested practices, monitoring and recording options, and references and further resources; and
- A series of practical tools and resources to help you.

We recommend that you choose to start in one the following two ways.

#### 1: Start with the Risk Assessments

The guidelines cover a total of 20 different topics, divided into eight numbered sections. At the start of each major topic, there is a simple Risk Assessment diagram that asks key questions about your enterprise.

HIGH

RISK

LOW

RISK

Work your way through each Risk Assessment, answering the questions.

If you end up at a **HIGH RISK** sign you need to take some action. Read the Suggested Practices for the topic to explore your options.

If you end up at a **LOW RISK** sign you probably do not have a significant problem. However, you may want to read on to check your understanding of the issues.

By working through all the Risk Assessments you can prioritise issues that may need attention on your property.

#### or 2: Start with the Review Checklist

Another way to get an overview is to start with the Review Checklist. The checklist covers the major environmental assurance issues dealt with in the main text of the guidelines. By working through the checklist you will get an idea of your priority areas and you can then read the relevant topic in more detail.

The Checklist is divided into tables covering a range of topics. Select a topic and turn to the matching section in the guidelines and look for the Risk Assessment diagram. By working though the Risk Assessment you can quickly determine if that particular topic is relevant to your business.

If the topic is relevant to for your business, complete the relevant checklist table and record the answers.

Read the Suggested Practices for the topic to explore your management options.

The guidelines also provide information about monitoring options and sources of additional information for each topic.



Further resources

**Review checklist** 



2 Introduction

### Introduction

4 6 7 8 8 9 11 12 13 13 14 14 15 16 16 16 16 19 20
1.1 1.2 1.8 1.12 1.19 1.28 1.34 2.1 2.2 2.13 2.20 3.1 4.1 4.2 4.8 5.1 6.1 7.1 7.2 7.5 7.8 7.10 8.1
A.1
B.1 C.1 D.1 E.1
F.1



### Introduction

#### Acknowledgements

Title:	Guidelines for Environmental Assurance in Australian Horticulture – Second Edition 2014
Published by:	Horticulture Australia Ltd (HAL)
Date:	First printed 2006 (First Edition); updated 2014 (Second Edition)
Email:	horticulturefortomorrow@horticulture.com.au
Internet:	www.horticulturefortomorrow.com.au
Citation:	Kelly, A. (Ed, 2nd Edition) (2014); Lovell, J. (2006). Guidelines for Environmental Assurance in Australian Horticulture, Horticulture for Tomorrow, Horticulture Australia Ltd, Sydney.

The first edition of the Guidelines for Environmental Assurance in Australian Horticulture was originally launched in 2006 as an output from Horticulture for Tomorrow, a national project supported by Australia's horticultural industries, funded by the Natural Heritage Trust through the Australian Government's Pathways to Industry EMS Program, and managed by Horticulture Australia Ltd (HAL). For more background on the Horticulture for Tomorrow initiative, see the website www.horticulturefortomorrow.com.au

The original guidelines were developed by Jane Lovell, Tasmanian Quality Assured, with input and advice from many people.

This second edition of Guidelines for Environmental Assurance in Australian Horticulture was edited by Alison Kelly, Principle, Alison Kelly Consulting, and acknowledges the continual hard work of all involved in updating these guidelines.

The 2014 revision has been funded by HAL as part of the across-industry program. The Australian Government provides matched funds for all HAL's R&D activities.

The following people are particularly acknowledged for their contribution to the review:

- Richard Bennett, Product Integrity Manager, HAL;
- Kevin Bodnaruk, AKC Consulting Pty/Ltd;
- Anne-Maree Boland, Partner, RM Consulting Group;
- Jordan Brooke-Barnett, Environment Coordinator, Ausveg;
- Andy Chambers, Director, Seed Consulting Services;
- Clare Hamilton-Bate, National Program Manager, Freshcare Ltd;
- Anthony Kachenko, Research and Market Development Manager, Nursery & Garden Industry Australia;
- Naomi King, Development Horticulturist, AgriScience Queensland Department of Agriculture, Fisheries and Forestry;
- Andreas Klieber, Coles;
- Brenda Kranz, Portfolio Manager Natural Resources, Integrated Pest Management;
- Peter Melville, Portfolio Manager Natural Resources, Climate, Training and Leadership, HAL;
- Jane Muller, Senior Research and Policy Officer, Growcom;
- Simon Newett, Principle Extension Horticulturist, Agri-Science Queensland, Department of Agriculture, Fisheries and Forestry QLD;
- Charles Thompson, Senior Fellow, RM Consulting Group; and
- Production of this document has been managed by Rachel Bennett.

As part of the 2014 revision of this document, a new section on Climate Adaptation was developed and included. The author, Alison Kelly, acknowledges the following people for their contribution to the new section:

- Snow Barlow, Professor of Horticulture & Viticulture, University of Melbourne;
- Andy Chambers, Director, Seed Consulting Services;
- Peter Deuter, Senior Principal Horticulturist, Horticulture and Forestry Science, Department of Agriculture, Fisheries and Forestry (Queensland); and
- David Putland, Manager Energy & Climate, Growcom.

Significant cross-reference was made between this document and:

- Banana Best Management Practice Environmental Guidelines (2013);
- EnviroVeg: The Environmental Management System cycle Plan, Do, Check, Review - of continuous improvement (Version 3, August 2011); and
- Freshcare Environmental Code of Practice (2011); and
- Nursery Industry Accreditation Scheme Australia (NIASA) Best Management Practice Guidelines (5th Edition 2013) and Nursery & Garden Industry Australia Environmental Management System (EcoHort) for production nurseries (2nd Edition 2013).

#### Disclaimer

This guide has been produced by Horticulture Australia Ltd (HAL) to provide general information on environmental management in horticulture. The information has been provided in good faith, and the views expressed do not necessarily represent the views of HAL. HAL does not accept responsibility for the accuracy of the information provided and no responsibility or liability will be accepted for any use or reliance on the material contained in the document.

Laws governing environmental protection and management by horticultural producers can be complex, and may be include numerous Federal, State and Local Acts and Regulations. In addition, changes to these laws may be made from time to time. The user of this guide should seek expert advice from an appropriate professional or the relevant government agency to ensure the precise effect of current laws is fully understood before implementing any course of action referred to in this guide.

© 2014 – Horticulture Australia Limited. Parts of this document may be reproduced provided they are not sold for profit and the copyright owner is clearly identified on any reproduction



### Introduction



#### Introduction

#### Australian horticulture industry

Australia's horticultural industries produce fruit, vegetables, nuts, mushrooms, herbs and spices, nursery products, turf, cut flowers and extractive crops. Together these industries generated a farm-gate value of approximately \$8.7 billion in 2011-12, making horticulture Australia's third largest agricultural industry by value.

While estimates vary, the industry is made up of approximately 18,200 businesses (including wine and table grape production). The majority of businesses are small-scale family farms - though there is a trend towards an increasing number of medium to large scale operations.

Horticulture is a highly diverse industry operating in a broad range of locations and environments, and using a wide variety of production methods. Horticultural enterprises commonly operate in highly sensitive environments such as the Murray-Darling Basin, in catchments that drain to the Great Barrier Reef, or in close proximity to urban areas. The total area under horticultural production in Australia is estimated to be 289,300 hectares. Production locations tend to be concentrated in areas where there are fertile, well-drained soils, appropriate topography, reliable access to high guality water, and reasonable access to transport infrastructure and labour supply.

Australian horticultural industries make an important contribution to the nation's prosperity, especially in terms of providing food security, health and nutrition, and are a stimulus for regional economies. On-farm, healthy soils and water rights are major assets, along with crop genetics. See Figure 1 below.



Figure 1: Chart of horticulture's assests and contributions, based on the framework established by the National Land & Water Resources Audit (2008). Source: Day P (2010).



Across society, there is a move towards improving the environmental sustainability of our activities. Most industries invest in research and development that seeks to:

- Optimise the efficiency of natural resource use;
- Reduce the impacts of production or products on the environment;
- Monitor and document the environmental outcomes of management efforts.

The Australian horticulture industry has a strong record of seeking to proactively address matters of environmental sustainability. Horticulture Australia Ltd (HAL) plays an important role in facilitating investment in research, development and extension (RD&E) aimed at improving environmental management.

The National RD&E Framework for Horticulture (the Horticulture Framework) delivers the provisions of the National Framework for the horticultural industries, state government agencies working in horticulture, and Australian Government programs related to horticulture (including HAL). The Horticulture Framework was developed by the National Horticultural Research Network (NHRN) in 2010 and has been endorsed by all the horticultural industries, except the vegetable industry. HAL is a member of the NHRN.

#### Future environmental constraints for horticulture

The future holds many challenges. These include increasing competition for land and water, limits on inputs, the effects of changing rainfall and temperature patterns, threats from pests and weeds, pollution, overexploitation of resources, the impacts of consumer food choices and consumption, and increasing community expectations about how our land and oceans are managed.

Climate change as a consequence of the enhanced greenhouse effect will have both positive and negative impacts on horticulture. According to international scientific evidence, the most likely changes in our climate will be an increase in temperature of up to 2 °C over the next 50-100 years and more variable rainfall events. The warming of the atmosphere may also influence precipitation intensity, types and patterns, wind directions and intensity, and the frequency and severity of extreme weather events.

For further information on GLOBAL scenarios visit http://www.ipcc.ch/. For further information on DOMESTIC scenarios visit the CSIRO website http://www.csiro.au/ Outcomes/Climate/Climate-Change-Book.aspx (Specifically Chapter 3: Future Australian climate scenarios).

Climate change will not proceed smoothly. There is a risk of abrupt changes as the climate shifts from one state to another as a result of feedbacks in the climate system. This will mean increased risk for growers and changes in the way crops are grown. Consequently, adaptation options will need to expand from incremental adaptation options to more transformational strategies as greater levels of climate change are observed.

Besides the negative effects, potential benefits and fresh opportunities also arise from climate change. Higher temperatures may enhance production from horticulture and pastures in the continent's cool regions and the positive effects of higher levels of carbon dioxide on plant growth may partly offset the negative effects of higher evaporation or decreased rainfall.

See Section: Climate adaptation for more information.

Aside from the physical impacts of climate change on horticultural products and businesses, the industry will also be impacted by the global demand for food, increasing demand for productivity growth in response to this global demand, and the impacts of climate change policy.

The world's population is projected to reach around 8 billion by 2025. As the world economy shifts from west to east, millions of people are likely to move out of poverty and the middle class is predicted to grow from 1.8 billion in 2010 to 3.2 billion in 2020 and 4.9 billion in 2030. Increased demand for food across the world will create new export opportunities for the Australian food industry.

### Introduction



The quantity and quality of food we produce is directly affected by the condition of natural resourcesî- including biodiversity, soil, water, native vegetation and oceans. The future reliability and access to these resources in the future will be directly impacted by climate change. However, the effects of climate change will vary greatly between regions and industries.

The Australian horticulture sector has historically been successfully adapting to the challenges of changes in climate, water availability and weather extremes, and the industry continues to value improvements in production efficiencies and best management practices as approaches to managing ongoing variability and change.

Our food industry will only be able to grow if it can produce more food with the same or fewer natural resources. At the same time, growers will need to adapt their practices to reduce the impacts of unavoidable climate change and climate variability, and take advantage of potential opportunities from these processes. This will require a very high standard of crop management in order for producers to use and manage productive natural resources sustainably so we can produce food today and for future generations.

#### Horticulture and environmental assurance

The diversity of industries within horticulture means that a wide variety of production systems are used and businesses are located in a broad range of locations. Environmental management is often complex and multifaceted.

Most horticultural production systems are highly intensive (often highly reliant on irrigation, fertilisers and pesticides) and usually highly efficient (generating high yields per unit of land, water and fertilisers used).

For a horticultural business, environmental assurance is a means of demonstrating the use of management practices that achieve the level of environmental protection expected of itself and by its customers, the community and other interested parties.

A key feature of the environmental assurance process is risk assessment and using recognised practices to minimise or prevent (current or future) environmental impacts. A business can demonstrate management of environmental issues through self-assessment, or by seeking assessment from its customers or an independent (or third) party, typically an auditor.

#### Drivers for environmental assurance in horticulture

Horticultural producers have a fundamental interest and pivotal role in protecting, and where necessary and practical, enhancing and restoring natural resources. Assuring the long-term sustainability of natural resources directly affects the long-term sustainability of horticultural businesses.

There are many drivers for the development of environmental assurance programs in horticulture. The three main drivers identified through an environmental audit in 2011 (HAL Project Reference AH11005):

- Practices that improve production efficiency. The environmental assurance process may highlight areas where improvements can be made to benefit the sustainability of the business, for instance by reducing waste or nutrient leaching and thereby saving money:
- Improved or alternative inputs that reduce environmental impacts or increase efficiency. An environmental assurance process gives a horticultural producer confidence that their chosen management practices are effective in protecting natural assets such as soils and water, and minimising the risk of causing negative environmental impacts; and
- Evidence to demonstrate sound environmental performance or improvement over time. Put simply, community interest in environmental issues is increasing and our competitors are catching up fast. So it makes good business sense for the horticultural sector to develop a way to demonstrate its commitment to sound environmental and natural resource management.

Underpinning the above drivers are reducing potential for regulation, maintaining market access/meeting market demand, addressing consumer perceptions and meeting the needs of future generations under a changing climate.

The notion of 'clean and green' has been used to market Australian horticultural produce for years. Industry and government have put considerable effort into developing and implementing food safety programs that clearly justify the 'clean' label. Environmental management systems provide horticultural producers the tools to prove its 'green' environmental credentials.

Many horticultural markets are beginning to demand that their suppliers demonstrate an acceptable level of environmental management. Access to key markets may be jeopardised if horticultural businesses cannot provide this in addition to food safety and quality assurance.

Developing an industry-driven and industry-wide approach to environmental assurance demonstrates a broad commitment to environmental management. In turn, this commitment should alleviate the need to resort to increased regulation. Industry will also be in a better position to maintain long-term access to natural resources (especially water) by demonstrating sustainable use and management of these resources.

A widespread uptake of environmental assurance processes that are credible and consistent will help to maintain community confidence in the professionalism of the horticultural industry and its commitment to careful environmental management and protection of Australia's natural resources.

#### Environmental impacts associated with fresh produce

Horticultural activities have both positive and negative impacts on the external environment. Many relationships are complex (e.g. biodiversity) and poorly understood, let alone valued.

Potential environmental impacts are changes that may occur in the environment as a result of horticulture production practices. Risk assessment is required to determine whether the potential environmental impact is likely to occur within a particular enterprise, and to determine the likely significance of the impact, based on the management of each practice.

A comprehensive summary of environmental impacts associated with fresh produce is provided in an impact identification Table 1. The nursery industry's Eco-Hort program also outlines the environmental risks associated with nursery production systems.

The impact identification table details potential on-farm and off-farm environmental impacts from horticultural operations. The impacts are considered for each process step, making the link between activities and environmental 'hazards'.

In addition to these physical environmental issues, a number of other management considerations are also relevant, including:

- Environmental risk assessment processes;
- Climate risk assessment and adaptation planning;
- Farm planning, environmental management planning, and adaptive management systems;
- Environmental monitoring at farm, catchment and regional scales;
- Participation in market-based environmental assurance/certification schemes;
- Participation in community-based regional or catchment scale natural resource and environmental management initiatives;
- Compliance with environmental regulations.



Introduction

### Introduction



### Introduction

Category	Hazard	Specific environmental impact
Land and soil	Soil erosion	Sedimentation of rivers/waterways Reduction of water quality – nutrients and agricultural chemicals entering rivers/ waterways – eutrophication
	Soil structure	Compaction Increased run off Soil erosion, sedimentation of rivers/waterways Nutrient depletion
	Salinity	Reduction of arable land Spread of saline water and land Adverse impact on flora and fauna – loss of biodiversity
	Soil acidity and alkalinity	Loss of productivity Reduction of arable land
	Sodicity	Reduction of arable land Soil erosion Soil waterlogging
	Soil degradation	Compaction Increased run off Soil erosion, sedimentation of rivers/waterways Nutrient depletion
Water	Inefficient use of resources	Insufficient water supply/environmental flow Depletion of water table Adverse impact on flora and fauna – loss of biodiversity Rising water table and waterlogging Salinity Soil erosion Nutrient leaching Contamination of waterways
	Inappropriate water quality	Reduction of water quality Contamination by fertiliser, eutrophication Contamination by agricultural chemicals Contamination by fuels and oils Sedimentation
Chemicals	Inappropriate storage of chemicals	Contamination of surface/groundwater Contamination of drinking water Adverse effect on flora and fauna – loss of biodiversity
	Inappropriate application	Contamination of surface/groundwater Contamination of drinking water Adverse affect on flora and fauna – loss of biodiversity Soil contamination Adverse impact on other crops Adverse impact on neighbours
	Inappropriate disposal of agricultural chemicals, surplus agricultural chemicals, rinsates, chemical containers	Contamination of surface/groundwater Contamination of drinking water Adverse affect on flora and fauna – loss of biodiversity Soil contamination
	Spray drift	Disruption of Integrated Pest Management strategies Health risk for local residents Contamination of surface/groundwater Contamination of drinking water Adverse affect on flora and fauna – loss of biodiversity Adverse affect on surrounding crops
Nutrients	Inappropriate use of resources	Soil acidification Adverse impact on flora and fauna – loss of biodiversity
	Misplacement of fertiliser	Reduction of water quality – eutrophication Adverse impact on flora and fauna – loss of biodiversity
Biodiversity	Loss of biodiversity	Clearing of land Reduction of wildlife corridors Loss of aquatic habitat Change in pest species present

Category	Hazard	Specific environmental impact
Waste	Inappropriate disposal of waste	Contamination of soil and water Adverse affect on flora and fauna – loss of biodiversity Greenhouse gas emission – global warming and climate change Inconvenience to local residents
	Inefficient use of resources	Wasting non-renewable resources Greenhouses gas emission – global warming and climate change Waste disposal sites required (landfill)
Air	Dust	Sedimentation of waterways Soil erosion Inconvenience for local residents
	Smoke	Creation of greenhouse gases – global warming Inconvenience for local residents
	Noise	Disturbance Inconvenience for local residents Adverse impact on fauna – loss of biodiversity
Energy & greenhouse gases	Inefficient use of resources	Creation of greenhouse gases – global warming & climate change Wasting non-renewable resources

Table 1: Environmental Impacts Identification Table. Source: Lovell J (2006).

#### Levels of environmental systems

Each of these levels, environmental assurance, Best Management Practices and Environmental Management Systems, involve assessment of environmental impacts or risks, and all advocate appropriate actions to address environmentally significant issues.

#### Environmental assurance

Environmental assurance provides a generic checklist of recognised environmental best practices. It does not allow for certification in its own right; being completed through selfassessment and not third-party audited.

#### **Best Management Practices (BMPs)**

Best management practices (BMPs) provide specific techniques, operational practices and industry guidelines for establishing, achieving and reviewing best production methods and management of resources within a business. BMPs provide more detailed and tailored information than environmental assurance guidelines. BMPs may relate to any aspect of a business operation, e.g. food safety BMPs, environmental BMPs, etc. BMPs provide:

- Information on suggested management practices based on research and development and recognized industry best practices;
- Resources that may be specific to regional conditions or to industry or commodity requirements;
- A mechanism against which an individual business can review their existing practices. Implementation of BMPs are often a business's first step towards achieving formal certification to a recognized program or standard.

#### **Environmental Management Systems (EMS)**

An Environmental Management System, or EMS, describes any systematic approach to managing the impacts an enterprise has on the environment. EMS is one of the tools available to help a business deliver environmental assurance.

EMS provides a process for the business to keep track of all the information needed to demonstrate to both business managers and external parties it is meeting the environmental assurance standard established. EMS also offers businesses 'continuous improvement', that is, a risk assessment- based pathway to continuously improve their management systems. It encourages a business to 'plan, do, check and review' at regular intervals and across all aspects of the production cycle (see Table 2). A business would not necessarily need a comprehensive EMS in place to demonstrate compliance with the environmental assurance standard.

### Introduction



### Introduction

System	Version	Training	Certification	Branding	Online Tool
Environmental Assessment – self-assessment only					
Horticulture Environmental Assurance Guidelines	2006	No	No	No	PDF only
Best Management Practice – industry or regional self-assessment	tool				
Banana BMP	2013	Yes	No	No	Yes
EnviroVeg Self-assessment Tool	2011	Yes	No	Yes	Yes (members only)
Nursery Industry Accreditation Scheme Australia (NIASA) BMP (production nursery)	2013	Yes	No	No	No
Growcom's Farm Management System	2013	Yes	No	No	No
Pineapple BMP	2009	No	No	No	No
Freshcare Winery/ Viticulture	2011/2013	Yes	Yes	Yes	No
DairySAT	2012	No	No	No	Yes
Environmental Management System – environmental certification	through third party	y audit			
Freshcare Environment Code of Practice	2011	Yes	Yes	Yes	No
EnviroVeg Platinum	2011	Yes	Yes	Yes	Yes (members only)
EcoHort (production nursery)	2013	Yes	Yes	Yes	No
ePar (golf and recreation)	2013	Yes	Yes	Yes	No
Eco-Warranty (banana and mushrooms)	2010	Yes	Yes	Yes	No
Certified Land Management	2013	Yes	Yes	Yes	Yes (members only)
Cotton BMP	2013	Yes	Yes	Yes	Yes
ISO14001	2009	Yes	Yes	Yes	No
GlobalG.A.P.*	2013	Yes	Yes	Yes	No
Linking Environment And Farming (LEAF)	2012	Yes	Yes	Yes	Yes (members only)

KEY: BMP - Best management practice - industry or regional self-assessment tool

EMS - Environmental Management System - environmental certification through third party audit \*This is a quality assurance system that has an environmental component.

Table 2: Classification of existing environmental systems within relevant to horticulture.

#### A note on organic farming

Organic farming is a unique form of agriculture that avoids using synthetic chemicals, artificial fertilisers or genetically modified (GM) organisms. A focus on environmentally sustainable practices is a requirement of organic certification and standards.

Basic principles of organic farming include:

- Achieving optimum quantities of produce and food of high nutritional quality without using artificial fertilisers or synthetic chemicals;
- Preferring renewable resources and conserving energy, soil and water; and
- Minimising the use of non-renewable resources and avoiding polluting activities.

The premium for certified organic products is based on both the basic principles of organic farming and the perception in the marketplace that organic food is produced in clean environments.

The Australian Certified Organic Standard (ACOS) 2013 outlines the requirements for marketing produce as certified organic in Australia. Its scope includes organic production, manufacturing, processing and retailing. While the standard describes parameters for compliance, it does not override legislative requirements but differentiates organic agricultural practices from traditional farming practices.

The ACOS considers all other primary production or food preparation that does not conform to the ACOS standard as 'conventional' production. These guidelines would therefore be seen to cover 'conventional' horticulture production.

For more information on organic farming methods visit the Australian Organics website http://austorganic.com.

#### Catchment scale - Natural Resource Management (NRM)

Natural Resource Management (NRM) is the way in which resources like soil, water and vegetation are managed. It is fundamental to sustainable agricultural production. It is also sound business practice to ensure the resources we depend on for production are used efficiently and are in optimal condition.

A commercial business approach to risk management requires that future operations are free from potentially adverse environmental consequences. This need is strengthened by the trend in Australia to set targets for regional environmental outcomes - and an expectation that industry will do its bit to help achieve them. Many growers also accept a 'duty of care' to protect and enhance the environment - leaving their land in good shape for the next generation.

The Australian Government, in association with state and territory governments, has identified 54 regions covering all of Australia. Natural resource management regions are based on catchments or bioregions.

For natural resource management to be successful, it is important that management activities are coordinated between growers and other natural resource managers in their regions, such as Landcare groups, NRM groups and catchment management authorities. This gives the region a greater likelihood of achieving more wholistic natural resource goals and reduces the risk of individual actions being fragmented and inefficient.

NRM groups and catchment authorities may have already developed environmental objectives or targets for your region. This information is useful because it:

- Assists your activities to feed into the regional targets;
- May provide you with opportunities for financial assistance in achieving your own property goals; and
- Can provide guidance on what the local environmental issues are. For instance, if salinity has been identified as an issue in your area this may prompt you to consider in greater detail whether your property might be at risk of developing salinity problems.

Information about NRM groups/catchment authorities in your region can be obtained from http://www.nrm.gov.au/about/nrm/regions/index.html

The Horticulture NRM Strategy is an industry-wide initiative that sets out horticulture's national NRM agenda. It fits with other activities under the umbrella of 'Horticulture for Tomorrow' and provides a blueprint for future activities.

#### Development of the guidelines

The original version of the guidelines launched in 2006 were produced by Horticulture for Tomorrow – a national project supported by Australia's horticultural industries and funded by the Natural Heritage Trust, through the Australian Government's Pathways to Industry EMS Program (2004-2006).

In late 2003, the Australian Government consulted with industry and community groups on its Environmental Management Systems (EMS) policy. Following these consultations, the Australian Government announced new directions, which included the Pathways to Industry EMS Program. Through the program, the Australian Government offered to assist industry to implement an EMS or environmental assurance 'pathway' that positioned its members for the future. Horticulture Australia Limited (HAL) and the Horticulture Australia Council (HAC) accepted the invitation to lead, coordinate and manage the project to ensure whole-of-industry goals were identified and addressed in developing a pathways framework for industry environmental assurance.



12 Introduction

### Introduction



The guidelines (1st Edition) were the key output for addressing environmental assurance developed over 18 months and comprehensively trialed by 190 growers and 40 industry experts across Australia.

A series of meetings reviewed and discussed existing environmental management programs and supporting resources developed for horticultural industries in recent years. A scoping study identified the key regulatory and market drivers in Australian domestic and export markets and provided context for the guidelines.

A risk assessment approach, consistent with HACCP (Hazard Analysis Critical Control Point) and traditionally used for food safety, identified processes used in horticulture industries and the potential for significant environmental impact if process steps were not managed appropriately. Further information resources were also identified to link these guidelines with other industry-specific guidelines and programs.

The guidelines were then created by building on existing material and work in progress, and by drawing on information from a number of sources to develop a consolidated reference document based on the most up-to-date scientific and industry knowledge available at the time.

#### **Revision of guidelines in 2014**

Since the release of the guidelines in 2006, Horticulture for Tomorrow has become a benchmark for HAL due to its success in delivery, evaluation and raising awareness of the environmental issues facing Australian horticulture.

There have been advances in this area since the first edition of the guidelines were launched, including the development of industry systems (vegetables, bananas, pineapples and nursery systems), changes to Government funding programs relevant to this area (Caring for Our Country), development of the horticulture recognition framework and retailer activity (Coles support for Ausveg EnviroVeg Platinum, Freshcare Environmental v2 and GlobalGAP v4).

The across-industry committee agreed that these developments needed to be incorporated within the guidelines and consequently HAL commissioned an external consultant, Alison Kelly – Alison Kelly Consulting, to undertake the revision in late 2014.

The revision included consultation with existing system owners, industry stakeholders and relevant experts. Concurrently, the Horticulture for Tomorrow website was significantly revised. See www.horticulturefortomorrow.com.au.

#### **Continual improvement**

It is recognised that the guidelines will need to be periodically reviewed due to changing understanding of the issues, risks, technologies and management approaches. Feedback on the guidelines is welcomed and can be directed to Horticulture Australia Limited.

#### References and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www. horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

Australian Bureau of Statistics 2013 - Australian Farming in Brief (Note: Source for horticulture business statistics) http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/7106.0Main+Features42013 Australian Certified Organic (2013) Australian Certified Organic Standard, Australian Organic http://austorganic.com/wp-content/ uploads/2013/11/ACOS-2013-final.pdf

Australian Government Caring for Our Country website - NRM regions http://www.nrm.gov.au/about/nrm/ regions/index.htm

Australian Government Department of Agriculture Forestry and Fisheries - Australian Horticulture Fact Sheet (Note: Source for value and land-use of horticulture statistics) http://www.daff.gov.au/agriculture-food/food/ publications/hort-fact-sheet

Australian Government Department of Agriculture Forestry and Fisheries - National Food Plan (Note: Source for future challenges). http://www.daff.gov.au/agriculture-food/food/publications/national\_food\_plan/white-paper

CSIRO (2011) Future Australian climate scenarios, CSIRO. http://www.csiro.au/Outcomes/Climate/Climate-Change-Book.asp

Day, P. (2010) Environmental Contributions from Australian Horticulture, An information stocktake, Final Report - HAL Project Reference HG08059, Horticulture Australia Limited, Sydney, NSW. www.horticulture.com.au

Day, P. (2006) Horticulture Natural Resource Management Strategy, Horticulture Australia Limited, Sydney, NSW. www.horticulturefortomorrow.com.au

Freshcare (2011) The Freshcare Environmental Manual, 2nd Edition. Freshcare, Sydney, NSW. www.freshcare. com.au

HAL (2011) Climate change: Managing variability and carbon, Vegetable Industry Development Program, Horticulture Australia Ltd, Sydney, NSW. http://ausveg.businesscatalyst.com/rnd/fact%20sheets/Cl Change%20and%20Managing%20Carbon.pdf

Intergovernmental Panel on Climate Change http://www.ipcc.ch/

Lovell, J. (2006) Guidelines for Environmental Assurance in Australian Horticulture – First Edition, Horticulture Australia Limited, Sydney, NSW.

Muller, J; Knudsen, L; Hauser, S; Putland, D (2012) Environmental investment audit for the Australian horticulture industry, Final Report - HAL Project Reference AH11005, Horticulture Australia Limited, Sydney, NSW. www.horticulture.com.au

National Horticulture Research Network (2010) Horticulture National RD&E Framework: A plan for future investment in Australia's horticultural industries, Part of the National Primary Industries Research, Development and Extension Framework, NHRN. http://www.npirdef.org/cms\_strategy/7

NLWRA (2008) Signposts for Australian agriculture: The Australian horticulture industry. National Land and Water Resources Audit, Canberra, ACT. http://lwa.gov.au/files/products/national-land-and-water-resourcesaudit/pn21405/pn21405.pdf

Nursery and Garden Industry Australia (Second Edition 2013) EcoHort - the environmental management system for Australian nursery production, NGIA, Sydney, NSW. http://www.ngia.com.au (Source for environmental impacts of production nurseries).

Stokes, CJ and Howden, SM. (2011) Adapting agriculture to climate change, Chapter 7 of Climate Change: Science and Solutions for Australia, CSIRO. http://www.publish.csiro.au/?act=view\_file&file\_id=CSIRO\_ Chapter%207.pdf



Horticulture Australia

### Introduction



#### How to use the guidelines

#### Purpose of these guidelines

The guidelines have been developed to provide a common platform for all horticultural industries in Australia interested in implementing an environmental assurance process.

They are the national, industry-wide approach to recognising sound environmental and natural resource management in the horticulture sector. They provide a variety of suggested practices to address common environmental issues, whether the enterprise produces fresh or processing produce, cut flowers, nuts, turf or potted plants. It is recognised that the suggested practices need to be considered in combination with other, sometimes competing, environmental and business issues. Changing one component of a farm system can impact on other practices. Management is often about balancing competing resource requirements.

#### Scope of the guidelines

The guidelines are designed to provide:

- An overview of the priority environmental management issues of concern in the horticultural sector in general;
- Guidance on how a business can assess its environmental risks;
- Guidance on practices that are recommended for addressing environmental and natural resource management issues; and
- Suggestions for monitoring and recording to demonstrate that environmental management outcomes are being met.

The guidelines cover the growing, harvesting, packing, storage and dispatch stages of horticultural production. Horticultural production includes fresh fruit, vegetables, flowers, nursery products, nuts, herbs, mushrooms and turf supplied for sale to customers in the wholesale, retail, and food service sectors or for further processing across all climate zones in Australia. It does not cover the production of sprouts and minimally processed products (e.g. fresh cuts) or grapes for wine production.

In effect, we have endeavoured to produce a guide that applies to a hypothetical Australian horticultural business that grows every conceivable horticultural product. Not all issues will apply to every business. If the issue applies, you should address it; if it doesn't apply, move on to the next one.

The guidelines provide a platform for creating environmental auditing and certification options that deal with Australian environmental issues, but do not in themselves deliver certification. Businesses further down the formal EMS pathway may need to seek more site and crop specific information.

It is important to recognise that the guidelines are not a substitute for local, state or national legislative requirements. Given the breadth and variation of legislative requirements across Australia, it is strongly recommended that specific information be sought from relevant authorities to ensure compliance.

#### Who are the guidelines for?

The guidelines are targeted at horticulture enterprises with a basic understanding of production and environmental issues.

The guidelines can be used by:

- Individual businesses to implement an environmental assurance process;
- Horticultural industry groups to guide the development of environmental programs tailored to their needs on a product or regional basis; established food safety and quality assurance schemes that offer optional environmental certification choices; and
- Other stakeholders such as government, education and training, catchment management bodies, retailers, financial institutions and utilities.

Horticultural commodities that are covered includes fresh fruit, vegetables, flowers, nursery products, nuts, herbs, mushrooms and turf supplied for sale to customers in the wholesale, retail, and food service sectors or for further processing across all climate zones in Australia. The guidelines do not cover the production of sprouts and minimally processed products (e.g. fresh cuts) or grapes for wine production. For wine grape growers click here.

Note that some commodities have there own industry-specific environmental systems available, including:

- Bananas
- Nursery
- Pineapples
- Vegetables

#### For banana growers:

The Banana Best Management Practices (BMP) – Environmental Guidelines were launched at the 2013 Banana Industry Congress to demonstrate the industry's commitment to the responsible management of natural resources.

The Banana Industry BMP provides a system for growers to assess their current practices against an industry benchmark and access information to assist with practice change. Key features are comprehensive information, intuitive layout for ease of use, and access to additional resources that are only a mouse click away in the online version.

These guidelines have been developed in collaboration with the banana industry, using an industry consultative group with producers from the major production regions in QLD, NSW and WA, to ensure that the focus is on practical, regionally appropriate best management practices. Technical specialists in government agencies and the private sector have also reviewed the guidelines and provided input.

The guidelines have been designed as a valuable resource for all banana farming businesses, whether they already have Environmental Management Systems (EMS) or are assessing their environmental performance for the first time. The Guidelines reflects the structure of the Freshcare Environmental Code, which is administered by Freshcare, of which ABGC is a stakeholder. Businesses audited under other systems will also find it highly valuable. For more information on the Banana BMP Environmental Guidelines, see www.abgc.org.au.

#### For pineapple producers:

The Pineapple Best Practice Manual provides a practical reference for Queensland pineapple growers to the current 'best practices'. The manual covers all elements of how to grow pineapples with minimal environmental risk. There are 35 chapters that cover all elements of production, from site selection to product specifications.

The manual is not available online, so for more information contact: Simon Newett

Principal Extension Horticulturist Agri-Science Queensland Department of Agriculture Fisheries and Forestry Maroochy Research Facility PO Box 5083 SCMC, NAMBOUR, QLD 4560 T: (07) 54 535 926 F: (07) 54 535 901 M: 0400 56 57 84 E: simon.newett@daff.gld.gov.au

Horticulture Australia

### Introduction



#### For vegetable producers:

The EnviroVeg Program is a highly successful and industry-led environmental program developed specifically for Australian vegetable growers. EnviroVeg provides growers with guidelines and information on how to manage their business in an environmentally responsible manner.

The basic annual self-assessment program is offered free to all National Vegetable Levy paying growers throughout Australia. Once growers have submitted a minimum of one Self-Assessment and achieved a score of 85% or greater and submitted an Environmental Action Plan to AUSVEG they can then opt to move to higher levels of membership. Moving to higher levels requires the grower to subject their operation to third party audits, which require additional proof of compliance. Auditing costs are met by the grower. In return, participating growers are rewarded through increased recognition of their environmental performance and access to rewards such as use of the EnviroVeg logo in their businesses.

EnviroVeg Platinum was launched in April 2013 and is an independently assessed version of the EnviroVeg Program that offers growers access to rewards including use of the EnviroVeg logo if growers choose to submit their operations to environmental audit. The Scheme involves a number of record keeping requirements on top of what is currently required by EnviroVeg, although has been designed to work into existing Quality Assurance schemes and be easy to implement for the grower. AUSVEG has teamed up with leading retailer Coles to deliver this Scheme.

All documentation relating to the scheme and further information is available on the AUSVEG website www.ausveg.com.au or contact the AUSVEG Environment Coordinator by calling (03) 9882 0277 or emailing jordan.brooke-barnett@ausveg.com.au.

#### For wine grape growers:

Freshcare Ltd together with the Winemakers Federation of Australia (WFA) have developed the Freshcare Environmental Code – Viticulture and associated workbooks to provide wine grape growers with an industry specific environmental program to assist them in gaining certification to a recognised environmental standard. Compliance with the Freshcare Environmental Code of Practice is one of the steps required in the process of gaining EntWine accreditation.

Entwine Australia is the wine industry's national environmental assurance program. It provides Australian winemakers and wine grape growers with formal certification of their practices according to recognised international standards.

For more information visit http://www.freshcare.com.au/COPVit or http://wfa.org.au/entwineaustralia/

# HAL

#### **Guideline sections**

#### Introduction

Introductory sections provide background information about the issues and the guidelines. A table summarises the key environmental impacts associated with horticultural production, giving a useful overview of topics covered in the main part of the document

#### Assessing and managing your enterprise – Chapters 1-8

The main working sections help you assess and manage your enterprise. These sections cover risk assessment, suggested practices, monitoring and recording options, and references and further resources.

Risk assessment diagrams are provided to assess the risk of potential environmental impacts occurring and the good agricultural practices required to prevent or minimise the impact. This section provides information to help understand why the risk of impact varies. Some of the risk diagrams contain environmental indicators that are based on research, professional advice, legislative requirements and other guidelines.

List of references and resources - References that may assist producers to find additional information regarding a specific issue have been included in each chapter.

#### **Climate adaptation**

Adaptation is the ability to adapt to unavoidable climate change. Successful adaptation to climate change will require flexible, risk-based approaches that deal with future uncertainty and provide strategies that are robust enough to cope with a range of possible local climate outcomes and variations.

Due to horticulture's dependence on natural resources, especially irrigation, it is inherently vulnerable to climate change and variability. This new section outlines the potential impacts of climate change, the potential approaches for adaptation and highlights further resources that growers can access in order to consider their adaptation options.

#### Practical tools and resources

#### Legislation requirements

As laws and regulations can vary considerably between topics, states and regions this section provides advice on how to find out what laws apply to your property.

#### Process steps and inputs

Flow diagrams which detail the process steps and inputs for the major stages of field crop production, nursery production, field packing and shed packing. The diagrams show the range of steps that may occur for each process and the inputs and practices that may result in environmental impact.

#### Environmental impact identification table

The impact identification table details potential on-farm and off-farm environmental impacts from horticultural operations. The impacts are considered for each process step, making the link between activities and environmental 'hazards'. A significant environmental impact is defined as any negative change to the environment resulting from business practices that varies from the environmental outcomes acceptable to industry, the community, regulators and markets.

#### Further resources

References that may assist producers to find additional information.

#### **Review checklist**

The checklist provides a way of recording your progress through the guidelines and for identifying actions needed to address any environmental issues you uncover. By completing the checklist each year you can track your progress over time and build up

Horticulture Australia



### Introduction

#### Glossary

Acid sulphate soil - the common name given to soils containing iron sulphides.

Acidity - the strength (concentration of hydrogen [H+] ions) of an acidic substance; measured as pH. Acid substances have a pH of 1 – 7. The opposite of alkalinity.

Alkalinity - the strength (concentration of hydrogen [H+] ions) of an alkaline substance; measured as pH. Alkaline substances have a pH of 7 – 14. The opposite of acidity.

Biochar - the carbon-rich solid product resulting from the heating of biomass in an oxygen-limited environment. Due to its highly aromatic structure, biochar is chemically and biologically more stable compared with the organic matter from which it was made. Consequently it is often used to sequester carbon and improve soil fertility.

Biodiversity - the variety of life on our planet, measurable as the variety within species, between species, and the variety of ecosystems.

Bund - an embankment, wall or other structure designed to trap or contain liquids.

Carbon footprinting - a measure of how much greenhouse gas emissions specific human activities release.

Carbon sequestration - the storage of carbon that would otherwise be released into the atmosphere.

Climate adaptation - the ability of a system to adjust to climate change (including climate variability and extremes) to moderate potential damage, to take advantage of opportunities, or to cope with the consequences.

Climate change - this term is commonly used interchangeably with 'global warming' and 'the greenhouse effect' but is a more descriptive term. Climate change refers to the buildup of man-made gases in the atmosphere that trap the sun's heat, causing changes in weather patterns on a global scale. The effects include changes in rainfall patterns, sea level rise, potential droughts, habitat loss and heat stress.

Climate mitigation - involves reducing the sources of greenhouse gases and creating or enhancing carbon sinks

Constructed wetland - conversion of an area into a wetland by building dikes, small dams and/or shaping land to provide an appropriate water regime for hydrophytic vegetation.

Crop coefficient (Kc) - the proportion of water used by individual crops compared to the water that is used by a reference crop. The reference crop is a green, uniform, actively-growing crop such as grass or lucerne.

Environmental flow - water provisions needed to sustain the ecological values of our water.

Eutrophication - the enrichment of water by nitrogen or phosphorus, causing algae and higher forms of plant life to grow too fast which disturbs the balance of organisms present in water and the quality of the water concerned.

Evapotranspiration (ETo) - refers to the total loss of water from a green, uniform, actively growing reference crop such as grass or lucerne. ETo is calculated from wind speed, solar radiation, humidity and temperature.

Field capacity - refers to the soil water content after rainfall or irrigation at the point where drainage stops.

Fertigation - the application of nutrients through irrigation systems.

Greenhouse gases - gases that trap the heat of the sun in the Earth's atmosphere, producing the greenhouse effect. The two major greenhouse gases are water vapor and carbon dioxide. Other greenhouse gases include methane, ozone, chlorofluorocarbons and nitrous oxide.

Groundwater - water tha infiltrates the soil and is stored in slowly flowing reservoirs (aquifers); used loosely to refer to any water beneath the land surface.

Leaching fraction - leaching is applying irrigation water in excess of soil moisture depletion level to remove salts in the root zone. The excess water, expressed as a percentage of the applied irrigation water, is the leaching fraction

Natural Resource Management (NRM) - describes the management of our natural resources - land, soil, native vegetation, biodiversity, and water (both fresh and marine).

Nutrient - element or compound essential for animal and plant growth. Common nutrients in fertilizer include nitrogen, phosphorus and potassium

Nutrient leaching - the process by which soluble nutirents in the soil are washed into a lower layer of soil or are dissolved and carried away by water.

Readily available water (RAW) - the amount of water in the soil that is readily available to the crop. This is between field capacity and a no-stress situation.

Regulated deficit irrigation (RDI) - an irrigation strategy to manipulate vegetative growth, yield and quality with water stress

Rhizosphere - the zone of soil surrounding a plant root where the biology and chemistry of the soil are influenced by the root.

Riparian land - any land that adjoins or directly influences a body of water and includes: land immediately alongside small creeks and rivers, including the river bank itself; gullies and dips that sometimes run with water; areas surrounding lakes; and wetlands and river floodplains that interact with the river in times of flood.

Salinity - a measure of how much salt there is in water or soil.

Sedimentation - the accumulation of earthy matter (soil and mineral particles) washed into a river or other water body, normally by erosion, which settles on the bottom

Sodicity – a sodic soil has an exchangeable sodium percentage (ESP) of more than 6. This means that sodium comprises more than 6% of the total exchangeable cations in the soil.

Soil carbon - soil carbon is all the carbon found in the soil from both living things and nonliving sources such as carbonates. It is sometimes referred to as total soil carbon.

Soil organic carbon (SOC) – as measured by laboratory analysis is all soil carbon from plant and animal sources at various stages of decomposition. It does not include new plant and animal material as much of this decomposes easily, with carbon released quickly to the atmosphere as carbon dioxide. It is also known as total organic carbon and organic carbon. Soil organic carbon is around 58% of soil organic matter.

Soil erosion - the wearing away of land surface by wind or water. Erosion occurs naturally from weather or run-off, but can be intensified by land-clearing practices related to farming, residential or industrial development, road building or timber cutting.

Soil organic matter (SOM) - soil organic matter is the matter found in the soil associated with living things. It includes living organisms, fresh residues, well rotted organic matter, silica-occluded plant carbon (phytoliths), charcoal, nitrogen, sulphur, phosphorus and compounds beneficial to horticultural production and soil health in general, such as plant promotant chemicals. Soil organic matter is not tested in soil analysis, but can be calculated by multiplying the soil organic carbon test result by 1.75.

Substrate - any growing medium used in place of soil, for example potting mix.

Turbidity - a measure of water clarity or 'murkiness'. Soil particles in water increase the turbidity.

Water quality - a term used to describe the chemical, physical, and biological characteristics of water, usually in respect to its suitability for a particular purpose.

Water table - the level below which soil or rock is saturated with water.

Water use efficiency (WUE) - WUE is calculated by the amount of yield produced per megalitre (ML) of irrigation water applied. It can also be calculated by the production value in dollars per ML of irrigation water applied.

Wildlife corridors - is a link of wildlife habitat, generally native vegetation, which joins two or more larger areas of similar wildlife habitat



20 Introduction

### Introduction





# Chapter 1 Land and soil management



Objective – to minimise soil degradation and loss from the property

Farm businesses should always implement management practices that maintain or improve soil condition.

To achieve this objective, horticulturalists need to consider issues such as:

- **1a** Soil erosion caused by water;
- 1b Soil erosion caused by wind;
- 1c Soil structure;
- 1d Salinity;
- 1e Soil acidity and alkalinity; and
- 1f Sodicity.

The priorities for soil management vary depending on soil type, topography of the land, surrounding environment, previous land use and climate. The priority given to the soil management practices discussed in this section will vary from farm to farm and between production regions.

References to help determine your soil management priorities:

- For access to key sets of Australian soil information visit
- Australian Soil Resource Information System www.asris.csiro.au
- For information on soil and land resources visit

   Australian Collaborative Land Evaluation System www.clw.csiro.au/aclep/
- For climate information visit Bureau of Meteorology www.bom.gov.au

Further references and resources can be located at the end of this chapter.



# Objective – to minimise the potential for water to erode soil on the property

Soil erosion caused by water happens when water contacts exposed and/or unstable soils (soils with poor structure). Erosion can happen as a consequence of heavy rain or excess irrigation, or when drainage water from paddocks, roadways and areas around sheds and buildings moves across the land. In tree crops, shaded soil under the tree canopy can be eroded during intense rainfall events because there is little groundcover. In some regions, low-lying ground can be subjected to regular flooding. Waterways can also be subject to erosion, with negative impacts on downstream water quality (see Topic 2b – Water quality).

The likelihood of soil erosion by water and the control measures needed depend on vegetation cover, soil type and texture, soil stability (structure) and the type of horticultural activity.

Evidence of soil erosion caused by water may include:

- Rills or gullies;
- Turbid water in farm dams or leaving the property, and soil build up on fencelines or at the bottom of slopes.

To manage soil erosion caused by water, you need to identify sites on your property that are at risk, assess the level of risk and, if necessary, put in place control measures. There are a number of key strategies you can employ, including maintaining soil cover, controlling run-off water, improving soil structure and establishing sediment traps.



LOW

of the issue.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

### 1a Soil erosion caused by water

LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

**HIGH RISK** – You need to take some action. Read the Suggested Practices for that chapter. Further references and resources can be located at the end of this chapter.





#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### **Suggested practices**

#### Maintaining soil cover

Soil cover protects the soil from erosion by reducing the displacement (movement) of soil particles caused by rain or overhead irrigation droplets, and by slowing the movement of water across the site.

Types of soil cover include:

- Grassed waterways on drainage and sump areas;
- Inter-row groundcovers in orchards, vineyards and ground crops;
- Green manure/cover crops planted between (in space and time) commercial crops;
- Organic mulches, plastic, slashed inter-row material or crop residues spread over the exposed soil; and
- Products such as PAM (polyacrylamide), PVA (polyvinyl acetate) or molasses that bind soil together.

#### Managing soil cover

Control measures may include:

- Avoiding soil tillage (where possible) during times of the year when heavy rainfall events are likely, especially in tropical areas;
- Avoiding cultivation of light sandy soils subject to regular flooding;
- Using minimum tillage systems that minimise mechanical disturbance of the soil;
- Using permanent bed systems that improve soil structure and soil stability through maintaining or improving soil organic matter levels;
- Planting green manure or cover crops during the period between commercial crops to cover the soil and increase soil organic matter levels for improved soil structure, stability and fertility; undersowing or planting in the inter-row area at the same time as commercial crops;
- Leaving crop residues (where possible) on site until the site is next required;
- Minimising the time soil is left exposed between harvest and planting of the next crop; and
- Establishing permanent grass or vegetation cover on areas that are not cropped.

### Controlling run-off water

Controlling the direction of flow, volume and speed of run-off water on the site can minimise soil erosion. Long, gentle slopes are just as prone as short, steep slopes. Good planning and drainage design before planting can prevent problems later.

Control measures may include:

- Utilising the natural contour lines (natural topography) of the property, where possible; cultivating rows across the slope of the land rather than up and down the slope, when practical and safe to do so;
- Establishing cut-off drains or banks (also known as diversion banks/drains) to divert and prevent water from other areas coming on to the site;
- Establishing contour drains/moulds/bunds to collect and slow run-off from site; establishing diversion drains to control excess water flow on and around exposed sites; establishing v-drains in inter-row areas to divert water to grassed waterways and away from exposed areas;
- Establishing grassed irrigator runs and waterways to control run-off water collected by contour drains, diversion banks and roads;
- Interrupting long slopes with a cut-off drain or grassed/mulched rip lines;
- Establishing in-paddock structures such as sediment basins and sumps along drainage pathways;
- Installing and maintaining barriers such as sediment netting, filter strips or secured straw bales in water drainage channels;
- Mulching rip lines;
- Positioning access roads on ridge lines or on the contour. If possible, on relatively flat ground, construct access roads so they are higher than surrounding cultivated land;
- Ensuring all measures work with natural watercourses within and adjacent to the area being managed; and
- Considering theîlikelihood ofîexcess rain orîpotential flooding events andî managing or avoiding associated run-off when establishing new horticulturalî sites, particularly where major groundworks are concerned.

#### Improving soil structure

Organic matter is an essential component of a healthy soil because it increases the soil's nutrient and water holding capacity, improves the soil structure and provides a food source for soil organisms. Adding organic matter increases soil resistance to erosion.

Organic matter can be increased with practices such as:

- Applying compost or manures or;
- Leaving organic matter on the soil surface as mulch;
- Growing fallow crops between crop rotations or interrows;
- Applying organic by-products, such as greenwaste;
- Reducing cultivation;
- Avoiding high rates of nitrogen fertilizer; and
- Encouraging earthworm activity, which incorporates organic material deeper into the soil.



Further references and

the end of this chapter

resources can be located at

Horticulture Australia

### 1a Soil erosion caused by water

Further references and resources can be located at the end of this chapter



#### Establishing sediment traps

Sediment traps or ponds (also called silt traps or ponds/sediment retention basins) aim to hold run-off water long enough to allow soil particles to settle. They can be small ponds or weirs, or large dams that capture and re-use run-off water. Artificially constructed wetland systems may be established to capture sediment and remove the nutrient in run-off waters.

Filter strips are areas of vegetation that catch sediment and nutrients in water that are flowing to a watercourse. Generally they run alongside watercourses or across a depression. They are not effective if the water is deep enough to flatten the vegetation and is not slowed down. Slopes of more than 10% are unsuitable for filter strips as the water moves across the ground too rapidly for sediment to be caught by vegetation.



#### Monitoring and recording

Erosion caused by water can be monitored by:

- Visual inspection;
- Assessing water turbidity; and
- Assessing soil erosion losses.

#### Visual inspection

Immediately after a rainfall event, look at how run-off is flowing across the farm. Is erosion occurring? How dirty (turbid) is the water?

Inspect the property for signs of scouring (drainage lines, channels) or for silt accumulation around plants or other obstructions.

Photographs can be useful to record problem areas (e.g. drainage lines, rills, gullies, prone slopes) before and after control measures are implemented.

#### Assessing water turbidity

Turbidity is a measure of water clarity or 'murkiness'. Soil particles in water increase the turbidity.

In addition to a visual inspection of water leaving the property or returning to farm dams, a turbidity tube can be made and used to gauge basic changes in water turbidity. Turbidity meters are also available for more precise assessments.

#### Turbidity tube

A turbidity tube is a length of clear pipe with a clear bottom. The general idea is to determine the depth at which you can no longer see through the water. This is an indicator of turbidity.

To measure turbidity:

- Collect a water sample in a clean bucket without disturbing sediment from the bottom of the dam or stream;
- Assess the water sample as soon as possible after collecting;
- Place the turbidity tube on a white piece of paper or card that has a cross or other mark on it;
- Shake water sample and pour into the tube until the cross or mark on the card can no longer be seen when viewed from the top (i.e. looking down through the water);
- Record the height of the water in the tube;
- The lower the height of water, the greater the turbidity;
- This may indicate there is a large amount of sediment in your farm run-off and action may be required to stabilise soils or reduce run-off.

The tube and card need to be stored to prevent the tube from getting scratched and the mark on the card from fading.

#### Assessing soil erosion losses

Place a piece of  $100 \times 50$  mm timber, or similar, on the ground and, over time, look at the amount of soil that accumulates behind it.

Pegs with depth markings can be placed in silt traps to measure the amount of accumulated silt. Paddock records can also be useful to demonstrate groundcover/ cropping history during times when high rainfall is usually expected.



#### **References and resources**

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter



Horticulture Australia

### 1a Soil erosion caused by water

Further references and resources can be located at the end of this chapter



### 1b Soil erosion caused by wind



Objective - to minimise the potential for wind to erode soil on the property

Soil erosion caused by wind happens when wind contacts exposed (uncovered) and unstable soils (soils with weak structure) at speeds that can physically move soil particles. Minimising the area of exposed soil and reducing the wind speed are the keys to minimising soil erosion. Once wind erosion starts, it is hard to control and repair. Prevention is best.

Evidence of soil erosion caused by wind may include:

- Dust:
- Rills or gullies on light or sandy soils;
- Exposed subsoil and rocks (rocks appear to be 'rising to the top' of the paddock);
- Exposed roots of trees and shrubs (long-lived vegetation); and
- Soil and/or organic matter (such as twigs and grass) building up against the side of fences or hedges.

To manage soil erosion caused by wind, identify the sites on your property that are at risk, assess the level of risk and put control measures in place if needed. There are a number of key strategies you can employ, including maintaining soil cover, controlling wind speed and improving soil structure.





#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



Suggested practices

#### Maintaining soil cover

Soil cover protects the soil from erosion by minimising soil exposure to the physical force of the wind. Types of soil covers include:

- Grass or native vegetation on areas that are not cultivated;
- Inter-row groundcovers in orchards, vineyards and ground crops;
- Under-tree groundcovers and mulchers in orchards;
- Green manure/cover crops planted between commercial crops; and
- Organic mulches, plastic, slashed inter-row material or crop residues spread over the exposed soil.

#### Managing soil cover

Control measures may include:

- Avoiding soil tillage (where possible) during times of the year when high winds are likely; planting green manure or cover crops during the period between commercial crops to cover the soil and increase soil organic matter levels for improved soil structure, stability and fertility; undersowing or planting between rows at the same time as commercial crops;
- Using cover crops for germinating seedlings;
- Leaving crop residues (where possible) on site until the site is next required;
- Minimising the time soil is left exposed between harvest and planting of the next crop; and
- Establishing permanent grass or vegetation cover on areas that are not cropped.

#### Moderating wind speed

Controlling wind speed on the site can minimise soil erosion.

Constructing or planting a shelterbelt/windbreak will slow the velocity of wind across a site (shelterbelts/windbreaks should be designed to allow 30-50% of the wind to pass through). The protective effects from a shelterbelt/windbreak reduce with distance away from it (protection extends no more than 20 times the height of the shelterbelt/windbreak). Vegetation shelterbelts/windbreaks also provide wildlife habitat, assist in minimising spray drift and reduce the visual and noise impacts of site activity and can have an influence in

high water tables.

Horticulture Australia

Further references and

the end of this chapter

resources can be located at

### 1b Soil erosion caused by wind

Further references and resources can be located at the end of this chapter



#### Improving soil structure

The wind will not pick up soil particles greater than about 0.5 mm. To control wind erosion keep soil aggregates greater than this size. This is relatively easy for soil heavier in texture than loamy sands. However with sands, especially water repellent sands, this is not possible and adequate ground covers are required.

Plenty of organic matter in the soil will strengthen soil structure and make it less prone to wind erosion.

Strategies to improve soil structure may include:

- Using minimum-tillage systems that minimise mechanical disturbance of the soil and improve soil;
- Structure and soil stability by maintaining higher soil organic matter levels;
- Using permanent bed systems that improve soil structure and soil stability by maintaining soil organic matter levels; and
- Incorporating organic matter into the soil to build up soil organic matter levels.

#### Other management strategies

Irrigation can be applied immediately prior to, or during, wind events to increase the cohesion between soil particles, thereby reducing erosion.

Cultivating so as to leave a rough, raised and very uneven surface can also reduce erosion.

Planning when setting up new sites, particularly where major ground works are concerned, should include consideration of the likelihood of wind extremes and managing or avoiding the periods when they are likely to occur. Using remnant vegetation or shelter belts within or adjacent to the new site can minimise soil erosion.



Monitoring and recording

Erosion caused by wind can be monitored by:

- Visual inspection, or;
- Assessing soil erosion losses.

#### Visual inspection

Wind erosion can be visually assessed - have a look at an exposed site with light soils on a windy day! However, the effects of erosion are often subtle and require an extended period of time to become obvious. In this case it may not be possible to clearly distinguish between the causes of erosion, but an understanding of your own property, soil type and weather patterns should help you determine the most significant influences so that appropriate control measures can be instigated.

Visual signs of erosion include:

- Rills or gullies;
- Exposed subsoil;
- Exposed rocks (rocks appear to be 'rising to the top' of the paddock);
- Exposed roots;
- Piles of organic matter such as twigs and grass forming 'debris dams'; and soil and organic matter caught in or building up against sides of fences.

Use of satellite imagery (to assist in mapping eroded areas), GPS (to pinpoint exact locations for measurement) and photography are good ways to more accurately record changes over time.

#### Assessing soil erosion losses

Measuring wind erosion can be difficult because of its patchy nature.

#### Natural benchmarks

For longer-term monitoring in some situations, use natural benchmarks such as big rocks or trees - mark the soil height now, and then check the soil height over time to see if it changes.

Be careful when selecting a natural benchmark (for instance rocks may be moved by livestock or cultivation).

Alternatively, choose a natural benchmark that has a 'soil mark' and measure the distance between this soil mark and the current soil level to gauge the erosion that has taken place up until now.

#### **Erosion** pin

An erosion pin is a metal bar driven into the ground with a portion protruding for a known height (e.g. 5 cm). Monitor the distance between the top of the pin and the soil surface over time.

#### **Erosion pipe**

An erosion pipe is like an erosion pin except that it contains soil that will not be affected by erosion. Monitor the distance between the soil height in the pipe and that surrounding it.

Be careful where you site the pin or pipe so that results are not affected by ploughing or other soil cultivation activities.

#### DustTrak technology

DustTrak devices use laser technology to measure Atmospheric Particulate Concentration: the concentration of dust particles in the air. The data obtained with this device is used to compare erosion events within and between years.

Further references and resources can be located at the end of this chapter



Horticulture Australia



**References and resources** 

For access to relevant references and further resources click here.

### 1b Soil erosion caused by wind

Further references and resources can be located at the end of this chapter



### **1c Soil structure**



Objective - soil structure is suitable for root growth, water infiltration, aeration and drainage needs of the crop

Deep well-structured soils grow the best crops. A well-structured soil has pores, channels and spaces between aggregates (clumps). Water can drain quickly, roots go through the soil easily and there is no hard crust on drying.

Degraded soil has a high proportion of small particles with few water stable aggregates. The reduction of pore size and continuity results in massive blocks that restrict root growth and plant productivity. Compacted soil requires more cultivation to prepare a seedbed and this additional cultivation causes further deterioration in soil structure.

Crop yields increase dramatically when soil improves. In Australian horticulture, crops grown on the poor soil types average 10t/ha and those grown on the best soils can achieve yields of 50t/ha. The cause of low productivity in irrigated agriculture lies in coalescence, a soil hardening process, and consequent low root activity, which also restricts what happens above the ground. The best soils overseas remain loose, soft and porous even after centuries of growing crops.

To maintain and improve good soil structure you should establish an appropriate crop rotation, increase organic matter in the soil and follow good tillage practices.

Risk assessment





#### Review checklist

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

#### Cultivation method

Most tillage for fruit and vegetable crops occurs prior to planting to enable suitable contact between the soil and the planted material. This primary tillage is an important part of initial land preparation and cannot really be avoided. Secondary tillage operations should be minimised where possible.

Machinery can cause compaction, so the following points should be considered:

- Minimise traffic in the paddock;
- Keep trucks to headlands;
- Use low weight spray rigs;
- Consider the effects of axle load, tyre width and inflation pressure, aiming to maximise weight distribution; and
- Make contractors aware of compaction issues.

Consider using a minimum-tillage approach such as tramlining or semi-permanent beds.

When choosing cultivation, consider the crop stage, soil moisture and soil condition. Rotary hoes are generally used to pulverise the soil for primary tillage. Along with disc cultivators, they should be used as sparingly as possible as their use leads to reduced soil organic carbon and soil compaction. Tyned and non-inverting implements are kinder on the soil structure.

Blunt tools can also add to compaction. Use sharp and correctly adjusted tools to till the soil. Use implements that mainly apply an upward force to the soil.

Minimise the number of soil workings.

Avoid overworking with powered implements. Consider using semi-permanent beds.

#### Cultivation timing

The soil moisture content during tillage has an important effect on soil structure. Where the water content is too great, the soil acts like plasticine, smearing and compacting with tillage and traffic. Don't go onto paddocks with machinery when the soil is wet. Similarly, soils can be too dry to work, requiring excessive amounts of energy to produce a seed bed.

Horticulture Australia

Further references and

the end of this chapter

resources can be located at

### 1c Soil structure

Further references and resources can be located at the end of this chapter



#### Cultivation timing cont.

Ideal moisture levels depend on soil type and texture. You can check by working some soil from the plough layer in your hands. If the soil is too wet it will work like plasticine, if it is too dry it will be hard to work and tend to shatter to dust.

#### **Remedial action**

If a hard pan or compaction layer is present, then additional cultivation may be needed depending on whether the cause is cultural or due to sodicity (see Topic 1f – Sodicity). If the condition is not due to sodicity, cross-ripping under the correct soil moisture levels will help to shatter the pan, loosening and breaking clods that will break down further when exposed to the weather.

Deep ripping needs to be done early enough to allow weathering, or else try to leave your deepest working to last in the soil preparation sequence, because after ripping the soil is highly susceptible to recompaction.

The benefits of deep ripping can be short term (~1 year) unless actively growing roots enter the fracture lines.

A sodic soil has an exchangeable sodium percentage (ESP) of more than 6. This means that sodium comprises more than 6% of the total exchangeable cations in the soil.

Soils with shallow sodic subsoils should not be ripped. This can bring sodic soils to the surface and create problems with surface crusting (see Topic 1f – Sodicity).

#### Increasing organic matter

The terms soil carbon, soil organic carbon and soil organic matter are often used interchangeably, but have distinct meanings. It is important to understand the differences. See glossary. Soil carbon is dealt with below in organic amendments.

Soil organic matter refers to the matter found in the soil associated with living things, such as living organisms, fresh residues, well rotted organic matter, silica-occluded plant carbon (phytoliths), charcoal, nitrogen, sulphur, phosphorus and compounds beneficial to horticultural production and soil health in general, such as plant promotant chemicals.

Increasing organic matter through use of crop rotations, such as rye grass, and green manure crops promotes good soil structure. Stubbles and crop residues can also be returned to the soil.

#### **Ryegrass in orchards**

In south eastern Australia it is important that fruit growers encourage winter grass in their orchards. Grass /grassroots:

- Adds organic matter to the soil;
- Opens the surface of the soil to let irrigation water in;
- Keep the soil loose, soft and porous;
- Penetrate into the subsoil, producing pores and increasing stability;
- Helps dry the soil in wet winters; and
- Forms a mulch in summer to reduce high soil temperatures.

Grass is much better at doing these things than clovers or weeds. Rye grass is the most effective of all the grasses. Ryegrass is suited to the climate of this area and in fact is a weed here. It will self seed each year if managed properly and will grow well. It has been demonstrated that rye grass develops a rhizosheath of soil particles adhering to each rye grass root. The soil within the rhizosheath builds up organic matter, arising from root exudates, root hairs and microbial organisms, all in very large numbers. These eventually become organic matter. An important property of the rhizosheath is that organic matter within it is protected from oxidation. Under rye grass, organic matter quickly builds to 8%, the level required to prevent soil coalescence. Fruitgrowers should note that full potential for increasing yield and quality relies on tree management as well as soil management. Tree management on soil that has become more responsive after treatment, may require changes to such practices as pruning and reduction of the leaf to fruit ratio.

#### Building soil organic carbon

The organic carbon content of soil is defined by the balance between inputs of carbonrich material (plant growth and additional material) and losses through decomposition, erosion and product removal. Where inputs are greater than losses, soil organic carbon increases.

Table 1 below summarises management practices that increase soil organic carbon by providing carbon inputs or decreasing carbon losses from the soil. The first seven management activities listed are appropriate and feasible for horticultural systems.

Table 1: Management practices to increase soil organic carbon. Source: Cox et al (2012).

Activity	Provides C inputs	Reduces C losses
1. Increase plant (biomass) production by applying sufficient nutrient and water	~	
2. Retain stubble/crop residue	<ul> <li>Image: A second s</li></ul>	<ul> <li>Image: A second s</li></ul>
3. Reduce fallow periods	<ul> <li>Image: A set of the set of the</li></ul>	<b>v</b>
<ol> <li>Include opportunity crop/rotations/ manure crops green</li> </ol>	<ul> <li></li> </ul>	<b>V</b>
5. Apply high carbon amendments such as compost, biochar, some manures	<ul> <li></li> </ul>	
6. Reduce erosion		<ul> <li>Image: A second s</li></ul>
7. Reduce cultivation		<ul> <li>Image: A second s</li></ul>
8. Introduce farm forestry	<ul> <li>Image: A second s</li></ul>	
9. Improve pasture management	<ul> <li>✓</li> </ul>	<ul> <li></li> </ul>





Horticulture Australia

### **1c Soil structure**

Further references and resources can be located at the end of this chapter.



Carbon-rich soil amendments such as mulch, compost and biochar offer horticulture the benefit of improved soil condition and potential for carbon trading for the most stable forms. See Energy & Greenhouse Gas Management Chapter.

Biochar application to soil, for example, has been shown to:

- Alter chemical functionality such as soil Cation Exchange Capacity (CEC), pH and nutrient availability;
- Improve the physical properties of the soil, particularly aggregation, water retention, water use efficiency, and reduce tensile strength in hard setting soils;
- Modify the biological functionality by providing a habitat for microorganisms due to its highly porous nature or by altering substrate availability and enzyme activities on or around biochar particles;
- Suppress some soil-borne diseases; and
- Increase yields (when used in conjunction with fertilisers).

Current methods of biochar incorporation use surface application, then mechanical incorporation into the topsoil, a method suitable for most annual and semi-permanent orchard crops. Alternative incorporation methods need to be developed to introduce biochar into permanent perennial horticultural crops without damaging existing root systems. Possible methods include coring using modified turf aerators, and combining biochar with a mulch material for surface application.

However, it is difficult to generalise the impact of biochar on soil properties due to this range of biochar production variables and biochar's complex interactions with soil organisms, chemical elements and physical structure. Considerable uncertainty still surrounds the use of biochar in farming systems given the range of production processes, the types of biochar, and the variety of soil types in horticulture.

Adding carbon-rich amendments to soils is becoming increasingly common in horticulture, but is only really effective when the amendments are selected for a purpose. Growers need to be clear whether they are addressing a specific soil constraint (e.g. lack of nutrients), or want to build soil organic carbon. Organic amendments unsuited to a particular soil, site or production type may have detrimental effects on the soil or environment.

For more information on biochar see HAL/NSW DPI publication on biochar in horticulture (2012) http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0008/447857/DPI-BioChar-in-Horticulture.pdf.

#### **Crop rotation**

Using rotations and green manure crops will provide short-term soil structure benefits through better soil aggregation. This helps optimise the soil's water-holding capacity, ability to hold nutrients, workability and water infiltration.

Rotating crops also assists soil structure, with crops such as grasses and legumes increasing the spaces or pores through the soil. Deep-rooted crops can also recycle excess soluble nutrients like nitrate and sulphur from deeper in the soil and these crops add organic matter as the deep roots eventually break down. Roots help break up the soil and create pores to assist with movement of water through the soil.

Monitoring and recording

Soil compaction can be assessed by determining how difficult it is to dig. The assessment should bear in mind any short-term tillage and effects of soil moisture.

#### Spade test

The following scale can be used:

Hand	Soil that can be dug easily by hand h sandy. Maintain and improve structur matter. Use minimum tillage for all
Spade	Soil that can be dug easily using a sp structure.
Standing on spade	If you have to stand on the spade, the have high clay content. Aim to break improved drainage.

#### Record the result of spade test.

#### Penetrometer (screwdriver) test

A simple test of compaction is to see how far you can push a screwdriver into the soil using reasonable hand pressure. It is a way of simulating the difficulty that roots have pushing through the soil. Try it after decent rainfall or irrigation.

#### Visual assessment

Soil compaction affects the ability of plant roots to penetrate the soil and root systems are often stunted. Dig up some plants and assess their root systems and also assess the overall vigour of the plants. Stunted or sharply bent roots mean small, feeble, low-yielding plants that are prone to drought. It can be useful to compare roots from different areas, such as under fencelines where compaction may be less.

Take a closer look at the clods and aggregates. Many large clods mean the soil will need to be kept wetter to allow roots to penetrate. Sharp angular aggregates with smooth faces indicate poor structure. Well-structured soils have a range of aggregate sizes (2 -10 mm), with irregular or rounded shapes and porous faces.

Look for areas where water ponds. Ponding is a way of measuring compaction and soil structure. Water lying around in the paddock means that there are few soil pores in and below the plough level. Soil compaction is one cause of this.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

### **1c Soil structure**

has a poor structure or is very ire by increasing organic l crops.

pade has good or very good

ne soil may be compacted or < up compacted areas and

a very high clay content. nprove the drainage. rops/pasture.

Further references and resources can be located at the end of this chapter.



### **1c Soil structure**

#### Soil test

Organic matter content can be included on regular soil tests. Natural levels of organic matter in the soil depend on factors such as climate, site drainage and soil texture. A heavy soil will generally have higher levels of organic matter than light, sandy soils.

Measured as organic carbon content, an approximate guide is as follows:

- Very low (below 1%)
- Low (1 2%)
- Satisfactory (2-4%)
- High (above 4%)



#### References and resources

For access to relevant references and further resources click here.



#### Objectives

- horticultural activities are managed to ensure soil or water salinity problems are not created or exacerbated
- horticultural production does not contribute to local, catchment or regional salinity problems

Salinity refers to the presence of soluble salts in soil or water. For the irrigator, there are 'good' salts and 'bad' salts. The good salts are the nutrients like nitrate, potassium and calcium because plants need to take up large quantities for growth. Bad salts include sodium and chloride; they are not required for plant growth and in some circumstances can build up to high levels in the soil and harm plant growth.

These salts may be naturally occurring, coming from the parent material from which the soil was formed. Other sources of salt can be rainfall, overuse of mineral fertilisers or poultry manure, or the use of saline irrigation water. Saline irrigation water may result from salts percolating out of naturally salty soil into waterways or groundwater, or from seawater intrusion into coastal groundwater. Seawater intrusion is usually a result of excessive groundwater drawdown from irrigation, or lack of groundwater recharge due to drought.

Primary salinity is naturally occurring, while secondary salinity is the result of human activity. In the context of these guidelines, salinity should be taken to mean secondary salinity.

Salinity can dramatically reduce agricultural productivity, as high salt levels can limit crop growth and even kill plants. Salinity makes it more difficult for plants to extract water from the soil. Salinity also has impacts beyond agriculture as it can affect infrastructure such as roads and buildings. Salinity reduces the diversity of native plants and animals and is linked to environmental degradation such as soil erosion, deteriorating water quality in streams, rivers and groundwater and loss of riparian vegetation.

The development and progress of salinity tends to be highly complex. Water table levels and potential salinity problems and discharge points may vary considerably. This will depend on the site conditions, groundwater processes and land management practices.

Rising water tables are one of the main causes of salinity. Crop and annual pasture plants use less water than perennial native vegetation, therefore allowing more water to travel down past the root zone and into the groundwater beneath the surface. This extra water makes the water table rise. As the water rises it dissolves the salts that are naturally in the soil, so the rising water becomes salty, contaminating the land and surface water. Rising water tables can also bring salt into the root zone, which may not be leached out of the soils by rainfall or irrigation. Evaporation from water tables within two metres of the soil surface also causes salt accumulation in the root zone and can dramatically affect plant productivity.

Undulating landscapes tend to have specific groundwater discharge points, resulting in discrete areas of salinity that can vary in severity and area. In contrast, rising water tables in flat landscapes tend to affect larger areas and salinity can be a more regional issue.

More recent studies indicate that the dryland salinity occurrences are closely related to naturally occurring salinity. With the rising water tables after the higher rainfall, salt is again being expressed on the surface in high-risk salinity catchments. Consequently, the expression of dryland salinity in the landscape is cyclical – related to rainfall.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

### 1d Salinity

d to ensure soil or eated or exacerbated contribute to local, blems

Further references and resources can be located at the end of this chapter.



### 1d Salinity

#### Salinity cont.

Salt movement and salinity outbreaks in landscapes dominated by sandy soils tend to be rapid (<10 years) and severe compared to landscapes with many metres of subsurface clays. Salt movement in clay landscapes is buffered by the clay and may take decades, or even centuries, to flow through the system.

To manage salinity it is important to understand whether it is caused by rising groundwater, irrigating with saline water or saline soils. Sites on a property that are at risk need to be identified, the level of risk assessed and control measures put in place if needed.

There are a number of key strategies in relation to managing salinity including:

- Careful site selection;
- Understanding the source/cause of salinity;
- Monitoring salt levels in irrigation water and adjusting irrigation strategies where
- necessary; and
- Minimising rising water tables through appropriate drainage/use of vegetation.

R

**Risk assessment** 

### GROUNDWATER AND SOIL SALINITY





Further references and resources can be located at the end of this chapter.



Horticulture Australia

### 1d Salinity

Further references and resources can be located at the end of this chapter.



### 1d Salinity



#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

Always avoid salinity problems in preference to attempting rehabilitation. Key considerations in avoiding salinity include choosing the right site and then employing good soil/water management strategies, combined with diligent monitoring.

#### Site selection

When choosing sites, consider the likelihood of salinity problems. Undulating landscapes tend to have specific groundwater discharge points, resulting in discrete areas of salinity that can vary in severity and area. In contrast, rising water tables in flat landscapes tend to affect larger areas and salinity can be a more regional issue.

Check for visual indicators of salinity problems, such as:

- Salt-tolerant vegetation; and
- Bare soil or salt scalds.

A soil survey or Electro Magnetic (EM) survey can help identify, assess and help manage saline soils and water.

#### Understanding the source

If salinity problems exist or are suspected, it is important to understand the source of the problem. Salinity can be due to saline irrigation water, groundwater salinity or soil salinity.

#### Irrigation water salinity

Salts dissolved in water can be easily measured by testing for electrical conductivity (EC). A small, hand- held meter is invaluable for checking the salinity of irrigation water and monitoring changes through the season. If irrigation water exceeds an EC of 0.8 dS/m (this is equivalent to 500 ppm or 500 kg of salt in 1 megalitre of water) a full chemical analysis, interpreted by a technical expert, should be undertaken and professional irrigation management advice sought.

If salinity is present, depending on the actual EC level and the soil type, consider implementing the following:

- Irrigate at night to avoid high evaporation rates which cause salts to concentrate;
- Avoid leaf contact (e.g. Use drip and not overhead sprinkler);
- Maintain low soil moisture deficit (making it easier to flush salts from the root zone);

A megalitre of irrigation water that has a salinity level of 1 dS/m will deposit 640 kg of salt.

dS/m stands for deci-Siemens per metre and is a measure of electrical conductivity, and therefore a measure of salinity

- Ensure good subsurface drainage;
- Shandy/dilute saline water with less salty supplies; and/or
- Consider growing salt-tolerant crops.

These practices will usually only be beneficial for a short time. It is important to consult a specialist for more site-specific assistance.

#### Some salinity benchmarks (in dS/m)

Distilled water = 0Absolute limit for people = 2.5Desirable limit for people = 0.83Limit for mixing herbicides = 4.7Tastes salty (depending on the ions present) = 1.7 Seawater = 55+

Table 1.2: Tolerance of plants to salinity in irrigation water. Source: Australian and New Zealand Guidelines for Fresh and Marine Water Quality - Volume 1: Paper 4 - Primary Industries (2000)

Сгор	EC **(dS/m) threshold for crops gro		
	sand	loam	
Apple	2.0	1.2	
Avocado	2.3	1.3	
Beans	1.9	1.0	
Beet	6.5	3.7	
Broccolli	4.9	2.8	
Carrot	2.2	1.2	
Cucumber	4.2	2.4	
Eggplant	3.2	1.8	
Grape	3.3	1.9	
Grapefruit	3.0	1.7	
Lettuce	2.7	1.5	
Olive	5.0	2.9	
Onion	2.3	1.3	
Orange	2.9	1.7	
Pea	3.2	1.8	
Peach	4.7	2.7	
Pepper	2.8	1.6	
Potato	3.2	1.8	
Rockmelon	4.6	2.6	
Tomato	3.5	2.0	
Zucchini	7.3	4.2	

Further references and resources can be located at the end of this chapter



\*\* EC is a measure of the salinity of the irrigation water. The threshold is the salinity level above which yield decline is likely to occur. The threshold is lower on soils with higher clay content because plants have to work harder to extract water from these soils to start with. Saline irrigation water makes it harder again.

Guidance is available on crop tolerance for water salinity, however this is highly dependent on soil types and the degree of associated waterlogging. Expert advice should be sought.

### 1d Salinity

Further references and resources can be located at the end of this chapter



#### **Groundwater salinity**

To minimise the chances of salinity problems, the water table should be kept two metres or more below the soil surface. In some areas this is an issue that requires regional management, such as establishing spear points, tile drains or groundwater pumps to increase the depth of the water table.

The water table should also be monitored over time to check if it is stable or rising. If the water table is high (within 2 m) then artificial subsurface drainage may be needed. If water tables are not yet high but are rising, subsurface drainage may be needed in the future. Also work on improving irrigation efficiency.

#### Soil salinity

Salinity levels in the soil are measured in a 1:5 soil solution (1 part soil to 5 parts water). These are called EC 1:5 readings. A reading of <0.2 dS/m is usually safe for horticultural crops. The heavier the soil texture (e.g. clay) the more sensitive the crop is. Thus EC 1:5 readings need to be adjusted for soil texture to reflect how happy the plant actually feels in the soil. The adjusted measurements are called 'saturation extract' or EC. For major horticultural projects, it is recommended that EC values are measured directly rather than the approximate conversions. A few laboratories in Australia provide this service.

In some cases, applying a 'leaching fraction' when irrigating may be necessary to flush salts through the soil profile. However for leaching to be effective, good drainage is needed and a plan for safe disposal of the saline drainage water must also be in place. Consider growing or changing to salt tolerant crops or varieties. Further advice should be sought from a specialist.

Sometimes high soil salinity can be found when the water table is low and the salinity of the irrigation water is also low. In these instances check the leaching efficiency of the irrigation system. Also check your fertiliser program – certain fertilisers such as muriate of potash can have a strong influence on soil salinity.

High salinity can be found on the edge of wetted areas or in other dry spots. Also check for poor drainage of soil and seek expert advice. Careful management of the soil chemistry is needed when saline, sodic soils are drained.

#### Soil salinity tolerance of horticultural crops:

Table 1.3 is a list of thresholds expressed as the electrical conductivity of soil water (ECsw) for maximum production of major horticultural crops, and likely yield reductions from higher salinity levels. These values were calculated using the relationship between soil saturated paste electrical conductivity (ECe) and the salinity by the suction cup soil water extractor developed by SARDI. The SARDI suction cup water extractor is now marketed under the brand name of SoluSAMPLER by Sentek Sensor Technologies (www.sentek.com.au).

http://www.environment. gov.au/resource/ australian-and-newzealand-guidelinesfresh-and-marinewater-quality-volume-1guidelines

Сгор	Threshold for maximun production <sup>2&amp;3</sup> (ds/m)	Threshold for reduced yield levels (ds/m)	
	100% yield	75% yield	
Orange	3.4	6.6	
Grapefruit	3.4	6.6	
Lemon	3.4	6.6	
Apricot	3.2	5.2	
Peach	3.4	5.8	
Carrot	2.0	5.8	
Onion	2.4	5.6	
Potato	3.4	7.6	
Tomato	5.0	10.0	

NOTE: These values are a guide only and can vary with soil type, leaching potential, irrigation method and plant age.

The term leaching fraction refers to the application of water to the point where water percolates down a soil profile.

Table 1.3: Horticultural crop thresholds. Source: Biswas et al. (2009).

#### Irrigation management

In areas affected - or at risk of being affected - by salinity irrigation requires careful management. It is a good idea to seek professional advice before developing an irrigation system in these situations.

Applying a leaching fraction can flush salts out of the topsoil. Rainfall may act as a leaching fraction. However, excessive leaching fractions can worsen the process of salinisation by causing the water table to rise, so they need to be carefully managed.

Monitoring the level of soil moisture provides a means of checking water table depth as well as efficiency of irrigation (see Topic 2a - Irrigation efficiency).

#### Improving drainage

Improve drainage in saline areas, particularly if salinity problems are associated with a rising water table and saline groundwater. If soils are waterlogged, removing excess water can help leach of salt from the root zone to lower levels in the soil profile. Consideration must be given to management of the drainage water.

Cut-off drains can divert and remove surface water that would otherwise become groundwater recharge. Surface drains should be stabilised with fencing and vegetation cover.

Raised beds with adjoining furrow drains can improve surface drainage and salt leaching. Sub-surface drainage can reduce waterlogging and increase the leaching of salt. Care is needed when considering drainage options as drains in dispersive soils can lead to soil instability and severe erosion.





Horticulture Australia

## 1d Salinitv

#### Further references and resources can be located at the end of this chapter



#### Vegetation cover

Vegetation can assist in preventing and managing salinity, particularly salinity associated with rising water tables.

Deep-rooted plants can assist in preventing rising water tables, by utilising water deep in the soil profile.

Maintaining vigorous plants will help use rainfall, preventing excess water soaking through the soil surface.

If salinity problems already exist, salt-tolerant tree species can be established to assist with water utilisation and gradual lowering of the water table.



#### Monitoring and recording

If you are beginning a new project or irrigating a new area, undertake the required investigation to determine the risk of salinity problems arising or to find existing problems.

If salinity problems exist or there is a risk of them developing, a monitoring program should be developed based on a salinity survey and specialist advice. The program needs to be tailored to suit the specific site.

From then on it is a matter of monitoring in areas where salinity issues occur. It is recommended you measure and record:

- Soil salinity (electrical conductivity, chemical composition of the salts and em survey);
- Trends in the salinity of irrigation water and groundwater;
- Trends in the depth of the water table;
- Salinity (sodium and chloride) in leaf samples; and
- Photographic points to track changes in scald area and severity.

A low-cost approach to monitoring for salinity is detailed over the following pages. This may not be appropriate for your situation, particularly if dealing with high-value perennial crops where a more detailed analysis of the salinity risk would be warranted.

#### Monitoring the water table

Equipment needed – a small 50 mm drilling rig (borrow or hire), a few lengths of 40 mm diameter PVC pipe and a bundle of steel posts.

#### Select suitable sites for monitoring stations

If you don't have any visible sign of salt-affected land, choose a site that is convenient for monitoring (i.e. not in the middle of land you will be regularly working up) and that could be at risk of salinity.

If you have a salt scald, choose a monitoring station where the boundary is suspected of spreading, and if possible where the boundary is easy to identify. Two or three monitoring stations for each area of salt-affected land should be sufficient. For each monitoring site, mark the boundary of the affected area with steel posts 10-20 m apart, exactly on the salt boundary, so that you will easily see any change in the boundary over time.

#### Establish the monitoring stations

At each monitoring station, drill holes (about 50 mm in diameter) with a hand or power auger to a depth of 2 m. If you have salt-affected land, locate the holes across the salinity boundary so that at least one hole is in the unaffected area, one is in the obviously affected area and one is on the boundary.

Use 3 m lengths of 40 mm PVC pipe (Class 6) to line the holes. Cut slots in the bottom metre with a fine-bladed hacksaw. When the pipe is in the hole, pack clay or cement grout around it at ground level to prevent inflow of rainwater. Keep rain out of the top with an upturned jam tin or old bucket. Identify the hole by a number on the top of the pipe. Prevent stock rubbing against the pipe by driving a steel post into the ground next to it or cutting the pipe off short, 20-30 cm above ground level.

#### Determining the depth of the water table

After establishing the monitoring sites, leave the holes for at least three weeks before measuring the depth to the water table from the top of the pipe. Subtract the height of the pipe above the ground surface to calculate the depth to the water table below the ground surface. Use a metric tape measure with a small wash basin plug wired on the end to detect the water surface by sound.

If no water is detected in a 2 m deep hole, it is unlikely that the land is in immediate danger of salt encroachment. Capillary rise (wick action) of saline water to the soil surface only operates from water tables that are 2 m deep or shallower in most soil types.

If the water table is 1.5 - 2 m below ground level, the soil surface must be considered at risk. If the depth to the water table is less than 1.5 m there may well be indications of salt on the surface.

With the water table at 1 m or less, salting is practically inevitable. In the worst cases, the water table level in the pipe may rise above ground level, indicating upwards pressure in the water table.

#### Soil profile sampling

Take soil samples from near each hole during dry months. Sampling should be conducted down the soil profile, ideally to a depth of at least 2 m.

Label each sample clearly with the number of the hole and depth. Get the soil samples analysed for electrical conductivity. Sites should be sampled every six months to three years depending on the results and risk of rising water table.

#### Take water samples

Using a weighted dipper on the end of a piece of cord (or a submersible pump), take a sample of groundwater. Have this sample analysed for electrical conductivity.

#### Long-term monitoring

To detect whether the water tables are rising, falling or stationary over a period of years, check water table depths regularly and record or graph them. The most suitable time for measuring depths is in dry months, when the risk of capillary rise is greatest. It is a good idea to take measurements across the seasons to establish the range of water table heights.

Annual checking of the saltland boundary between the steel posts will also indicate whether the problem is getting worse, is stable or is retreating. Take photographs to record changes.

**A** 

References and resources

For access to relevant references and further resources click here.

Horticulture Australia

Further references and

the end of this chapter

resources can be located at

### 1d Salinity

Further references and resources can be located at the end of this chapter.



### **1e Soil acidity**



Objective - soil pH is maintained within the optimum range for crop production

Soils can be naturally acid or alkaline. Soil pH may also change with irrigation, fertiliser and crop management practices. As soil pH changes, the availability of soil nutrients may also change. Therefore it is important to monitor soil pH changes over time.

### Soil acidity

Soil acidification is a major land degradation issue, which can lead to reduced availability of nutrients, lower yields and fewer crop options. Soil acidity can be naturally occurring and can be made worse by prolonged and heavy use of nitrogen fertilisers like sulphate of ammonia and MAP (monoammonium phosphate). It can also be exacerbated by the removal of hay and alkaline materials.

The speed with which soil becomes acidic depends on many factors including soil type, soil texture (sandy soils become acidic more easily), organic matter, cation exchange capacity, the amount of crop product removed and the type of fertiliser used.

Older and more highly weathered soils are likely to have become acidic due to the natural processes of time and weathering. Calcium and, in particular, magnesium can be leached out of the soil profile under these conditions, contributing to acidity.

Acid sulphate soils - defined as soils with a pH reading below 5.5 - are formed when seawater or sulphate-rich water mixes, in the absence of oxygen, with land sediments containing iron oxide and organic matter. Acid sulphate soils are commonly found less than 5 m above sea level. Mangroves, salt marshes, floodplains, swamps, wetlands, estuaries and brackish or tidal lakes are ideal areas for acid sulphate soil formation.

The presence of acid sulphate soil may not be obvious on the soil surface as it is often buried beneath layers of more recently deposited soils and sediment.

When exposed to air due to drainage or disturbance, these soils produce sulphuric acid, which in turn can release toxic quantities of iron, aluminium and heavy metals.

#### Soil alkalinity

Alkaline soils have a pH greater than about 7.5 or a high sodium content, or both.

Alkaline soils may be deficient in zinc, copper, boron and manganese. Soils with an extremely alkaline pH (>9) are likely to have high levels of sodium.

Some soils in the semi-arid and arid regions have naturally high pH caused by significant quantities of free calcium carbonate. Irrigated well/bore water may also contain significant quantities of calcium carbonate.



Further references and resources can be located at the end of this chapter



Horticulture Australia



of the issue.

### 1e Soil acidity and alkalinity

LOW RISK - You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

HIGH RISK - You need to take some action. Read the Suggested Practices for that chapter.

Further references and resources can be located at the end of this chapter



### 1e Soil acidity



#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### **Suggested practices**

### Soil acidity

Lime or dolomite is usually added to maintain soil pH within a desirable range and can reverse the acidifying process in surface soils. Soil testing can help determine the correct rate to apply. Over-application can take years to remedy and can decrease uptake of nutrients by plants. It is usually easier to apply lime before planting. Thorough incorporation improves results, although incorporating lime into subsoil layers is difficult.

Not all soils can be maintained at pH 5.5 or above. Unless acidic soils are already cultivated and acid tolerant crops are to be grown, consider leaving them in their natural state, as drainage and cultivation can cause extreme acidification. If this occurs, liming is often expensive and often fails to achieve a lasting increase in pH.

Nitrogen leaching is a common form of soil acidification; reducing nitrogen, reducing leaching or using less acidifying forms of nitrogen can assist in reducing soil acidification. In some fertilisers the conversion from the applied form to one the plant can take up is a process that acidifies the soil. The acidification potential of different fertilisers is:

- Severely acidifying ammonium sulphate and monoammonium phosphate (MAP);
- Moderately acidifying diammonium phosphate (DAP);
- Slightly acidifying urea and ammonium nitrate; and
- Non-acidifying potassium nitrate, calcium nitrate and composted poultry manure.

Nitrates are highly mobile under the influence of high rainfall or over-irrigation and will readily leach in permeable soils. This process can cause further soil acidification and contamination of surface and groundwaters.

Legumes or nitrifying crops can also contribute to soil acidity.

Soil acidity can also develop under drip irrigation where soils are highly leached.

The following approaches should be considered:

- Efficient nitrogen application in order to applying smaller amounts of fertiliser more often. Fertigation (that is, applying fertilisers through irrigation systems) is one technique that can help match fertiliser application more effectively with crop demand and also allows for use of soluble fertilisers like potassium nitrate and calcium nitrate:
- Increase organic matter in order to help to buffer soil from pH changes; and;
- Do not drain, clear or expose acid sulphate soils in order to prevent mobilising the acids and toxic elements in the profile.

#### Soil alkalinity

Alkaline soils need to become more acidic. One way of achieving this is to use fertilisers such as crushed sulphur and some ammonium-based nitrogen fertilisers.

Elemental sulphur combines with oxygen and water to become sulphuric acid. This process can take some time and its effect on soil pH will depend on how much free calcium carbonate there is as this acts as a buffer.



Monitoring and recording

A check of the soil pH is an ideal way of monitoring the change in acidity of soils over time. It is important that pH is determined in soil samples taken to a depth of at least 60-80 cm to represent the root zone, and because surface lime applications often only increase pH to the depth the lime was incorporated. When collecting samples, be careful to separate the 0-15 cm and 15-30 cm samples from the deeper layers so that the acidity profile can be identified.

You can also measure pH with a simple test kit available from rural merchandise stores. The kit uses colour to indicate the pH level and is easy to use. It measures pH in water and not calcium chloride, so the results may be different from laboratory tests conducted by your local provider.

Soil water extraction tubes can be used to collect soil water samples at different depths for analysis of pH as well as other factors such as salinity and nitrates.



#### **References and resources**

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter



Horticulture Australia

### 1e Soil acidity and alkalinity

Further references and resources can be located at the end of this chapter



### **1f Sodicity**



Objective - soil permeability is adequate for water infiltration and drainage needs of the crop and erosion of sodic soils is minimised

Sodic soils are those where the amount of sodium held on to the clay particles is 6% or more of the total cation exchange capacity.

Sodic soils have an unstable structure and are poor places for plants to grow.

Following rain or irrigation with fresh water, clay particles in sodic soils force each other away, because of the sodium bound to the clay. This causes the soil to disperse, leaving a cloudy suspension.

Soil sodicity and soil salinity are often related because both involve sodium – a metal element widespread in Australian soils. Sodicity may be the more obscure problem, but it is a more widespread form of land degradation. It affects nearly a third of all soils in Australia (including one-third of all agricultural soils) and can cause poor water infiltration, low water storage, toxicity, surface crusting or sealing and waterlogging.

Many duplex soils (sandy topsoil over clay) in Australia have sodic clay subsoils. The structure of the subsoil clay is described as prismatic or columnar, which is hostile to plant roots. Tunnel erosion is also a risk on slopes with sodic clay subsoils.

The impacts of sodic soils extend to water catchments, infrastructure facilities and the environment. Run-off from sodic soils carries clay particles into waterways and reservoirs causing water turbidity, or cloudiness.



Further references and resources can be located at the end of this chapter



Horticulture Australia

### **1f Sodicity**

HIGH RISK – You need to take some action. Read the Suggested Practices for that chapter.

Further references and resources can be located at the end of this chapter



### 1f Sodicity



#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### **Suggested practices**

Improving sodic soils is not straightforward. Good soil management practices will generate as much benefit as high-input, costly remedial action, especially if the soil is in reasonable condition to begin with. Practices such as minimising tillage and compaction, returning large amounts of organic matter to the soil and improving surface drainage should be the first consideration before undertaking specific remedial action.

Once basic good soil management practices are adopted then attention can be turned to the sodicity problem itself. If sodic soils are identified or suspected seek professional advice.

#### Application of gypsum

Calcium in the form of gypsum can be added to sodic soils to improve their structure. Gypsum may also be applied to the soil via the subsurface drip irrigation system.

Gypsum improves soils in two ways. The first is a short-term benefit. The calcium salts in the gypsum increase the salt level in the soil solution around the clay particles. This prevents dispersion of sodic clays and promotes aggregation.

The second effect is a longer-term one. Sodium ions held on the clay particles are swapped with the calcium ions from the gypsum. This helps reclaim the soil and allows the displaced sodium ions to be leached out below the root zone.

The gypsum can be applied to riplines to help stabilise fracture lines, or broadcast and either incorporated or left on the surface. The best option depends on the nature, depth and extent of soil sodicity. Other factors to consider are soil pH, soil salinity, irrigation water quality, drainage, irrigation systems and the horticultural enterprise being undertaken. Because there are many factors to consider when dealing with sodic soils, it is advisable to seek professional advice.

It should also be remembered that gypsum adds to the overall salinity of soil water (though not in proportion to the amount used) so under pre-existing saline conditions, large applications should be made after completion of the growing season. Certainly it is better to apply smaller amounts annually rather than large, infrequent applications; this reduces opportunities for structural decline. The amounts required depend on the quality of irrigation water and soil type but are generally in the order of 5-10 t/ha in irrigated permanent plantings.

#### Further references and resources can be located at the end of this chapter



Application of lime

Lime (calcium carbonate) can also be used if the soil is not alkaline. However lime does not have the short-term benefits of gypsum.

Generally sodic soils take many years to improve using lime and gypsum. In the meantime it is important to manage these soils appropriately:

- Minimise tillage and avoid aggressive, deep working;
- Maximise returns or additions of stubble and organic manures to stabilise structure and maintain permeability;
- Install surface drains and cutoff drains to minimise waterlogging. Consider raised beds;
- Sodic soils can be irrigated with slightly saline water. Fresh water will maximise dispersion. Seek professional advice, as this can have negative effects on plant health due to increase in sodium and chlorine levels; and
- Avoid deep ripping unless soils are stabilised with gypsum, the fracture lines will collapse during the first full saturation, resulting in a more compacted and impermeable state than prior to ripping.

Technical advice should be sought to identify the suite of management options that are relevant and practical for the management of sodic soils in your particular situation.



### Monitoring and recording

Monitoring of soil physical properties and sodium levels is required to check the results of your management strategies.

Test the surface and the subsoil separately to determine the distribution of any sodicity problem.

- Collect samples from both the surface and the subsoil using a 5 cm soil auger or similar. Place the samples in clean buckets, one bucket for the subsoil and one for the surface soil;
- Collect samples randomly from a minimum of five locations over a uniform 1-2 ha representative area of the paddock;
- If it isn't clear where the subsoil begins, take a sample from the top 10 cm of the soil profile. Then take a second sample from somewhere deeper in the profile, within the range of 20-60 cm below the surface;
- Spread the soil from each bucket into a thin layer on a clean plastic sheet. Place in a well- ventilated location to air-dry, which may take several days;
- You need pieces of soil approximately 1 cm in diameter for this test. If necessary, break the air-dried soil into pieces and then mix thoroughly;
- From each surface and subsoil sample weigh 100 gm of soil into a clean 600 ml glass jar with lid;
- Measure out 500 ml rainwater or distilled water to give a 1:5 ratio of soil to water. Gently pour this water down the side, without disturbing the soil at the bottom;

### 1f Sodicity

Further references and resources can be located at the end of this chapter



### **1f Sodicity**

- Invert the jar once, slowly and gently, and then return to its original position (avoid any shaking). Then let stand for four hours, with no vibrations or bumping; and
- Check the suspension above the sediment at the bottom of the jar: Place a white plastic spoon or spatula in the solution (without stirring) to assist in determining the level of sodicity. Score the cloudiness using the following scale:



Estimating turbidity (soil sodicity) in a 1:5 soil/water suspension:

- 1. Clear or almost clear not sodic
- 2. Partly cloudy medium sodicity
- 3. Very cloudy high sodicity

Estimating turbidity using spatula visibility:

- 1. Plastic spatula visible not sodic
- 2. Plastic spatula partly visible medium sodicity
- 3. Plastic spatula not visible high sodicity
- Make up another soil suspension and repeat the process if unsure. Record the results.

A laboratory test for ESP (Exchangeable Sodium Percentage) and salinity of both surface and subsoil will provide the necessary information to decide management options.

For more on salinity monitoring information see 'Sodicity - a dirty word in Australia, Activity 1, A field test for sodicity', Australian Academy of Science.

**References and resources** 

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter



Horticulture Australia

#### Land and soil management - References and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www. horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

#### NATIONAL DATA SETS AND GUIDELINES:

Agricultural Production Systems slMulator (APSIM) www.apsim.info

This website contains a suite of modules which enable the simulation of systems that cover a range of plant, animal, soil, climate and management interactions.

#### ApSoil www.apsim.info/Products/APSoil.aspx

This tool provides access to a database of soil water characteristics enabling estimation of Plant Available Water Capacity for individual soils and crops. It covers many cropping regions of Australia and is regularly updated. It is designed for use in simulation modelling and agronomic practice.

Australian and New Zealand Guidelines for Fresh and Marine Water Quality - Volume 1: Paper 4 -Primary Industries (2000)

#### http://www.environment.gov.au/resource/australian-and-new-zealand-guidelines-fresh-and-marine-waterquality-volume-1-guidelines

This resource outlines the requirements for fresh and marine water quality. This version (2000) is currently being revised, with next version to be released mid-2014 - see www.environment.gov.au/topics/water/water-guality/ national-water-quality-management-strategy

Australian Bureau of Statistics http://www.abs.gov.au/ausstats/abs@.nsf/ featurearticlesbytitle/1DF2ACCC5071A76ACA2569DE00267E5A?OpenDocument This reference outlines some statistics for soil erosion.

Australian Collaborative Land Evaluation System www.clw.csiro.au/aclep/ This website is a national collaborative approach to Australian land and soil resources. This includes the agreed Australian Soil Classification framework www.clw.csiro.au/aclep/asc/index.htm

Australian Soil Resource Information System www.asris.csiro.au

This website provides online access to publically available information on soil and land resources.

AUSVEG, Healthy Soils http://ausveg.com.au/Default

#### Saltland Genie www.saltlandgenie.org.au/all-about-saltland/unit-4-plant-and-animal-performance/ nderstanding-salinity-and-waterlogging.htm

This website provides up to date information resulting from the National Dryland Salinity Program (1993 – 2005), national Sustainable Grazing on Saline Lands initiative, but built on wider work from the CRC for Plantbased Management of Dryland Salinity, and the National Land & Water Resources Audit.

SoilMapp www.csiro.au/soilmapp

This resouce provides direct access to best national soil data and information from the Australian Soil Resource Information System (ASRIS) and ApSoil, the database behind the agricultural computer model: Agricultural Production Systems SIMulator (APSIM).

#### SoilQuality http://www.soilquality.org.au

This website provides links to chemical, biological and physical calculation tools, as well as capturing soil quality information within a number of selected regions within WA, TAS, SA, QLD, and NSW only.

Soil Science Australia www.soilscienceaustralia.org

This website is the official site for the peak body on soil science.

**STATE - SPECIFIC LINKS** 

NSW – Department of Primary Industries

Soil erosion www.dpi.nsw.gov.au/agriculture/resources/soils Soil structure - SoilPak for Vegetable growers www.dpi.nsw.gov.au/agriculture/horticulture/vegetables/soil/ soilpak

Soil acidity www.dpi.nsw.gov.au/agriculture/resources/soils/acidity Salinity www.dpi.nsw.gov.au/agriculture/resources/soils/salinity General horticulture resources www.dpi.nsw.gov.au/agriculture/horticulture

### Land and soil management - references and further resources



NT - Department of Land Resource Management www.lrm.nt.gov.au/soil#.UncCWqVtzjA

#### QLD – Queensland Government

http://www.qld.gov.au/environment/land/soil/ Soil pH www.nrm.qld.gov.au/factsheets/pdf/land/l47.pdf WetlandInfo http://wetlandinfo.ehp.qld.gov.au/wetlands/index.html

#### SA – Primary Industries and Regions South Australia

Soil erosion www.pir.sa.gov.au/pirsa/more/factsheets/soil General horticulture resources www.pir.sa.gov.au/horticulture Irrigation and soil http://www.sardi.sa.gov.au/water/irrigation\_management/soils

#### TAS - Department of Primary Industries, Water and Environment

Soil erosion www.dpiw.tas.gov.au/inter.nsf/WebPages/TPRY-5YW4WC?open Field Mapped Land Capability Maps and Reports www.dpiwe.tas.gov.au/inter.nsf/WebPages/TPRY-6BF9EF?open

Soil structure www.dpiw.tas.gov.au/inter.nsf/ThemeNodes/EKOE-4ZG6PP?open Hamlet, A.G (Ed) 2002, Soil Management – A Guide for Tasmanian Farmers, DPIWE Tasmania www.dpipwe.tas.gov.au/inter.nsf/Attachments/TPRY-68TVRC/\$FILE/Soil%20Guide.pdf

#### VIC - Department of Environment and Primary Industries

Soil and water http://www.depi.vic.gov.au/agriculture-and-food/farm-management/soil-and-water Soil erosion www.dpi.vic.gov.au/agriculture/farming-management/soil-water Soil – Victorian Resources Online http://vro.dpi.vic.gov.au/dpi/vro/vrosite.nsf/pages/soil-home General horticulture resources http://www.depi.vic.gov.au/agriculture-and-food/horticulture Regional focus – Gippsland Soil Directory www.wgcma.vic.gov.au/index.php/component/content/ article/45/284-the-gippsland-soil-trial-directory.html

#### WA – Department of Agriculture and Food

Soils https://www.agric.wa.gov.au/climate-land-water/soils Soil salinity - A simple way to monitor your saltland (2005) www.agric.wa.gov.au/PC\_92345.html

Moore, G (2001) Soilguide: a handbook for understanding and managing agricultural soils. National Library of Australia. www.agric.wa.gov.au/objtwr/imported\_assets/content/lwe/rpm/landcap/soilguide\_introduction

#### LITERATURE

Alt, S; Jenkins, A; Lines-Kelly, R. (2009) Saving Soil – A landholder's guide to preventing and repairing soil erosion. Northern Rivers Catchment Management Authority, State of New South Wales through NSW Department of Primary Industries. http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0008/270881/saving-soilcomplete.pdf

Australian Government (2006) State of the Environment Report: Issue: Catchment scale influences - Land and vegetation condition - Erosion. Department of Environment, Canberra, ACT. http://www.environment.gov.au/ node/22498 (Note: Source of information on turbidity)

Australian Government (2011) State of the Environment Report. Department of Environment, Canberra, ACT. www.environment.gov.au/topics/science-and-research/state-environment-reporting/soe-2011/soe-2011report/5-land

Chapter 2.2 Soil | 2 Current state and trends of the land environment | 5 Land www.environment.gov.au/ node/22965

Australian Online Coastal Information, OzCoasts (2013) Measuring turbidity. GeoScience Australia. http://www.ozcoasts.gov.au/indicators/turbidity.jsp

Barson M, Mewett J. & Paplinska J. (2012) Land management practice trends in Australia's horticulture industry (soil acidity and maintaining ground cover). Caring for our Country Sustainable Practices Fact Sheet 4. Department of Agriculture, Fisheries and Forestry, Canberra, ACT. http://www.daff.gov.au/\_\_data/assets/ pdf\_file/0010/2148715/national-factsheet-farm-practices-horticulture.pdf

Bennett L. T., Mele P. M., Annett S. & Kasel S. (2010) Examining links between soil management, soil health, and public benefits in agricultural landscapes: An Australian perspective. Agriculture, Ecosystems & Environment 139, 1-12.

Biswas, T.; Bourne, J; Schrale, G. & McCarthy, M. (2009) Sustainable Irrigation Management Update - Salinity Management Practice Guidelines: Managing root-zone salinity for irrigated horticultural crops in winter rainfall zones of Australia. National Program for Sustainable Irrigation, Land & Water Australia, Canberra, ACT. http://npsi.gov.au/files/products/national-program-sustainable-irrigation/pn22225/pn22225.pdf

Bui, E; Hancock, G; Chappell, A; Gregory, L, (2010) Evaluation of tolerable erosion rates and time to critical topsoil loss in Australia, CSIRO Publishing. This report identifies regions of Australia at risk of exceeding 'tolerable' rates of soil loss.

Campbell A. (2008) Managing Australia's Soils: A Policy Discussion Paper. Department of Agriculture, Fisheries and Forestry, Canberra. http://www.clw.csiro.au/aclep/documents/Soil-Discussion-Paper.pdf

Caring for our Country (2013) Factsheet: Soil acidification in Australia, Caring for Our Country, Australian Government, Canberra, ACT. http://www.nrm.gov.au/funding/agriculture/innovation/pubs/soil-acidification.pdf

Cockcroft B. (2012) Soil Management For Australian Irrigated Horticulture. National Program for Sustainable Irrigation, Canberra, ACT. http://lwa.gov.au/files/products/national-program-sustainable-irrigation/npsi0212/ npsi0212-soil-management-australian-irrigated-hort.pdf

Cork, S; Eadie, L; Mele, P; Price, R; and Yule, D. (2012) The relationships between land management practices and soil condition and the quality of ecosystem services delivered from agricultural land in Australia, Caring for our Country, Australian Government, Canberra, ACT. http://www.daff.gov.au/\_\_data/assets/pdf\_ file/0008/2210597/soils-final-report-full.pdf

Cox, J. (2009) Effects of Sustainable mulching systems on soil health and production in SoilQuality http://www.soilquality.org.au This website provides links to chemical, biological and physical calculation tools, as well as capturing soil quality information within a number of selected regions within WA, TAS, SA, QLD, and NSW only.

DEFRA (2011) Protecting our water, soil and air, Department of Environment Food and Rural Affairs, UK. https:// www.gov.uk/government/publications/protecting-our-water-soil-and-air

Department of Employment, Economic Development and Innovation (2011) Wetland Management Handbook: Farm Management Systems (FMS) guidelines for managing wetlands in intensive agriculture, Queensland Wetlands Program, Brisbane, QLD. http://wetlandinfo.ehp.gld.gov.au/resources/static/pdf/resources/reports/ fms/fms\_025\_handbook\_web.pdf

Department of Natural Resources, Mines and Water (2005) Land Manager's Monitoring Guide: Soil pH indicator, Department of Natural Resources, Mines and Water, QLD. http://s3.amazonaws.com/zanran\_storage/ www.nrw.qld.gov.au/ContentPages/2343095.pdf

Doolittle, W.E. (2004), Measuring erosion. Available from http://uts.cc.utexas.edu/~wd/courses/373F/notes/ lec17ero.html. This report provides information on how to measure erosion.

ERIN (2009a) Indicative locations where improving soil and land management practices to reduce soil loss from wind erosion will provide the biggest benefits (map). Environmental Resources Information Network Canberra, http://nrmonline.nrm.gov.au/catalog/mgl:2348

ERIN (2009b) Indicative locations where improving soil and land management practices to reduce soil loss from hillslope (sheet and rill) erosion will provide the biggest benefits (map). Environmental Resources Information Network, Canberra, ACT. http://nrmonline.nrm.gov.au/catalog/mgl:2386

ERIN (2009c) Indicative locations where improving soil and land management practices to increase soil organic matter will provide the biggest benefits (map). Environmental Resources Information Network, Canberra, ACT. http://nrmonline.nrm.gov.au/catalog/mgl:2390

Glowinski, L. (Updated 2008) The Complete Book of Fruit Growing in Australia, Hachette Australia. http://www. publish.csiro.au/pid/7098.htm (for purchase via this link).

HAL (2011) Benchmarking Soil Health for Improved Crop Health, Quality and Yields in the Temperate Australian Vegetable Industries. Horticulture Australia Limited, Sydney, NSW. www.horticulture.com.au

Hunt, N. and Gilkes, B. (1992) Farm Monitoring Handbook. University of Western Australia, WA.

Isbell, R. (2002 – revised edition) The Australian Soil Classification – Revised Edition – Australian Soil and Land Survey Handbook Series 4, CSIRO Publishing. www.publish.csiro.au/pid/3529.htm (for purchase via this link). This publication provides a hierarchical soil classification system of five categorical levels from general to the most specific: order, suborder, great group, subgroup, and family. The Australian Soil Classification describes fourteen soils orders across Australia with further subdivision under these broad groups.

Kelly, J. (2011) Australian Vegetable Industry Soil and Land Management Knowledge Exchange (HAL Project Number: VG07146). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Kelly, J. (2009) Soil and Land Management Extension Resource Manual (HAL Project Number: HG06165). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au





LEEPER, G; and UREN, N, (1993), Soil Science, An Introduction, 5th edition, Melbourne University Press. http://trove.nla.gov.au/version/44659070 (Note: link provides advice for borrowing this edition)

Maas and Grattan (1999) Salt tolerance of crops, FAO. www.fao.org/docrep/005/y4263e/y4263e0e.htm McKenzie, N; Jacquier, D; Isbell, R; and Brown, K. (2004) Australian Soils and Landscapes – An Illustrated Compendium, CSIRO Publishing. http://www.publish.csiro.au/pid/3821.htm (for purchase via this link). This publication provides information on how soil properties and landscape processes should guide our use of the land.

McPhee J. E. (2009) Controlled Traffic Farming Systems for the Tasmanian vegetable industry (HAL Project Reference VG07058). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

McTainsh G. H., Leys J. F., O'Loingsigh T. & Strong C. L. (2011) Wind erosion and land management in Australia during 1940- 1949 and 2000- 2009. Report prepared for the Australian Government Department of Sustainability, Environment, Water, Population and Communities on behalf of the State of the Environment 2011 Committee. DSEWPaC, Canberra.

Murray, R.S. (2007) A Review of Dr Bruce Cockroft's Work for Australian Irrigated Horticulture, Milestone Report 2, National Program for Sustainable Irrigation, Narribri, NSW. http://npsi.gov.au/files/products/ national-program-sustainable-irrigation/pn21945/pn21945.pdf

Murray, R.S.; McCarthy, M; Grant, C, (2011) Guidelines for Managing Soil Structure in Irrigated Vineyards, National Program for Sustainable Irrigation, Narribri, NSW. http://npsi.gov.au/files/products/national-program-sustainable-irrigation/npsi2311/npsi2311-guidelines-managing-soil-structure-irriga.pdf

NPSI (2012) Irrigation Essentials Updated – Research and innovation for Australian irrigators. National Program for Sustainable Irrigation. Cotton RDC, NSW. http://npsi.gov.au/files/products/national-program-sustainable-irrigation/npsi06121/npsi06121-irrigation-essentials-updated.pdf

Pattison T., Moody P. & Bagshaw J. (2010) Soil Health for Vegetable Production in Australia. Queensland Government, Brisbane. http://www.daff.qld.gov.au/26\_17025.htm

Pattison, T. (2008) Soil and root health for banana production (HAL Project Number: FR02025). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Porter, I. (2008) Management of soil health for sustainable vegetable production (HAL Project Number: VG06090). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Quirk, J, Sodic Soils, Australian Academy of Science www.science.org.au/nova/035/quirk.html

Reid G. (2002) Soil and Nutrient Loss in Macadamia Lands. (HAL Project Reference MC98011). Horticulture Australia Ltd, Sydney.

Slavich P. G. & Cox J. (2010) Improving soil and orchard floor management in macadamias. (HAL Project Reference MC05001). Horticulture Australia Ltd, Sydney.

Smith J. & Leys J. (2009) Identification of areas within Australia for reducing soil loss by wind erosion. Bureau of Rural Sciences, Canberra. http://nrmonline.nrm.gov.au/catalog/mgl:2246

Soils Research Development and Extension Working Group. (2011) A stocktake of Australia's current investment in soils research, development and extension: A snapshot for 2010-11. Department of Agriculture, Fisheries and Forestry, Canberra. http://www.daff.gov.au/\_\_data/assets/pdf\_file/0003/2085816/soils-stocktake.pdf

Sylvain Z. A. & Wall D. H. (2011) Linking soil biodiversity and vegetation: Implications for a changing planet. American journal of botany 98, 517-27.

Synder, C, pH chapter of Efficient Fertiliser Use Manual http://www.pegasuskó2.com/efficient-fertilizer-use-guide

Ugalde D., Brungs A., Kaebernick M., McGregor A. & Slattery B. (2007) Implications of climate change for tillage practice in Australia. Soil & Tillage Research 97, 318-30. For access to relevant references and further resources click here.





# Chapter 2 Water management



Objective – to maximise water use efficiency without compromising water quality on-farm and downstream

Water is a valuable resource and in many areas is becoming increasingly scarce. Good irrigation management is essential to maximise yields and control product quality.

Availability of water is increasingly subject to government regulations. These are designed to ensure waterways and groundwater. Extractions are at sustainable levels and protect the health of aquatic environments

Water management considers both the crop's water demand and the amount of water available. It also involves management of irrigation to maximise efficient use of water applied.

Drainage water and run-off also need to be managed to avoid any impact, such as nutrient pollution, on groundwater or waterways and wetlands.

This section is consequently split into the following sections:

- 2a Irrigation efficiency;
- 2b Water quality;
- 2c Managing wastewater.

Further references and resources can be located at the end of this chapter.

### 2a Irrigation efficiency



#### Objective

- uniform application of water to match crop needs
- drainage impacts are managed in accordance with environmental, community and regulatory standards

Irrigation efficiency is a term that helps us define the proportion of irrigation water that is actually taken up and used by the crop. Improvement in irrigation efficiency is normally associated with water savings, production gains and better long-term environmental management.

Irrigation efficiency is determined by irrigation management factors such as:

- Ensuring irrigation systems are operating to design specification and applying water as evenly as possible;
- Ability to time, or schedule irrigation, based upon crop water needs and clear understanding of soils' water holding, infiltration and drainage capacity.

To manage irrigation efficiently a number of management practices need to be considered, starting with an understanding of water availability and crop requirements.

#### Efficient irrigation management practices

There are nine basic steps involved in the efficient management of irrigation:

#### Identify

1. Define property goals and implications for water management

#### Plan

- 2. Know your soils
- 3. Design the most suitable irrigation system
- 4. Develop a farm water budget
- 5. Know your water supply/ies

#### Do

- 6. Determine a basic irrigation schedule
- 7. Implement strategies to manage nutrient input and salinity

#### Monitoring and recording

- 8. Monitor, record and evaluate
- 9. Check irrigation system performance

See the following two sections for more detail on these steps.



Horticulture Australia

the end of this chapter

### 2a Irrigation efficiency



#### HIGH RISK

iew produ ent Rev his chapte



neck performanc the irrigation system Review the Suggested Practices in this chapter

> HIGH RISK Review the ggested Practice in this chapter



LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

HIGH RISK - You need to take some action. Read the Suggested Practices for that chapter.

Further references and resources can be located at the end of this chapter





#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### **Suggested practices**

### 2a.1 Identify your goals

Your goals will largely depend on the crop(s) you are growing and desired yield and quality. The property goal can be made up of a series of block or paddock goals. Once the goal is defined, you can identify the right irrigation management strategies to help meet your goal, e.g. growing 150 tonnes of Class A onions for the export market, growing 200 tonnes of processing potatoes, or establishing 2 ha of cherry trees and producing 20 tonnes of processing peaches from existing orchard.

The goal may also be influenced by average annual rainfall, capacity of the irrigation water source, regulatory restrictions on water storage or access to surface and underground water. A fairly accurate estimate of expected crop water use will assist in balancing property water supply with yield and quality targets.

#### 2a.2 Know your soils

A soil survey is a fairly comprehensive analysis of soil types and their distribution across your property. Soil surveys establish a better understanding of your soil's ability to hold water and any potential physical and chemical limitations to growing your crops in that soil.

Soil surveys assist in determining if the land is suitable for developing particular crop types and help identify the irrigation system types that may be most suitable, manageable and efficient.

Soil surveys are also useful for identifying:

- Soil structural issues that may result in limited drainage, surface run-off, soil structure decline and root growth problems; and
- Soil chemical and nutritional characteristics that may directly effect plant growth or result in long-term soil quality decline (i.e. soil acidity, salinity and sodicity).

The types of measurement often referred to in a farm soil survey are:

- Readily Available Water (RAW) this is the water content value most relevant to irrigators. As the name suggests, Readily Available Water is the water component that can be readily used by the crop under ideal growing conditions. It is not the total soil water content. Ideally irrigation should replace water that has been removed by growing crops (there are exceptions to this generalisation but it stands as a sound 'rule of thumb');
- The infiltration rate of soil another valuable parameter which assists with better matching the application rate of the irrigation system with the soil's capacity to absorb this water without wastage or run-off.

#### **Readily Available Water (RAW)**

RAW is determined by soil texture and the rooting depth of the crop. Table 2.1 below helps you estimate how many millimetres of readily available water<sup>\*</sup> would be held in a metre of topsoil.

Table 2.1: Readily available water for different soil types. Source: Lovell (2006).			
	Readily available water (mm/m) between - 8kPa and		
Soil texture	-40Kpa	-60Кра	
Sand	35	35	
Loamy sand	50	55	
Sandy loam	60	65	
Loam	65	75	
Sandy clay loam	60	70	
Clay loam	55	65	
Clay	45	55	

Readily Available Water, is the water available between the soil being at full point (-8kPa) and dried to a tensiometer or gypsum block reading of -40kPa and -60 kPa. More information can be found in Growcom Water for Profit irrigation fact sheets here.

#### Infiltration rate

Infiltration rate is the speed at which water can move through a soil. Infiltration rate is related to soil texture, bulk density, organic matter, surface soil stability and groundcover.

The infiltration rate of a soil determines the maximum rate at which irrigation should be applied. Applying irrigation at a higher rate results in surface run-off. Table 2.2 below provides indicative irrigation rates.

Table 2.2: Average infiltration rates or some soil types. Source: Wise Watering Irrigation Management Course 2001.

Suggested application rate (mm/hr)						
Soil texture	Average soil structure Well-structured soil Infiltration rate range (mm/hr)					
Sand	50		20-250			
Sandy loam	20	45	10-80			
Loam	20	45	1-20			
Clay loam	20	40	2-15			
Light clay	2		0.3-5			
Medium – heavy clay	0.5		0.1-8			

(see http://www.dpiw.tas.gov.au/internnsf/Attachments/JMUY-5FP77U/\$FILE/IntroductiontoWiseWatering.pdf).

Further references and resources can be located at the end of this chapter



Horticulture Australia

Understanding soil characteristics across the property is vital for determining irrigation frequency and depth of application. It also allows soils with similar characteristics to be grouped into the same irrigation management area, based on the amount and rate at which irrigation water can be applied. It can be used to select representative sites to monitor soil moisture.

### 2a Irrigation efficiency





#### 2a.3 Design irrigation systems

Crop production will suffer if the irrigation design or the irrigation method does not suit your property goals or the soil type. One of the key aspects of design is to match irrigation delivery with water demand.

Consideration should be given to issues such as crop type, variety, harvest dates, soil type, topography and exposure to drying winds. Poor designs reduce irrigation efficiency, cause uneven water application and uneven crop yield and quality. Efficient fertiliser application, particularly fertigation, depends on uniform water application.

Different irrigation system options include:

- Drip irrigation (both surface and sub surface buried tape);
- Micro-sprinklers;
- Capillary bed (for containers);
- Sprinkler irrigation;
- Travelling gun irrigation;
- Centre pivot and linear move irrigation; and
- Surface (flood or furrow) irrigation.

In general, pressurised irrigation systems are a more efficient form of water delivery than surface flood or furrow irrigation. Growers in many irrigation regions are moving away from flood/furrow irrigation systems and changing to pressurised systems that enable more accurate and manageable water delivery. Whatever irrigation method is chosen, the system must be designed to accurately match soil type and plant water demand and protect the environment.

The table below gives some range of expected irrigation application system efficiencies.

Table 2.3: Expected irrigation application system efficiencies. Source: Lovell (2006).

System	% Efficiency
Rain gun (cannon, travelling irrigator)	50 - 75
Fixed Sprinkler	65 - 85
Linear move	75 - 90
Centre pivot	75 - 90
Drip	80 - 90

#### There are many factors in determining the most suitable irrigation system. Factors to consider include:

soil types and variation	topography
Readily Available Water (RAW)	maximum crop water demand
yearly water allocation	water quality
maximum extractable or stored water supply	fertigation
markets/end use of crop	allowances for leaching fraction
the need for frost protection	the need for crop cooling
the option of a cover crop	potential pests and diseases
maintenance and longevity of system	average annual rainfall recycle
structure	microclimate variation
cost to install and maintain system	climate
long-term reliable extractable or stored water supply	

Using experienced irrigation designers and installers for your crop is important. The Irrigation Association of Australia has a list of certified irrigation designers. See www.irrigation.org.au for more information.

#### 2a.4 Developing a farm water budget

A farm water budget is about making sure you have enough water to meet the property goals. Water budgeting helps determine the amount of water you expect to use over the season and attempts to match this with intended irrigated crop area so that the horticultural business can check that planned irrigation needs are within water entitlements.

Water requirements need to be budgeted using measurements of crop water demand at different times of the year, the irrigation system and knowledge of the soil water holding capacity. Individual farm data is best, but in some districts average crop water demands have been calculated and are available from agronomists and irrigation specialists.

Farm water budgets can be refined to crop-specific water budgets and irrigation schedules. There are two main types of irrigation schedule, indirect and direct. These are discussed further below (see 'Determine a basic irrigation schedule' later in this section).

An example of a water budget can be seen below in Table 2.4.

Table 2.4: Example of a water budget. Source: Lovell (2006).

An example of a water budget:				
Property name: Year: Intended crops:				
Сгор	Variety	Crop area (ha)		Water requirements / ha
Total water requirement for property				
TOTAL water allocation for property				
Sufficient water available to grow intended crops?				

#### 2a.5 Know your water supply

Understanding your crop water requirements and reliability of water supply is crucial.

The availability of water will affect the choice of crop type, the irrigation system and your irrigation management strategy. For example, drip systems (above ground or sub-surface) require a water supply that allows irrigation at short notice (e.g. within 24 hours) particularly during hot weather. Check with your state water agency to ensure the necessary licences/permits are obtained. In some districts with water allocation schemes, delays in ordering and then receiving water may limit your ability to adopt these practices. On-farm storage will reduce this problem, however, it is advisable to check whether licences/permits are required to construct dams. Interference with a waterway or obstruction of flow may require a licence/permit.

Comparing your crop water requirements against the quantity of water supply available will determine if you have sufficient water. Water may be limiting on an annual basis or sometimes in peak demand periods. If water supplies are limiting or uncertain, more efficient irrigation techniques and drought management strategies need to be considered.

Further references and resources can be located at the end of this chapter



Horticulture Australia

### 2a Irrigation efficiency

#### Further references and resources can be located at the end of this chapter


# 2a Irrigation efficiency

Strategies to lessen the impacts of drought include:

- Identify limits to systems and management and investigate crop options by preparing an irrigation and drainage management plan;
- Prepare a risk management strategy for reductions in water availability;
- Maintain soil structure to allow the full root zone to be exploited;
- Retain groundcover for erosion protection and to provide mulch; and
- Seek advice on technical matters, financial matters, and sources of assistance.

#### Following the drought:

- Reassess your irrigation and drainage management plan;
- Assess the market prospects of various crops and enterprises;
- Determine when water will be available again;
- Assess options for planting crops with or without assured water supply; and
- Watch out for weeds introduced by stock and bought-in fodder.

### 2a.6 Determine a basic irrigation schedule

Irrigation scheduling includes determining when and how much to irrigate. Growers have traditionally relied on their knowledge and experience to schedule irrigation. However many growers are now using other measures, such as soil moisture monitoring, to fine-tune their irrigation scheduling. Irrigation scheduling can be done by indirect or direct means.

A correct irrigation schedule can optimise yield and crop quality, and reduce the amount of water used. In turn, pumping and water costs may be reduced and drainage minimised. Applying the right amount of water will also avoid leaching of nutrients beyond the root zone. This careful irrigation management helps to achieve production targets and has environmental benefits.

#### Indirect irrigation scheduling

Crop water demand can be determined by estimating the total loss of water from the crop based upon information such as local evaporation and expected crop water loss through the leaf surface referred to as 'evapotranspiration' (ET). Refer to Growcom Water for Profit - irrigation fact sheets http://www.growcom.com.au/land-water/waterfor-profit/resources-water-for-profit/

The Bureau of Meteorology website provides maps of average monthly evaporation for Australia for each month of the year. Example of use: if the monthly map shows that an average evaporation of less than 250 mm during that month (~8mm per day) that represents a monthly water demand of 2.5 ML per hectare.

Some growers use software to develop a daily water budget. An example is shown below. This uses crop coefficients or crop factors to convert daily evaporation readings into estimated crop water use. Refer to Growcom Water for Profit – irrigation fact sheets http://www.growcom.com.au/land-water/water-for-profit/resources-water-for-profit/

Table 2.5: Example of an irrigation scheduling spreadsheet. Source: Lovell (2006)

SITE: SHEPPARTON YEAR 2003/04				Volume applied to this crop: 42 ML					
			CR	OP = PE	ACH	IES			
Fotal hectares as entered: 30   Fotal volume used for year: 42			He	CROP = PEACHES     Hectares   30     Crop Factor   Effective rainfall     0.9   0.8     Potential ET   Effective     Rainfall   Applied     deficit   (mm)     (mm)   (mm)     4.95   0     4.05   0     4.05   0     4.05   0     5.3   0     5.3   0     5.3   0     1.8   8     -2.15   7.92     0   5.3.3     0   11.89     5.3   0     1.8   8     -2.25   0     19.45   4.95     4.95   0     33.3   8.9     0   33.3     8.9   0     4.95   0     3.56   8.9					
Entor data in	white squares						all		
	n soil moisture deficit is gr	eater than 50 mm		0.9			0.8		
			F		Rai	nfall	Applied	and the second second	
Date	Evaporation (mm)	Rainfall (mm)			(ញ	m)	( <u>mm</u> )	( <u>mm</u> )	
23 Nov	5.5			4.95	1	0	40	0	
24 Nov	4.5			4.05		0		4.05	
25 Nov	2	10		1.8		8		-2.15	
26 Nov	8.8			7.92		0		5.77	
27 Nov	6.8			6.12	1	0		11.89	
28 Nov	5.9			5.3		0		17.2	
29 Nov	2.5			2.25		0		19.45	
30 Nov	5.5			4.95		0		24.4	
1 Dec	9.9			8.9		0		33.3	
2 Dec	9.9			8.9	1	0		42.22	
3 Dec				6.39		0	50	-1.39	
4 Dec	5.5			4.95		0		3.56	
5 Dec	9.9			8.9		0		12.47	
6 Dec	10.			9.09		0		21.56	
7 Dec				10.89	(	)		32.45	
8 Dec	9.9			8.9		0		41.36	
9 Dec	9.7			8.73		0	50	0.09	
10 Dec	8.8			7.92	0	)		8.0v	
11 Dec	1.	50		0.99	4	0		3	

#### **Direct irrigation scheduling**

Direct irrigation scheduling is achieved by measuring soil moisture. Soil moisture monitoring tools can provide information that can be used to check and improve irrigation scheduling. Such tools can help estimate the size of the wetting pattern (particularly depth), the level of plant stress and the additional moisture provided by rainfall. Monitoring soil moisture allows the initial estimates of how much and when to irrigate to be adjusted to suit a specific irrigation system and soil type.

Soil moisture monitoring tools normally measure soil water content or soil water tension.

Examples include tensiometers and resistance blocks (e.g. gypsum blocks) for measuring soil water tension, and neutron probes and capacitance probes for measuring soil water content.

Modern soil moisture systems track and graph soil moisture over time and can be used to help plan future irrigation dates. How much water to apply (hours) per irrigation to fully refill the soil can be calculated from the amount of water held in the soil and the system application rates.





Horticulture Australia

Further references and resources can be located at the end of this chapter



Chapter 2 Water management 2.9 The operational and maintenance requirements/costs of each system should be evaluated before making a decision. The nature of the crop may also have a bearing, for example annual crops require a more mobile system.

Regardless of the type of tools used, applying the right amount of water at the right time depends knowing crop water requirements, and stage of growth;

- Consideration of effects of weather;
- Water availability in the root zone (root zone raw);
- Wetting pattern (for micro-irrigation);
- Irrigation system efficiency; and
- Any leaching requirement (for managing high salinity).

Measuring if the volume of irrigation is adequate or excessive can also be achieved using 'wetting- front' detector-type sensors such as the CSIRO FullSTOP™

### 2a.7 Implement strategies to manage nutrient input and salinity

### Manage nutrient inputs

For nutrients to reach the crop roots and to avoid losses from over irrigation, fertiliser should be applied when soils are close to field capacity, i.e. late in the irrigation run. Over-irrigation or application of a leaching fraction will wash the nutrients past the root zone. For more information see Chapter 4 – Nutrient management.

### Manage salinity

Soil salinity can potentially reduce production by up to 100% due to reduced plant growth. This is because soil salinity makes it difficult for crops to obtain water and nutrients from the soil. Affected plants show similar symptoms to under-watering or can show visual symptoms such as burning on leaves. Soil salinity can also affect the biological health of the soil, which can have serious long-term effects on soil fertility. There is also a risk to soil structure if the soil becomes sodic. Soil salinity testing should be done regularly to monitor root-zone salinity.

There are a number of possible causes for high soil salinity (saline irrigation water, high water table, poor drainage, inadequate leaching, low rainfall). For more information see Topic 1d – Salinity.

If slightly saline water is used for irrigation, or deficit irrigation practices are used, in low rainfall areas, then it is essential to monitor the build up of salt in the root zone and apply leaching fractions when necessary.

When applying leaching irrigations there needs to be a balance between removing salt and minimising the loss of nutrients. Therefore, specific leaching irrigations should consider the potential for nutrient losses from the rootzone; e.g. avoid leaching when using fertigation or if having recently applied fertilisers.

### Checking the salt index of fertilisers

All fertilisers have a salt index, which indicates what the fertiliser contributes to soil salinity.

If your irrigation water or soils are saline, changing to fertilisers with similar nutrients but with a lower salt index may help. For example, potassium chloride has a salt index of 114 but potassium sulphate has a lower salt index of 46.

Information on the salt index of each fertiliser should be available from your local supplier.

Salinity is measured as Total Dissolved Salts (TDS) in eithermilligrams per litre (mg/L) or the equivalent, parts per million (ppm); 100 mg/L = 100 ppm. Electrical Conductivity (EC) is a good indicator of TDS. It is measured as EC units (equivalent to micro Siemen per centimetre) and can also be expressed as milli Siemen per cm (mS/cm) or the equivalent, desi Siemen per metre (dS/m); 1mS/cm = 1dS/m. NB:1,000 EC units = 1 mS/cm= 1,000 uS/cm = 1 dS/m =640 mg/L.



Monitoring and recording

### 2a.8 Monitor, record and evaluate

Monitoring, measuring and recording activities are essential for the overall management of the property. A range of factors should be monitored and evaluated but the following are important:

#### Monitor crop performance

Keeping records of crop productivity is important to understand the effects of different irrigation practices. Measuring and recording yield, quality and maturity for each crop allows yearly comparisons and evolution against the goal of the property, and helps to refine management decisions.

#### Document water budget

Record irrigation schedules, amount of water applied, rainfall, soil moisture and crop evapotranspiration.

### Assessment of economic yield

One measure of irrigation efficiency is through assessment of economic yield. This can be expressed in gross income per megalitre (\$/ML) and/or production water use efficiency (tonnes of produce/ ML). While no definitive figures exist for these criteria, historical on-farm or district comparisons will provide useful benchmarks.

#### Monitor water quality

Monitoring the quality of your drainage water can give an indication of nutrient loss. See Topic 2b – Water quality.

## 2a.9 Check irrigation system performance

You need to regularly check and maintain your irrigation system to make sure it is operating correctly and delivering what it should. If the system is not operating at maximum efficiency, irrigation scheduling and management strategies, such as controlling salinity, will not be effective.

Checks that should be undertaken include:

- Visual inspection of irrigation system and crop performance;
- Discharge or flow rate variation;
- Uniformity of water distribution;
- Pressure variation;
- Presence of cuts, blockages, leaks in dripper lines;
- Sprinkler/dripper malfunction;
- Filters; and
- Pumps.

It is also important to measure output uniformity or distribution uniformity. Uneven distribution causes areas of over/under irrigation and has consequences for crop yield and quality. The manufacturer's specifications should be referred to when assessing distribution uniformity and should be within the range specified in the table showing expected irrigation application system efficiencies table earlier in this section.

Further references and resources can be located at the end of this chapter



Horticulture Australia

# 2a Irrigation efficiency

#### Further references and resources can be located at the end of this chapter



#### Farm Dam Management - monitoring and mitigation

Conservative estimates suggest that in excess of 8,000 GL is stored in farm dams (i.e., 9% of total stored water) and that there are more than 2 million farms dams across Australia. Reducing this loss using either physical or chemical covers therefore has the potential to significantly increase agricultural water use efficiency in Australia and therefore generate millions of dollars for the Australian economy.

Unlike losses due to evaporation, which will vary based on seasonal variations of evaporative demand, seepage losses will be affected by soil permeability and dam construction.

For some storages it is possible to lose more water from seepage than evaporation. Storages that have excessive seepage losses will require some form of mitigation.

By measuring changes in water depth for periods when there is no inflow, outflow or rainfall, the components of evaporation and seepage can be directly measured as the change in depth. The Evaporation and Seepage Ready Reckoner or technologies such as EvapCalc, Irrimate<sup>™</sup> Seepage and Evaporation Meter or 'Pressure Sensitive Transducer' can be used. See Farm Dam Management Resource Kit for more information http://ncea-linux.usg.edu.au/farmdammanagement/



### **References and further resources**

For access to relevant references and further resources click here.



Objective - water quality is suitable for its intended use on the property and does not negatively impact downstream water quality

There are two aspects of water quality that need to be considered. The first is to make sure that the quality of water being used is suitable for the intended purpose (e.g. irrigation, agricultural sprays, packing sheds), and the second is to make sure that your operation is protecting the quality of water leaving your farm so it does not negatively impact on downstream users and the environment.

If you are sourcing water from rivers or streams then upstream farms and businesses may impact on you.

Problems caused by using poor quality water on-farm include:

- Salinity (high total soluble salt content);
- Sodicity (high sodium content);
- Toxicity (high concentration of specific salts in the soil);
- Blue-green algae, which may be toxic; clogging of irrigation equipment; and corrosion of pipes and other equipment.

One of the factors that needs to be considered is the proportion of dissolved minerals and salts in your irrigation water.

All groundwater and stream waters contain dissolved minerals. When irrigation water is used, the mineral salts are either taken up by the crop, left in the soil after the crop has used the water, leached down past the root zone, or washed out with run-off. Most of these salts are beneficial, but in some cases they may be harmful to the crops and to the long-term sustainability of the property. See Topic 1d - Salinity.

Other chemical contaminants of water may include heavy metals, and agricultural or industrial chemicals.

The potential impact of poor quality water leaving the farm includes:

- Harm to aquatic species in waterways from water eutrophication due to nutrient and organic matter pollution, and from chemical pollution;
- Sedimentation of waterways and marine environments, causing disruption and damage to these ecosystems; and
- High nutrient levels in waterways contributing to blue-green algae outbreaks.

Further references and resources can be located at the end of this chapter



Horticulture Australia

# **2b Water quality**

Further references and resources can be located at the end of this chapter



# **2b Water quality**



# **2b Water quality**

HIGH RISK Risk assessments

opics 1a and 1b. Review Suggested Practices in relevant chapters.

#### HIGH RISK

Review the uggested Practices in this chapter.

## HIGH RISK

ee Risk assessments in bics 3 and 4. Review the uggested Practices in relevant chapters.

### HIGH RISK

ee Risk assessments in bics 3 and 4. Review the buggested Practices in relevant chapters.

LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

**HIGH RISK** – You need to take some action. Read the Suggested Practices for that chapter. Further references and resources can be located at the end of this chapter.



# **2b Water quality**



### **Review checklist**

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

#### Check water source quality

This should be a priority when considering new enterprises. Good data is often available from your water supply authority/company/State government agency.

Where use of saline water is unavoidable, regularly check salinity to plan suitable irrigation management options.

It is important to remember that water quality can change from month to month and summer flows in a river system can have quite different water quality to winter/spring periods.

Recycled water can be derived from various wastewater sources and has many uses. Recycled water can be used for a variety of purposes. For each use the water must be treated to a level where it is considered fit-for-purpose. The Environment Protection and Heritage Council, the Natural Resource Management Ministerial Council and the National Health and Medical Research Council developed guidelines for the safe use of recycled water. These guidelines are now stored at http://www.environment.gov.au/resource/ national-water-quality-management-strategy-australian-guidelines-water-recyclingmanaging-0. It is important to check whether your relevant state authorities approved the recycled water for the intended use (e.g. fit for purpose). For more information on using recycled water see http://www.recycledwater.com.au/index.php?id=96

#### Check quality of water leaving the farm

It is also worth checking the drainage and run-off water leaving your own property. How does it compare with the water upstream or your neighbours? If the water is high in nutrients and turbidity (water cloudiness) then you should consider how fertiliser management, soil erosion, protecting watercourses and agricultural chemical management could be improved. See 1a - Soil erosion caused by water; 1b - Soil erosion caused by wind; 3 – Chemical management; and 4 – Nutrient management.

Your water authority, local catchment authority or Landcare group can usually provide access to water testing laboratories and information on run-off water quality targets.

#### Protect water quality

Water quality is impacted by activities both on and off farm. It is important to be aware of on-farm activities that can negatively affect water quality as this may impact the suitability of the water for use on the farm as well as having significant environmental impacts. Farm activities may affect water quality by increasing levels of salts, nutrients, suspended sediment, chemicals or organic matter.

#### Protect watercourses

Watercourses such as rivers, creeks and streams as well as their riparian areas (areas on or near creek and river banks) should be protected. Areas that have significant protected riparian zones have the ability to capture and filter soil sediment and soluble nutrients, improving water quality before it leaves the farm. A strip of undisturbed vegetation should be left to protect waterways.

Revegetate riparian areas with a mixture of native grasses, shrubs and/or trees to provide a buffer and stabilise waterway banks. In some regions there are legislated separation distances (for example 100 m) between sources of groundwater or surface water and obvious pollution sources such as fertiliser or agricultural chemical storage, packing sheds and workshops. It is advisable to check with the relevant government agency.

In known drainage lines or areas where run-off enters waterways install filter strips or buffer strips to minimise sediment and nutrient entering waterways. Seek information regarding design of the buffer strips, particularly in relation to the most appropriate vegetation and width of strip. Fencing waterways to keep stock out and providing offstream drinking points also help protect watercourses.

Financial assistance may be available to fence riparian zones. Contact your local Landcare, catchment authority or government representatives.

#### Soil erosion

Soil erosion is an important issue for both soil protection and water quality protection. High turbidity of run-off indicates soil loss is occurring. This is most common after intense rainfall, particularly after a dry spell. Buffer zones or grassed areas can be established to filter run-off and storm water. Often nutrients, especially phosphorus, and farm chemicals are carried attached to soil particles. Controlling soil erosion will help to retain nutrients and reduce nutrient pollution downstream. See Section 1a - Soil erosion caused by water, and 1b – Soil erosion caused by wind, for more details.

#### Nutrient management

Nutrient management is important to ensure that the nutrients applied are either used by the crop (some of which will be exported off-farm in the harvested product) or safely stored in the soil for the next crop.

Fertiliser or nutrients can be applied through fertiliser or nutrient rich water (e.g. recycled water can have significant concentrations of nitrogen and phosphorus). All nutrient sources should be considered when deciding crop nutrient requirements (nutrient budgeting).

Inaccurate or over-application of fertilisers can contaminate ground and surface water. This can result in the enrichment of water with nitrogen or phosphorus (eutrophication) causing rapid growth of algae and aquatic plants. This disturbs the balance of organisms present in water and the quality of the water within waterways. Nitrogen leaching can also cause soil acidity problems.

There are no blanket answers to reduce nutrient loss. Each farm is different and will require a different response.

A property is more likely to be susceptible to nutrient loss where:

- Soil types are very heavy and there is surface run-off (worse when surface is cultivated);
- Soil types are sandy and there is high leaching;
- High fertiliser inputs are used;
- Crops have high irrigation requirement (more irrigations mean more chances to wash nutrients out);
- A flood or furrow irrigation system is used;

Horticulture Australia

Further references and

the end of this chapter

resources can be located at

# **2b Water quality**

Further references and resources can be located at the end of this chapter



# **2b Water quality**

- The property is located in a high rainfall area;
- The cultivated area is located close to a watercourse;
- Water is not well managed; and
- Time of nutrient application is not well managed.

Underlying most of the best practices is the need to keep nutrients in the plant root zone and to manage water to minimise irrigation run-off via the surface or into the groundwater.

Good nutrient management involves:

- Deciding what nutrients are needed, e.g. budget nutrients removed in the crop versus nutrients added in fertiliser applications;
- Applying fertilisers the right way;
- Minimising nutrient leaching to groundwater, especially nitrogen, by applying lighter than normal irrigation after fertiliser application or fertigating with lighter irrigations than normal; ensuring any in-line fertiliser injection systems have back flow prevention measures;
- Storing fertilisers properly; and
- Reducing possible harm to the environment by ensuring broadcast application of fertilisers involves leaving a buffer (no fertiliser) zone between the crop and sensitive areas such as watercourses and native vegetation.

For more information on selection and application of fertilisers see Chapter 4 – Nutrient management.

Correct storage and application of fertilisers will reduce environmental harm. Controlling soil erosion and reducing run-off and sediment loss will assist with loss of nutrients from target areas.

#### Agricultural chemical management

Agricultural chemicals can contaminate waterways through inappropriate application and storage.

Agricultural chemicals should not be applied where they could drift onto water, unless they are specifically approved for use in or near water. Make sure there is a margin between where the spray falls and the bank of any watercourse. For some chemicals, a minimum width for the no-spray zone is specified on the label.

Storage of agricultural chemicals, disposal of waste agricultural chemicals and empty containers must be undertaken with care.

For more information see Chapter 3 – Chemical management.

### Prevent pollution from fuels and oils

Oil and fuel spills can pollute waterways and soils, and are a major threat to flora and fauna. State legislation and environmental protection authorities treat the matter very seriously. Theft, vandalism and accidental damage by moving vehicles can cause oil spills and should be guarded against. For more information see Chapter 3 – Chemical management.

### Packaging shed water

Some packing sheds use large amounts of water as part of the packing process. Steps should be taken to ensure used water is safe to release back into waterways. This can be achieved through regular monitoring and if necessary filtering or treating water to remove organic material and chemicals. Organic material in water affects the amount of oxygen available and can have significant impact on fish and other aquatic life.

#### Organic matter

Septic tanks, and manure storage and waste produce dumping areas should be located well away from waterways and from water sources such as bores and dams. Run-off containing leachates from manure storage heaps and dumped piles of waste produce should be contained to prevent entry into waterways.

Ensure sewerage and septic systems are regularly maintained to prevent leakages into surface or groundwater.



### Monitoring and recording

Where there is a risk of poor quality irrigation water, testing should be undertaken regularly and at times of greatest risk. For instance, water should be tested for agricultural chemicals (on farm and off farm) when spraying has taken place near the water body. Copies of water tests should be maintained to track changes over time.

Testing parameters will vary with each situation, however the most common tests will be for pH, pesticides (as above), key nutrients (nitrates and phosphates), electrical conductivity (EC) to test salinity, and biological oxygen deficit (BOD) to test for organic matter presence and its potential effect on aquatic species. Accredited commercial testing laboratories are available Australia-wide. Check with your local council or State natural resources or primary industries department for further information.

A regular nutrient stocktake is a cost-effective way of checking fertiliser stocks, storage facilities, purchases and usage. The record should include storage location, type and amount of fertiliser. You may wish to also include details of dates into and out of store, and link usage to the fertiliser application records. A regular chemical stocktake should also be done. The farm map should indicate fertiliser and chemical storage locations and nearest watercourses. Records of weather conditions when applying agricultural chemicals can be used to substantiate minimisation of spray drift.

Records for disposal of unregistered chemicals and chemical containers can also be useful.

Monitoring the use of fuel and oils can detect potential threats to water quality caused by wastage, spills or leaks.

Monitoring any drainage lines for run-off water quality (e.g. nitrate test strips and turbidity tests) is also recommended. This can indicate areas and location of nutrient and soil loss.



### References and further resources

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

# **2b Water quality**

Further references and resources can be located at the end of this chapter.





Objective - to manage wastewater and run-off appropriately to minimise or eliminate potential negative environmental impacts

Appropriately managing all waste and run-off water from production areas is important to minimise the release of polluted wastewater into the environment. This is particularly so for containerised nursery operations, where leaching of irrigation water (and rainfall) from containers, and consequent nutrient losses in run-off water, is difficult to avoid. Collecting and recycling it as irrigation water or re-using it in non-production areas such as lawns, gardens and windbreaks also saves on water and fertiliser use, further reducing costs. Wastewater from packing sheds is also addressed in Topic 2b – Water quality and managing water from chemical dips is addressed in Chapter 3 - Chemical management.





### Review checklist

To go straight to the worksheet for this chapter click here.



### Relevant legislation and regulation

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

An effective re-cycling/re-use system involves the following activities.

#### Retain water onsite

Plan layout for drainage and prepare drains to collect run-off water and deliver to a collection dam or tank. Follow local authority or industry guidelines for designing drainage and storage capacity.

Set up filters and treatment systems to remove sediment, litter and undesirable chemicals from run-off water.

Collect run-off water into storage dams/tanks. Storage dams/tanks need to be specially constructed for holding run-off water. Natural wetlands on your property should not be considered as storage for run-off water due to potential pollutants in the water affecting the wetland. Follow local authority or industry guidelines for storage capacity and construction.

Monitor the quality of run-off water to determine effectiveness of treatments. Monitor:

- pH;
- Nutrient levels (particularly nitrates and phosphates); and
- Electrical conductivity (EC), which is an indicator of total salts in the water.

### Minimise discharge

Where possible, to minimize the effect of discharge, intercept and divert water from outside your property away from the production area. This reduces the volume of water to be managed and stored, and prevents pollutants and diseases being transferred.

### Monitor the quality of discharge water as above.

Seal production areas, if applicable, to minimise infiltration of water into the soil and to direct run-off water to drains. In nurseries, 200 µm thick plastic is commonly laid with at least 75 mm depth of 10-25 mm diameter gravel on top. Other options are concrete or bitumen. Level the growing beds to a minimum grade of 1:70 to ensure movement of run-off water to drains.

Further references and resources can be located at the end of this chapter



Where applicable, risks associated with pathogens need to be managed (e.g. disinfestation). This is applicable for the nursery industry - see NIASA Best Management. Practice Guidelines section 1.1.1.

# **2c Managing wastewater**

### Disinfect run-off water to remove pathogens before recycling or reusing

#### Further references and resources can be located at the end of this chapter



# 2c Managing wastewater

A useful reference that provides detailed information about setting up a recycling/re-use system is Managing water in plant nurseries - a guide to irrigation, drainage, and water recycling in containerised plant nurseries, 2nd Edition (2002), by NSW Agriculture (now NSW Department of Primary Industries).



#### Monitoring and recording

Regularly monitor run-off water to determine effectiveness of treatments. Monitor:

- pH;
- Nutrient levels (particularly nitrates); and
- Electrical conductivity (EC), which is an indicator of total salts in the water.

Keep a record of run-off water monitoring results.



### **References and further resources**

For access to relevant references and further resources click here.

# Water management chapter - references and further resources

### Water management chapter - references and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www.horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

#### NATIONAL ORGANISATIONS

Cooperative Research Centre for Irrigation Futures http://www.irrigationfutures.org.au This website is an archive of irrigation research, education and training activities between 2003-2010 (active to 2015).

CSIRO Water for a Healthy Country Flagship

http://www.csiro.au/Organisation-Structure/Flagships/Water-for-a-Healthy-Country-Flagship/WIRADA\_WFHC\_ ResearchProfile.aspx

This Flagship provides science and technologies that improve the social, economic and environmental outcomes from water.

Growcom Water for Profit http://www.growcom.com.au/ This website provides access to irrigation fact sheets.

Healthy Waterways http://www.healthywaterways.org/HealthyWaterways/Home.aspx This website is managed by a not-for-profit community organization focused on protecting and improving waterways in Australia.

Irrigation Association of Australia Ltd www.irrigation.org.au This is the website for the peak body for urban and rural irrigation industry.

Murray Darling Basin Authority http://www.mdba.gov.au The Authority provides basin-wide strategy, policies and planning. This website provides links to the Water Quality and Salinity Management Plan, Sustainable Rivers Audit and other publications.

National Program for Sustainable Irrigation (NPSI) http://www.npsi.gov.au NPSI was the national research program for irrigated agriculture. This program was completed June 2012, but website provides access to research publications.

#### See also NPSI Knowledge Harvest - Irrigation Essentials

NPSI (2012) Irrigation Essentials Updated – Research and innovation for Australian irrigators. National Program for Sustainable Irrigation. Cotton RDC, NSW. http://npsi.gov.au/files/products/national-program-sustainable-irrigation/ npsi06121/npsi06121-irrigation-essentials-updated.pdf

National Water Commission http://www.nwc.gov.au/home The Commission drives national water reform under the Federal Governments National Water Initiative.

Recycled Water in Australia www.recycledwater.com.au

The national initiative for recycled water in horticulture and wider community. This program was decommissioned, but this website still provides access to relevant information on use of recycled water in horticulture

STATE-SPECIFIC INFORMATION

New South Wales NSW Department of Primary Industries - Water and irrigation http://www.dpi.nsw.gov.au/agriculture/resources/ water

Northern Territory Government Department of Land Resource Management – Water http://www.lrm.nt.gov.au/water#.UnjLKaVtx-U

#### QLD

Queensland Government - Rural water use efficiency http://www.dnrm.qld.gov.au/water/access/rural-water-useefficiency

Growcom - Water for Profit initiative - Resource centre http://www.growcom.com.au/land-water/water-for-profit/ resources-water-for-profit/



2.22 Chapter 2 Water management



# Water management chapter - references and further resources

#### South Australia

Government of South Australia http://www.sardi.sa.gov.au/water/irrigation\_management/irrigation\_efficiency

#### Tasmania

Department of Primary Industries, Water and the Environment, Tasmania, (2012) Wise Watering Irrigation Management Course www.dpiwe.tas.gov.au/inter.nsf/WebPages/JMUY-5FJVP7?open

#### Victoria

Department of Environment and Primary Industries, Victoria http://www.depi.vic.gov.au/water/rural-water-andirrigation

#### Western Australia

Department of Agriculture and Food, WA https://www.agric.wa.gov.au/climate-land-water/water/water-management

#### **CHAPTER - SPECIFIC REFERENCES AND RESOURCES**

#### 2a Irrigation efficiency

Boland, AM., Ziehrl, A. and Beaumont, J. (2002) Guide to Best Practice in Water Management-Orchard Crops. Department of Natural Resources and Environment, Victoria. http://catalogue.nla.gov.au/Record/691259 or http:// apal.org.au/wp-content/uploads/2013/07/fo-ow-handout-09-sept-best-practice-water-mgmt-boland.pdf

Boland, AM. (2008) Guidelines for irrigation management (HAL Project Number AP06055). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Boland, AM. (2010) Refinement of Guidelines for Avocado Irrigation Management Under Australian Conditions for Different Phenological Stages (HAL Project Number: AV08021). Horticulture Australia Ltd, Sydney, NSW. www. horticulture.com.au

Brebbia, C.A.; Marinov, A.N.; Bjornlund, H (2010) Sustainable Irrigation Management, Technologies and Policies III, WIT Press, UK. http://trove.nla.gov.au/version/48873367

#### Bureau of Meteorology www.bom.gov.au

Charlesworth, P. (2005) Irrigation Insights - Soil Water Monitoring (second edition). Land and Water Australia, Irrigation Insights No.1, Canberra. http://www.irrigationfutures.org.au/imagesDB/news/soilwatermonitoring2ed.pdf

CSIRO FullStop - FullStop Wetting Front Detector helps you to "see" what is happening down in the root zone when you irrigate the soil. www.fullstop.com.au

Cresswell, G (2002) Water retention efficiency of potting mixes. Nursery Papers Issue 7, 2002, Nursery & Garden Industry Australia, Sydney. http://www.ngia.com.au/Story?Action=View&Story\_id=1237

Gordon L. J., Finlayson C. M. & Falkenmark M. (2010) Managing water in agriculture for food production and other ecosystem services. Agricultural Water Management 97, 512-9.

Henderson, C (2011) Driving better vegetable irrigation through profitable practice change (HAL Project Number: VG07023). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au Hickey, Hoogers, Singh, Christen, Henderson, Ashcroft, Top, O'Donnell, Sylvia & Hoffmann (2006) Maximising returns

from water in the Australian vegetable industry (multiple reports), NSW Department of Primary Industries, NSW. http://www.dpi.nsw.gov.au/agriculture/horticulture/vegetables/industry/maximising-returns-from-water

Jensen, ME (ed.) (1983) Design and operation of farm irrigation systems, ASAE Monograph 3, American Society of Agricultural Engineers, St Joseph, Michigan.

Jobling, J; Rogers, G; Anderson, A; and Vicic, A. (2008) Managing Water for Yield and Profit: A training guide for Irrigators in the Australian Vegetable Industry, Arris. http://www.horticulture.com.au/librarymanager/libs/165/ Managing\_Water\_Yield\_Profit.pdf

Kriedemann, P. and Goodwin, I. (Updated in 2008) Irrigation Insights – Regulated deficit irrigation and partial rootzone drying. Land and Water Australia, Irrigation Insights No. 4. (Ed. Anne Currey), Canberra. http://npsi.gov.au/files/products/national-program-sustainable-irrigation/pr020382/pr020382.pdf

NSW Department of Agriculture (2003) Best management guidelines for irrigated vegetable crops (CD). http://www.dpi.nsw.gov.au/agriculture/resources/soils/salinity/crops/saline-irrigation

Nursery Papers Technical (September 2013 Issue no.78) - Automating irrigation scheduling in nursery Production. Nursery & Garden Industry Australia, Sydney. http://www.ngia.com.au/Story?Action=View&Story\_id=2192 Mitchell, P. and Goodwin, I. (1996). Irrigation of vines & fruit trees. Agmedia Pubs.

Rolfe, C, Yiasoumi, W and Keskula, E (2002) Managing water in plant nurseries: A guide to irrigation, drainage and water recycling in containerised plant nurseries: NSW Department of Agriculture, Orange, NSW. http://www.publish.csiro.au/pid/2943.htm (for purchase via this link)

Tee, E., Burrows, D., Boland, A-M and Putland, S. (2004) Best Irrigation Management Practices for Viticulture in the Murray Darling Basin. Cooperative Research Centre for Viticulture. http://www.crcv.com.au/resources/Irrigation%20and%20 Water/Additional%20Resources/MDB%20Irrigation%20Booklet.pdf

Wise Water - Irrigation Management Course 2001 - course notes http://www.dpiw.tas.gov.au/internnsf/Attachments/ JMUY-5FP77U/\$FILE/IntroductiontoWiseWatering.pdf

#### 2b Water quality

### Caring for Our Country - Reef Rescue program

http://www.nrm.gov.au/funding/reef-rescue/components.html This initiative supports projects in the areas of water quality, water quality monitoring and reporting, crown of thorns starfish, systems repair, the land-sea country partnership and the Great Barrier Reef Marine Park Authority.

Department of the Environment and Heritage (now Australian Government Department of the Environment) (Version 1 -2002) Water Quality Targets: A handbook, Environment Australia. http://www.environment.gov.au/resource/water-qualitytargets-handbook or http://nrmonline.nrm.gov.au/downloads/mgl:383/PDF

Department of Primary Industries, Water and the Environment, Tasmania, Water Quality Monitorin www.dpiwe.tas.gov.au/inter.nsf/WebPages/LBUN-4YH4AK?open

Great Barrier Reef Marine Park Authority (Revised Edition 2010) Water Quality Guidelines for the Great Barrier Reef Marine Park, GBRMPA, Townsville. http://www.gbrmpa.gov.au/\_\_data/assets/pdf\_file/0017/4526/GBRMPA\_ WQualityGuidelinesGBRMP\_RevEdition\_2010.pdf

Jarwal, S; Boland, AM; Stevens, D; and Faggian, R (2006) Using recycled water in horticulture: A growers guide, Department of Primary Industries Victoria. http://www.recycledwater.com.au/uploads/File/documents/Growers%20 Guide%20web.pdf

#### Landcare Australia www.landcareonline.com

Lovett, S., Price, P. & Edgar, B. (eds) (2007), Salt, Nutrient, Sediment and Interactions: Findings from the National River Contaminants Program, Land & Water Australia, 2007. http://lwa.gov.au/files/products/river-landscapes/pk071328/ pk071328.pdf

National Water Quality Management Strategy (NWQMS) (2000) Australian and New Zealand guidelines for fresh and marine water quality: Volume 3 - primary industries http://www.environment.gov.au/resource/australian-and-new-zealandguidelines-fresh-and-marine-water-quality-volume-1-guidelines (note that the guidelines are currently [2013] under review and next edition due to be released in June 2014).

NSW Department of Primary Industries – Water quality and testing http://www.dpi.nsw.gov.au/agriculture/resources/ water/quality

Queensland Government - ReefWise Farming initiative http://www.qld.gov.au/environment/agriculture/sustainablefarming/reef/

Recycled water in Australia www.recycledwater.com.au (decommissioned but still provides access to relevant information on use of recycled water in horticulture).

Schrale, G; Biswass, T, (2009). Salinity management practice guidelines: Managing root-zone salinity for irrigated horticultural crops in winter rainfall zones of Australia, National Program for Sustainable Irrigation, Land and Water Australia. http://npsi.gov.au/products/pn22225

Stevens, D; Smolenaars, S; Kelly, J (2008). Irrigation of amenity horticulture with recycled water, Horticulture Australia Ltd http://www.recycledwater.com.au/uploads/Amenity%20Hort%20HB%20Recycled%20Water%20(reprint).pdf

Waterwatch Australia – decommissioned May 2013 – archived at http://nrmonline.nrm.gov.au/downloads/mgl:2873/ PDF Waterwatch Australia (2005). Waterwatch Australia National Technical Manual, Department of the Environment and Heritage, ACT. The following modules are available (as separate stand alone documents) in the Manual:

- 1. Background http://nrmonline.nrm.gov.au/downloads/mgl:2882/PDF
- 2. Getting Started: the team, monitoring plan and site http://nrmonline.nrm.gov.au/catalog/mgl:2871
- 3. Biological Parameters http://nrmonline.nrm.gov.au/downloads/mgl:2883/PDF
- 4. Physical and Chemical Parameters http://nrmonline.nrm.gov.au/downloads/mgl:2880/PDF
- 5. Data to Information to Action (could not be sourced online)
- 6. Groundwater Monitoring http://nrmonline.nrm.gov.au/downloads/mgl:2875/PDF 7. Estuarine Monitoring http://nrmonline.nrm.gov.au/downloads/mql:2869/PDF
- Horticulture Australia

Chapter 2 Water management

2.24



#### 2c Wastewater

Badgery-Parker, J, (2003). Managing wastewater with a wetland. NSW Agriculture (now NSW Department of Primary Industries). Fact sheet - http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0005/119372/horticulture-waste-water-wetland-system-eng.pdf

EPA South Australia (2009). Wastewater irrigation management plan (WIMP) - a drafting guide for wastewater irrigators, EPA, SA. http://www.epa.sa.gov.au/xstd\_files/Waste/Guideline/guide\_wimp.pdf

NGIA (2013). Nursery Industry Accreditation Scheme Australia (NIASA) Best Management Practice Guidelines – 5th Edition, NGIA, Sydney. https://www.ngia.com.au/Category?Action=View&Category\_id=135 (for purchase via this link).

NSW Agriculture (2002). Managing water in plant nurseries – a guide to irrigation, drainage, and water recycling in containerised plant nurseries, 2nd Edition. http://www.publish.csiro.au/pid/2943.htm (for purchase via this link).

The Nursery Papers 2003:7 – Designing a nursery reed bed, NGIA, Sydney. https://www.ngia.com.au/ Story?Action=View&Story\_id=1221

The Nursery Papers, 1996:4 – Monitoring and managing recycled water quality in nurseries. NGIA, Sydney. https://www.ngia.com.au/files/nurserypapers/NP\_1996\_04.pdf

Nursery Industry Water Management Best Management Practice Guidelines (updated 2005), Nursery Industry Association of Australia, Sydney. http://www.horticulture.com.au/librarymanager/libs/172/Water%20 Management%20BP%20guidelines%20-%20nursery%20industry.pdf

Rolfe, C and McPhee, G (1999) WaterWork: A competency-based homestudy and workshop package version 2.1 on CD-ROM, produced by Australian Horticultural Training, published by Nursery & Garden Industry Australia and NSW Agriculture. http://www.ngia.com.au/files/nurserypapers/NP\_1996\_10.pdf

SAI Platform (2010) Water Conservation Technical Brief: TB7 – Wastewater Use in Agriculture, Sustainable Agriculture Initiative Platform. http://www.saiplatform.org/uploads/Library/Technical%20Brief%207.%20Wastewater%20use%20 in%20agriculture.pdf





# Chapter 3 Chemical management



Objective – agricultural chemicals are used in accordance with label or permit instructions; and all chemicals, including fuels and oils, are stored, handled, applied and disposed of in a manner that minimises environmental impacts

Agricultural chemicals are by nature potentially dangerous to humans, flora and fauna and ecosystems.

Pesticides can have serious effects on natural ecosystems if they move off-site via water, air or soil. Of particular concern is the effect of pesticide residues on sensitive neighbouring or downstream ecosystems such as wetlands, freshwater and marine habitats, and national parks and reserves.

Spray drift, in particular, is a potential source of friction between farmers and their neighbours. Spray drift can also cause much damage to wildlife. Liability for damage, illness or injury caused by spray drift is also becoming a serious issue.

Chemicals other than pesticides are widely used for cleaning and sanitising around growing and production sites and for treating water. Care needs to be taken to ensure these chemicals do not enter waterways and drains or accidentally spill on to soils and vegetation.

To minimise harm to the environment, all aspects of chemical use, from justification of the need for chemical intervention to storage, handling and disposal of empty containers, need to be considered.

Further references and resources can be located at the end of this chapter.

# **3** Chemical management





Spray drift is a potential source of friction between farmers and their neighbours. Spray drift can also harm wildlife. Liability for damage, illness or injury caused by spray drift is becoming a serious issue.

Spray drift is caused by a combination of factors, including:

- Wind speed;
- Instable local atmospheric conditions;
- Wrong nozzles or pressure choice affecting spray quality;
- Vehicle speed;
- Boom height; and
- Poor equipment maintenance, including incorrect equipment setting.

Horticulture Australia

Further references and

the end of this chapter.

HAL

resources can be located at

Further references and resources can be located at the end of this chapter.





### **Review checklist**

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

### Minimise application

Pesticides are only one strategy for controlling pest insects, diseases and weeds. Integrated Pest Management (IPM) involves a range of strategies:

- Training in recognition of the pests and diseases that can attack your particular crops, their symptoms of attack and life cycles;
- Regular monitoring of crops for pests, diseases and weeds;
- Only using pesticides if pest numbers exceed threshold levels, and consider using:
- environmentally friendly pesticides, such as oils, soaps and biological control agents such as bacillus formulations;
- narrow-spectrum pesticides instead of broad-spectrum pesticides;
- spot applications of pesticide instead of blanket sprays;
- strategic application when the pest or disease is most vulnerable; and
- resistance minimisation strategies;
- Practicing good hygiene to limit disease in particular;
- Having an all-year-round weed management program in place, both in and around the growing area - weeds can harbour pests and diseases and act as a constant source of reinfestation (although weeds may also be a refuge for natural predators);
- Maintaining good soil health, including an open, well aerated structure, high organic matter levels and a diverse and active soil biology, which in turn promotes healthy crops that are more resistant to disease and pests; and
- Encouraging natural predators. See Chapter 5 Biodiversity.

Further information about IPM is available from State departments of agriculture/ primary industries. Laws relating to chemical storage and use change frequently. Check with your State authority for the most up-to-date information.

#### Safe storage

Agricultural chemicals can contaminate watercourses if not stored appropriately. Any new chemical storage should meet the highest standards of design and construction. Existing chemical sheds may need to be improved.

Chemical sheds should not be built where there is a risk of polluting watercourses or groundwater, or in areas subject to flooding.

All farm chemicals must be stored in a secure storage area that meets local regulations/ legislation. Farm chemical safety and handling training courses address chemical storage issues as do Managing Farm Safety courses. It is recommended that growers obtain specific local advice as each state and, in some cases, each local government area may have particular requirements.

In general, when storing agricultural chemicals use the following guidelines:

- Store pesticides and chemicals away from residences and other occupied buildings;
- Store pesticides and chemicals in a lockable, weatherproof, fireproof, well-ventilated area away from production facilities, waterways, water supplies and flood-prone areas;
- Keep the storage area clear of extraneous or combustible waste materials and control ignition;
- Ensure adequate lighting for a safe working environment;
- The floor should be impermeable and easy to clean, and the storage area should be able to contain spills. Spilt liquid can be contained by bunding (an embankment or wall);
- Maintain a chemical spill cleanup kit near the area (see 'Dealing with spills' later in this topic). Store pesticides and chemicals in their original containers;
- Store pesticides separate from fertilisers and chlorine. Store liquid pesticides and chemicals below powders. Maintain an up-to-date inventory of stored pesticides;
- Store only enough pesticide on site to meet short/medium-term needs. Regularly check pesticide and chemical containers for any leakage or damage. Running water should be available;
- Chemical storage areas should be appropriately signed for the size and nature of the storage, including a 'No Smoking' sign;
- Consider a central storage point based with a local spray contractor or at the airstrip;

• Keep current Material Safety Data Sheets for all chemicals in use. All States have strict regulations concerning storage of pesticides on farm and business sites, including occupational health and safety requirements. Check with your local authority to ensure you conform to these regulations.

#### Safe transport

Ensure chemical containers are leak-proof and adequately secured when transporting on farm or between farms.

Observe safe handling practices when transporting pesticides and chemicals.

### Dealing with spills

It is a good idea to have an emergency plan in place to deal with spills of different groups of chemicals so you are prepared if it ever happens.

Know how to deal with spills and have spill kits close to storage and mixing areas.

Spill kits should include:

- A shovel;
- Dustless absorbent material, such as 'kitty litter', activated charcoal, vermiculite, hydrated lime, clay or earth and dry sand (avoid using sawdust or other combustible materials); and
- Containers to hold the absorbent material or other leaking containers.

3.4 Chapter 3 Chemical management

© 2014 – Horticulture Australia Limited

Further references and resources can be located at the end of this chapter



## **3** Chemical management

For more information on appropriate signage for chemical sheds refer to the Workplace Health and Safety agency in your State.

Further references and resources can be located at the end of this chapter



**Chapter 3** Chemical management 3.5

#### Also:

- Wear protective clothing, including gloves, respirator, boots and eye protection when dealing with spills;
- Properly dispose of the chemical-drenched absorbent clean-up material;
- Use hydrated lime or bleach to decontaminate spill surfaces, but never use these two materials together;
- Report the spill to authorities if it is large or if it enters waterways.

#### Mixing and application

#### Responsible use of pesticides and chemicals

Ensure at least one person in the business has completed an accredited chemical user's training course (similar or equivalent to ChemCert) and ensure all staff that apply pesticides have adequate training. Regularly calibrate and maintain pesticide application equipment.

Only use pesticides that are registered or permitted by the Australian Pesticides and Veterinary Medicines Authority (APVMA) or otherwise authorised for use in your State for the particular:

- Crop;
- Application timing;
- Method of application; and/or
- Tank mix.

Obtain and keep copies of any APVMA permits that are relevant to your spraying.

NSW requires all people who apply chemicals in the business to have accredited training.

Growers in Victoria do not require APVMA off-label permits for certain off-label uses. Visit the VIC DEPI Chemical Use website for more details, http://www.depi.vic.gov.au/ agriculture-and-food/farm-management/chemical-use

WA Health (Pesticide) Regulation 87 allows an agricultural user of pesticides in WA to use a pesticide registered for use on a crop against any pest/disease/weed on that crop provided the label directions are followed.

Always follow label and permit instructions. Seek professional advice if you have concerns.

Keep Material Safety Data Sheets (MSDSs) for all pesticides used close to areas where chemicals are handled and consider placing copies in tractors used with spray equipment. These provide information about health hazards and safe handling, including transport, storage and spill clean-up.

In case of poisoning or exposure to pesticides, contact the Poisons Information Centre on 131 126. This s a 24-hour service.

#### Mixing

Make sure to site the mixing and washdown area away from water sources, drains and streams. The area should be constructed to contain spills for collection and disposal.

Never leave a spray unit unattended while filling.

As some drums are difficult to pour, use a specifically-designed drum pourer to minimise spillage. Triple-rinse empty chemical containers and mixing equipment back into the vat.

#### Minimising spray drift

There are many strategies to minimise or prevent the chances of spray drift, starting with how you establish new horticultural sites.

Managing spray drift should be included in property management plans and specific spray plans can be developed which include identification of sensitive areas and options for minimising spray drift into those areas.

You should also consider the following strategies:

- Check the weather forecast before starting off. Do not spray if the wind direction and speed would cause the spray to drift on to sensitive areas or neighbours (see Weather Conditions topic below). Under light wind conditions, wind direction is often variable and may result in unpredictable off-target movement;
- Avoid spraying on hot days (>30°C) or dry days (<40% relative humidity) as these conditions can increase the rate of evaporation of water-based sprays and may subsequently increase spray drift;
- Spray when winds are most consistent, generally early morning, early evening or at night. Select the right combination of spray unit, nozzle type and size, and pressure;
- Use the largest appropriate spray droplet size (except if using CDA equipment);
- When using a boom spray, keep the boom as low as possible, consistent with an even spray pattern at the correct target height. Check spray angles and adjust the height accordingly. Reduce the operating pressure and forward speed but maintain the dose, volume and spray quality within recommendations on the label;
- Consider not treating the boom-width or part of the boom-width closest to the boundary. This can also provide a useful tool to check spray efficiency and to help maintain beneficial insects. Modify spray equipment to help minimise spray drift. These modifications include low drift nozzles, shielded and covered spray booms and air-assisted spray equipment;
- For air-blast sprayers pay particular attention to weather conditions, good set-up and maintenance of equipment and barriers/buffer distances from neighbours;
- For aerial spraying, ensure that the spray contractor is aware of sensitive areas/ neighbours and of their responsibility for any off-target application or spray drift;
- Ensure operators have appropriate training/competencies and relevant licences;
- Erect or plant barriers to catch possible spray drift, and establish buffer zones between production areas and neighbours or sensitive natural areas, such as wetlands and waterways (see Buffer zones later in this topic).

#### Weather conditions

Wind speed in the spray release zone is an important factor in determining spray drift. Meteorological measurement of wind speed is taken 10 m above ground, so care is needed in interpreting weather advice and actual wind speed at nozzle height.

Prior to spraying, an assessment should be made of the wind speed and direction by using a simple wind speed meter or by watching the movement of plants, trees and clouds. See Table 3.1 below. This monitoring should continue during application.

Table 3.1: Wind speed guide. Source: Lovell (2006).

Approx. wind speed at boom height	Description	Visible signs	Spra
Less than 2 km/hr	Calm	Smoke rises vertically	Not ′coa
2 - 5 km/hr	Light air	Direction shown by smoke drift	Acce
6 - 11 km/hr	Light breeze	Leaves rustle, wind felt on face	Idea
12 - 19 km/hr	Gentle breeze	Leaves and twigs in constant motion	Incre
20 - 28 km/hr	Moderate breeze	Small branches moved, raises dust or loose paper	Spra

Further references and resources can be located at the end of this chapter.



Horticulture Australia

## **3** Chemical management

- t recommended. Use only 'medium' or arse' spray quality
- ceptable spraying conditions
- al spraying conditions
- reased risk of spray drift, take special care
- raying inadvisable

Further references and resources can be located at the end of this chapter



Delta-T calculations can be used as indicators of acceptable spray conditions. It is indicative of evaporation rate and droplet lifetime. When applying pesticides, Delta-T should ideally be between 2 and 8. For more information see Weather for Pesticide Spraying, Bureau of Meteorology or ChemCert website.

#### **Buffer zones**

Buffer zones are a very good way to reduce the risk of spray drift. Buffer zones can be established by planting trees and shrubs downwind of a production area. Vegetation that is tall, rough and thin is better at catching droplets than vegetation that is short, smooth and thick.

A droplet-catching barrier should be less than 50% porous, with a height about 1.5 times the height of spray release. Use multiple vegetation layers as a screen rather than a single layer. Vegetable growers should also consider establishing intercrop buffers such as rye corn or sweet corn as barriers.

#### **Protecting water supplies**

Ensure pesticide cannot be back-syphoned into the water supply when filling spray tanks by installing an anti backflow device or pumping from a separate tank filled from the main water source.

Fill spray tanks with water and then move them away from waterways or water storages before adding chemicals.

Minimise movement of pesticides into stormwater or wastewater run-off by avoiding application of pesticides immediately before rain or irrigation, unless stated otherwise on the APVMA product label or permit.

#### **Consider community relations**

Disputes involving environmental nuisance (for example issues related to application of agricultural chemicals, noise or dust) can lead to a breakdown of good neighbourly relations.

The best way to avoid problems is to attempt to get on well with your neighbours. Having a 'good neighbour' policy and discussing aspects of farming with neighbours is one way to achieve this. Neighbours, particularly non-farming people, need to recognise that primary producers make their living through agricultural activities and that these activities are an important part of the economy and food chain.

Primary producers need to recognise that some activities can negatively impact on their neighbours and that at times it may be appropriate to adjust activities as far as reasonable to minimise the impact.

#### **Disposal of pesticide containers**

Under various State regulations, businesses are required to dispose of empty chemical containers safely. When purchasing, ask if used pesticide containers can be reused, returned, refilled or recycled.

Un-rinsed containers can hold as much as 3% of product concentrate. This means they can present a hazard to people who handle them and have potential to contaminate the environment.

Used containers that cannot be returned or recycled should be triple rinsed or pressure rinsed immediately after emptying the container as residues are more difficult to remove when they are dry. This is done by filling the container with clean water to approximately a quarter of capacity, replacing the cap, shaking and then adding the wastewater to the spray tank. This is repeated three times. Other disposal methods

(e.g. pumping to sump or limed disposal pit) are not acceptable. Pressure rinsing is also an option and special equipment is available.

Puncture steel containers after rinsing so that they cannot be re-used. Pass a steel rod or crowbar through the neck/pouring opening and out the base of the container. Do not puncture plastic 20-litre containers included in reconditioning/reuse programs. Empty pesticide containers must be stored in a designated, secure area (preferably locked), and disposed of either through a controlled approved disposal scheme, or according to a documented procedure that meets state or territory regulations. Access to this area must be restricted for both people and animals.

Ideally, property managers should be involved in any disposal scheme to remove rinsed chemical containers, and/or use the services of a licensed waste collector to remove unwanted pesticides.

The drumMUSTER scheme operates in all states. From January 2004, only containers carrying the eligible container logo are collected under the drumMUSTER program. The drumMUSTER website (www.drummuster.com.au) contains details of collection days and locations. It also includes contact details for regional field officers. Pending disposal via drumMUSTER or other approved disposal methods, containers must be rinsed and stored in a separate secure area.

### Disposal of surplus spray and washings

Avoid leftover pesticide by carefully calculating how much is needed for the area to be sprayed.

Do not allow leftover spray, rinsings from a spray tank or from empty pesticide or chemical containers to enter streams or drainage from the property. Make sure that any disposal method you use is safe for your chemical waste, location and circumstances as incorrect disposal can result in prosecution. Check your local laws regarding use and management of farm chemicals.

Disposal methods may include:

- Storing rinsate or surplus spray in an appropriately labelled container and use to make up the next compatible spray mix;
- Diluting rinsate/surplus spray and spray on to target crop in a manner that will not exceed label rates or wash off chemical previously applied;
- Spraying leftover pesticide and washings from rinsing after spraying on to an area of ground away from where people will be and from drains, low drainage areas, waterways and water storages (follow label guidelines);
- Emptying into a lime-filled pit (obtain advice as to quantities of lime and appropriate sites before using this method); or
- Consider using enzyme products (new technology enzyme products are capable of almost complete breakdown of organophosphate insecticides, either in the spray tank or in a holding tank, and provide a useful option for rapid chemical clean-up).

Post-harvest dips may also be treated with lime to deactivate the agricultural chemical. Again specific advice should be sought regarding the most appropriate disposal practices. Information may be available on the chemical label and this should be checked first.

Spray equipment should be filled and washed in an area chosen and established for that purpose. Spillages and rinsings should not be able to escape from the area. Ensure the area is well away from watercourses and dams.

Remember, damage to the environment such as fish kills and the like are prosecutable offences and should be avoided.

Further references and resources can be located at the end of this chapter.



## **3** Chemical management

Further references and resources can be located at the end of this chapter



#### Disposal of old, de-registered or unwanted pesticide concentrates

Unwanted chemicals, such as those that are no longer registered for use, should not be stored on farm for longer than is necessary to arrange for their disposal.

Make sure these chemicals are appropriately stored to prevent misuse. Storage in the chemical shed is recommended as long as the chemicals are clearly identified as not for use and, preferably, are segregated.

#### Ensure the containers do not leak

One option for disposal of unwanted agricultural chemicals is ChemClear<sup>®</sup>. The ChemClear<sup>®</sup> program has a web-based booking system where growers can register chemicals for collection. There are two categories of chemicals – Group 1 and Group 2. Group 1 chemicals are collected free of charge, while disposal of Group 2 chemicals attracts a fee. For more details visit www.chemclear.com.au or call 1800 008 182.

Alternatively, a certified or approved chemical waste contractor or supply company can be used. If transporting these pesticides to a collection centre, place them on the back of a utility or truck, never in the boot or cabin of a vehicle, or back of a station wagon, where fumes may affect the driver or passengers. Ensure containers cannot leak during transport.

Contact your local council or waste management authority for advice on methods of waste disposal available in your local area. See Chapter 6 - Waste management for information on controlled/ prescribed wastes.

#### Use and disposal of other chemical products

If rat and mouse baits are used, ensure they are enclosed in bait stations to prevent native birds and animals eating them. Dispose of used rodenticides or other pesticide baits, as well as carcasses, in accordance with the product label. If carcasses are being buried and the label does not give any special instructions, take care to bury them so that there is no risk of polluting surface or groundwater, and where dogs or native animals will not dig them up. Some baits have been developed that do not cause secondary poisoning.

Dispose of contaminated wastes, such as protective clothing and materials used to deal with spillages in accordance with local regulations. Some waste disposal sites can deal with this sort of waste, while others cannot. Contact your local council or waste management authority for advice on methods of waste disposal available in your area. See Chapter 6 – Waste management for information on controlled/prescribed wastes.

#### Storing and handling fuels and oils

Take reasonable steps to secure vulnerable tanks against interference; this may be as simple as locking pumps or taps. Bund above ground fuel tanks and provide some form of leakage protection for underground tanks. Materials for soaking up any spillages should be available at the storage area.

Check for leaks frequently and repair them promptly, especially with underground tanks (even slow leaks can have a major impact if allowed to continue).

If fuel tanks are bunded, all valves should be inside the bund and should still be closed and locked when not in use. Store flexible hoses for refuelling vehicles with the hose outlet in the bund.

Take water or fuel/oil out of the bund and dispose of it safely, e.g. by using a blanket that is specially made to absorb fuel/oil.

Consider installing an anti-siphon device if the inlet is lower than the highest fuel level of the tank.

Fuel storage facilities should be away from watercourses and with sufficient surrounding space to permit easy access, thus reducing the chance of accidental damage.

The risk of fire should be minimised. Mobile fuel tanks should:

- Be designed to protect them from accidental damage;
- Have a contents gauge and be stable enough to travel around the property; • Have all connections and valves, where fuel could empty by gravity, kept locked when not in use;
- Use with care, especially when near watercourses; and
- If possible, not be left near or uphill from a watercourse.

If fuel tanks are bunded, this should be done in accordance with the relevant State legislative requirements. For example, 125% of the largest container plus 25% of total volume.



### Monitoring and recording

Records that can (and in some cases must) be kept include:

- Staff training records;
- Pesticide application records (spray diary), including details of date and time of application, operator, location/area treated, pest/disease target, pesticide used, rate, application equipment and weather conditions (particularly to support miminisation of risk of spray drift); chemicals register or inventory;
- Waste disposal records (see Chapter 6 Waste management) such as:
- Chemclear<sup>®</sup> documentation, issued to prove chemicals have been booked in for collection and also when chemicals are collected, and
- Receipts and invoices from recycling or commercial disposal businesses;
- Disposal of surplus agricultural chemicals (can be recorded on spray records, particularly if sprayed back over the crop);
- Machinery maintenance;
- A farm map showing any buffer zones around paddocks, including sensitive areas and neighbours; and
- Ensure that your pesticide spraying records comply with your State or Territory regulatory requirements.



### **References and further resources**

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter.



3.10 Chapter 3 Chemical management

© 2014 – Horticulture Australia Limited

# **3** Chemical management

Further references and resources can be located at the end of this chapter



#### Chemical management - references and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www. horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

#### Aerial spraying www.aerialag.com.au (Aerial Agricultural Association of Australia Limited)

#### Australian Pesticides and Veterinary Medicines Association www.apvma.gov.au

Bianchi FJJA, Ives AR, Schellhorn NA. (2013). Interactions between conventional and organic farming for biocontrol services across the landscape. Ecological Applications 23:1531-1543. http://www.esajournals.org/ doi/abs/10.1890/12-1819.1 (for purchase via this link).

#### Bureau of Meterology (BOM) Weather for Pesticide Spraying www.bom.gov.au/info/leaflets/Pesticide-Spraying.pdf www.bom.gov.au/announcements/ag\_bulletins.shtml

#### Chemcert® - Chemical use training and accreditation www.chemcert.org.au

#### Collection programs:

ChemClear® - Collection of unwanted chemical or out of date chemical, handling and storage tips www.chemclear.com.au

DrumMUSTER® - Collection of empty chemical containers www.drummuster.com.au

CropLife Australia - resistance management - fungicides, herbicides and insecticides http://www.croplifeaustralia.org.au/industry-stewardship/resistance-management/

CSIRO (2002) Spray Drift Management - Principles, Strategies and Supporting Information, CSIRO Publishing. http://www.publish.csiro.au/Books/download.cfm?ID=3452

Spray Drift Management for purchase here www.publish.csiro.au/nid/22/pid/3452.htm

DEFRA (2011) Protecting our water, soil and air, Department of Environment Food and Rural Affairs, UK. https://www.gov.uk/government/publications/protecting-our-water-soil-and-air

Department of Agriculture, WA (revised 2007) Code of Practice for the use of Agricultural and Veterinary Chemicals in WA. Department of Agriculture, WA. http://www.agric.wa.gov.au/objtwr/imported\_assets/ content/pw/chem/cop\_bulletin.pdf

Health WA (2013) A guide to the use of pesticides in Western Australia. Department of Health, Government of Western Australia. http://www.public.health.wa.gov.au/3/1139/2/pesticide\_use.pm

Department of Environment and Primary Industries, Victoria – Chemical use in agriculture http://www.depi.vic.gov.au/agriculture-and-food/farm-management/chemical-use/agricultural-chemical-use

Department of Primary Industries, Fisheries and Mines, Northern Territory www.horticulture.nt.gov.au

Department of Primary Industries, Water and Environment, Tasmania - Guidelines for managing leftover spray mixes, rinsings and washings (Tasmania) www.dpiwe.tas.gov.au/inter.nsf/Attachments/CPAS-5U326X?open

Department of Primary Industries, Water and Environment Tasmania - On-farm Pesticide Storage www.dpiwe. tas.gov.au/inter.nsf/WebPages/TTAR-62Q5Y2?open

Huwer, R. (2007) Advancing integrated pest management in Macadamias & Towards adoption of IPM in macadamias (HAL Project Reference MC02048 & MC05005). NSW Department of Primary Industries (NSW DPI), NSW. Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

#### Legislation about pesticide use:

Search the databases under 'A' for Agriculture and Veterinary Chemicals

Australian Capitol Territory http://www.legislation.act.gov.au

New South Wales www.legislation.nsw.gov.au

Northern Territory www.legislation.nt.gov.au (or direct link: http://www.dcm.nt.gov.au/strong\_ servicedelivery/supporting\_government/current\_northern\_territory\_legislation\_database)

Queensland https://www.legislation.qld.gov.au/OQPChome.htm

South Australia http://www.legislation.sa.gov.au

Tasmania http://www.legislation.tas.gov.au

Western Australia www.legislation.wa.gov.au (or direct link http://www.slp.wa.gov.au/legislation/statutes. nsf/default html)

#### Victoria http://www.legislation.vic.gov.au

Llewellyn,R; Papacek, D; Altmann, J; Ryland, A; Seymour, J; Steiner, M; Wilson, C; Horne, P; Proctor, R; Thompson, J; and Grundy, P., (2002). The Good Bug Book, 2nd Edition. Integrated Pest Management Pty Ltd, Richmond, New South Wales.

Material Safety Data Sheets are available from manufacturer's websites or the following websites www.pestgenie.com.au or www.msds.com.au

Mattner, S. (2005) Identification of sustainable soil disinfestation options for the temperate Australian strawberry industry (HAL Project Reference BS01004). VIC Department of Primary Industries, VIC. Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

McCrystal, R. (2010) Improving the management of sweet potato soil insect pests (HAL Project Reference VG05037). Department of Employment, Economic Development & Innovation. Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Murray Darling Basin Commission (2004) Current recommended practice. A directory for broadacre dryland agriculture. See particularly the Integrated Pest Management section. http://www.mdba.gov.au/sites/default/ files/archived/mdbc-NRM-reports/2231\_Current\_recommended\_practice\_directory\_broadacre\_dryland\_ ag.pdf

National Chemical Reference Guide - Produced by the Australian Government Department of the Environment and Heritage. This online guide is the first resource of its kind for Australia. It brings together information from around the world, on a range of environmental standards and guidelines. The comprehensive, searchable database contains over 600 chemicals, encompassing various environmental standards for air, water, soil, sediment and biota (including plants). http://apps5a.ris.environment.gov.au/ pubgate/crg\_public/!CRGPPUBLIC.pStart

NSW Environment Protection Authority http://www.epa.nsw.gov.au/pesticides/Pesticides.htm

Nursery and Garden Industry, (2002) Australia Best practice manual for pesticide application in nursery and garden industry, NGIA, Sydney. http://www.ngia.com.au/Story?Action=View&Story\_id=1804

Oke, S. (2011) Advanced oxidation for the purification of horticulture produce, water and air (HAL Project Reference HG09028). Oztec Pty Ltd. Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Organic Pest Management information - Department of Primary Industries Victoria.

Porter, I. (2011) Best-practice IPM Strategies for Control of Major Soilborne Diseases of Vegetable Crops throughout Australia (HAL Project Reference VG07125). Victorian Department of Primary Industries (VICDPI). Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Safe Work Australia - Hazardous Chemicals - Safety Data Sets http://www.safeworkaustralia.gov.au/sites/ swa/whs-information/hazardous-chemicals/pages/hazardous-chemicals-other-substances

Schellhorn NA, Nyoike TW, Liburd OE. (2009). IPM programs in vegetable crops in Australia and USA: Current status and emerging trends. In: Peshin R, Dhawan AK (eds.). Integrated pest management: Innovation-Development Process, Volume 1. Springer Press. pp. 575-597.

SMARTtrain - Chemical use training and accreditation www.smarttrain.com.au

Storrie, A (2004) Reducing herbicide spray drift (Agnote DPI-477), NSW DPI, NSW. http://www.dpi.nsw.gov. au/\_\_data/assets/pdf\_file/0006/156993/herbicide-drift.pdf

Taverner, P; Wood, G; Jevremov, D; Doyle, B. (2006) Revegetation by design handbook http://www.sardi.sa.gov.au/\_\_data/assets/pdf\_file/0008/44945/reveg\_by\_design\_guidebook.pdf

Weatherzone www.weatherzone.com.au

Woods, N.; Cowles, G.; Crome, J; Lambourne, R.; Simpson, P.; Webster, R, (2005) Agricultural Chemical User's Manual, Department of Agriculture, Fisheries and Forestry, QLD. http://www.daff.qld.gov.au/plants/agvetchemicals-and-residues/chemical-use/agricultural-chemical-users-manual

Workplace Standards Tasmania Guidance note - How to HAZCHEM placard premises storing dangerous goods http://workplacestandards.tas.gov.au/resources/law/work\_health\_and\_safety\_laws



Horticulture Australia

# **Chemical management - references and further resources**





# Chapter 4 Nutrient management



Objective – the productive capacity of the soil is maintained without detriment to the environment

Most Australian soils are naturally low in nutrients. By managing nutrient application and soil fertility, production targets can be achieved without environmental harm.

It is important to apply fertilisers correctly because, if wrongly used, they may contribute to off-site degradation of groundwater and waterways, increase soil acidity, salinity and sodicity problems and contaminate soil.

Fertilisers may be lost from the production area through:

- Inaccurate application;
- Leaching past the root zone and into groundwater;
- Moving as dissolved nutrients in surface water leaving farm paddocks;
- Attaching to soil sediments and within organic particles in surface water leaving farm paddocks;
- Attaching to wind-eroded soil particles; and
- Volatilisation into the atmosphere.

Not only are these nutrients lost for crop production, such losses will potentially have downstream or off-farm impacts on the environment. The nutrients most at risk of causing off-farm impacts are nitrogen and phosphorus.

### Nitrogen

Nitrogen is an important macronutrient in soil and is essential to plant growth and high yield. It is present in soils, either as part of organic matter, which is unavailable for plant uptake, or in mineralised form (nitrate or ammonium ions), which is available to plants. In most soils, more than 95% of the nitrogen is present in organic form.

Further references and resources can be located at the end of this chapter.

# 4a Nutrient requirements

Nitrogen is a highly soluble element that is easily leached from the soil profile, dissolved in run-off water or volatilised into the atmosphere.

### Phosphorus

Like soil nitrogen, phosphorous is also a macronutrient and critically important to crop yield. Phosphorous can be added to the soil through the application of phosphate fertiliser to maintain the productive capacity of the soil. However, only a small proportion of total phosphorous is accessible to plants (1-4%) and its availability is highly dependent on soil pH.

Phosphorus binds strongly to soil particles and so can be lost by soil erosion through water and wind. Environmentally significant quantities of phosphorous can also be dissolved in run-off water when soil phosphorus levels are high.

### 4a Nutrient requirements



Objective – to effectively manage nutrient inputs to meet crop requirements and soil characteristics

To effectively manage nutrient inputs it is important to determine the amount and type of nutrients to apply for each cropping situation rather than using recipe-type application rates. This can be done through soil testing and nutrient budgeting.



resources can be located at the end of this chapter.



Horticulture Australia

Read the Suggested Practices for that chapter.

Further references and resources can be located at the end of this chapter.





### Review checklist

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

### Selecting nutrient types and amounts

Objective methods such as soil testing, plant tissue testing and sap testing, combined with yield data and visual assessments of crop or tree health, provide the basis for good fertiliser management. Fertilisers should be applied efficiently, taking seasonal conditions into account. This means applying just enough nutrients for good crop growth without providing excess nutrients that may be lost off farm into groundwater and surface waterways.

An understanding of the role of different nutrients in plant growth, the levels needed for good growth by particular crops and the cycles of key nutrients such as nitrogen is important. It is a good idea to either do formal training to learn to estimate the quantity and type of fertiliser to use, or to use the services of a soil consultant or agronomist to decide on the nutritional needs of crops. Both over- and under-application of fertiliser can create problems.

The Australian fertiliser industry has established the Fertcare® program to assist in the sustainable use of fertilisers. Fertcare® provides training and accreditation for agronomy, sales and logistics staff. Training options also exist for growers, depending on their particular needs.

Nutrient deficiencies can lead to low yields, poor quality crops and financial loss. They can also lead to indirect environmental damage because other nutrients are used inefficiently and will be vulnerable to loss off-site. Poor crop growth also leads to reduced organic matter returns to the soil, which can reduce soil health and cause erosion and still poorer crop growth.

The relatively low cost of fertiliser compared to other inputs means growers have tended to over-compensate with fertiliser application rather than face the risk of production losses.

Over-supply of fertiliser or incorrect use of blended fertilisers (e.g. NPK mixes) also brings problems. It is not only an unnecessary cost, but can cause reduced yields through toxic levels of nutrients or by inducing deficiencies through nutrient imbalances. Over-supply can also lead to long-term, off-farm impacts like degradation of sensitive environmental areas, algal blooms in waterways and groundwater contamination, and on-farm impacts such as increased soil acidity, soil sodicity and soil salinity.

Inorganic fertilisers are only one method for supplying nutrients. Cover crops, fallow crop residues, composts and animal manures not only add organic matter to the soil but also release significant amounts of nutrients as they break down. The gradual release of nutrients from organic sources can help to reduce the loss. For example researchers have found that banana plant crop residues can contribute 60 kg N/ha to the following ratoon crop in Far North Queensland conditions. Crop residues from a cauliflower crop can contain about 170 kg of nitrogen (N), 27 kg of phosphorus (P) and 180 kg of potassium (K) per hectare. Animal manures can add significant amounts of phosphorus to the soil. Manures should be tested for nutrient content, chlorine levels and presence of contaminants such as heavy metals before application.

Fertilisers should be selected that have low levels of contaminants such as heavy metals (cadmium, lead or mercury). High levels in fertilisers may lead to accumulation in the soil or uptake by crops in excess of maximum levels for human safety (for information regarding maximum levels, see the Food Standards Code www.foodstandards.gov.au).

### Soil and sap testing

Soil testing is a useful way to objectively measure the nutrient status of your soil. It is a particularly valuable nutrient management tool before planting a crop or orchard. Ongoing soil testing (say every one to three years) also provides valuable insights into longer-term trends in soil properties that may alert managers to developing sustainability problems. Soil organic carbon decline or the build-up of high available phosphorus levels are examples of this.

Soil testing, plant tissue testing, sap testing and visual crop inspection can all be used post-planting to monitor nutrient availability and determine an appropriate post-planting fertiliser program.

Soil tests measure soil properties that influence nutrient availability to the plant. These include pH, electrical conductivity (a measure of salt content), organic carbon, individual macro- and micronutrients and other elements. It is a good idea to use a laboratory that is accredited to carry out the required tests. Look for NATA (National Association of Testing Authorities) accreditation, a laboratory that uses NATA methods or one that participates in Australian Soil and Plant Analysis Council (ASPAC) proficiency trials. See http://www.nata.asn.au or http://www.aspac-australasia.com.

For soil test results to be meaningful, the sample must be carefully collected. When collecting a sample make sure it represents the area being tested, by taking into account the total area of the block, any changes in soil type within the block and the depth of sampling. The samples should be representative of the root zone. Taking a 10-15 cm sample from soil that has been worked to rooting depth will achieve this. In orchards, where soil is not worked, a shallow sample may not represent actual nutrient availability. Subsoil samples may be of value to determine nutrient availability for deep-rooting crops and identify possible causes for nutrient imbalances.

Sufficient samples should be collected to be representative of the site and should not include soil from any unusual areas. These samples should be bulked, mixed well and then a sub-sample of this bulk soil sample sent in for testing. The testing laboratory or its agent should provide instructions on taking the sample. Soil consultants or agronomists can also collect samples for you and there are accreditation programs for these services.

Nitrogen levels undergo dynamic changes during the season, being influenced by factors such as soil organic matter content, soil temperature and moisture. It is therefore important that soil samples for nitrogen budgets be taken separately from conventional soil samples and sampling should occur as close to the proposed nitrogen application date as possible to give an accurate picture of current nitrogen availability.

Soil test results and optimum soil nutrient levels should be discussed with an appropriately qualified person, such as an agronomist or soil consultant. Based on this interpretation and consideration of soil type, cropping history, specific crop needs and agronomy, a written fertiliser recommendation should be provided. This may include recommendations for adding lime, dolomite or gypsum.



Horticulture Australia

Further references and

resources can be located at

## 4a Nutrient requirements

#### Further references and resources can be located at the end of this chapter.



Soil testing and analysis needs to be completed early enough to allow nutrients and soil ameliorants to be applied in a timely manner.

Sap testing can also be used to develop nutrient uptake graphs, so fertiliser applications can be timed to the appropriate growth stage of the crop. Samples for sap testing need to be collected carefully and tissues analysed by a suitably proficient laboratory.

## **Nutrient budgeting**

Nutrient budgeting can help growers better understand the whole nutrient cycling and transformation system. This can lead to the design of more sustainable, integrated nutrition strategies.

A nutrient budget is like an accounting system for nutrients. It involves:

- Estimating the amount of nutrients available from the soil (soil test results);
- Obtaining uptake and removal figures for the target crop and the previous crop (to account for nutrients in crop residues, for example, consideration should also be given to the contribution of legumes to nitrogen availability). Figures should be in kg/ tonne of crop grown (for uptake) and harvested (for removal);
- Determining the target yield to calculate actual uptake and removal figures;
- Calculating the amount of nutrients, especially nitrogen, that will be applied with irrigation water (50 ppm nitrate in irrigation water will add about 1 kg n/ha with every 10 mm of irrigation water applied);
- Calculating the amount of nutrients already applied to a paddock;
- Estimating the amount of nutrients that will be removed through harvested product; determining possible nutrient losses through leaching (see Topic 2b – Water quality), volatilisation or soil erosion (see Topic 1a – Soil erosion caused by water, and Topic 1b – Soil erosion caused by wind). Deep soil nitrate testing can be an important tool in assessing leaching; and
- Replacing nutrients lost to the system through appropriate fertiliser applications.

Nitrogen, phosphorus and other major nutrients are the main elements considered in nutrient budgeting. Along with soil, leaf and sap testing and visual assessments, nutrient budgeting is another tool for fine-tuning the nutrient management program.

A nutrient budget should be prepared for a 3–5 year rotation. Break or cover crops should be considered as 'catch crops'. Nutrients that have not been used by the previous crop will be taken up by the break crop, thus avoiding leaching past the root zone.

Reviewing local research and advice from agronomists can also assist in determining fertiliser requirements, particularly in situations where key information such as crop nutrient removal rates are not known.

It is also important to determine if any nutrients are required in 'luxury' amounts (that is over and above the nutrient removal figures). For example, potassium may be applied at higher rates because of its role in preventing bruising.



Monitoring and recording

Soil test results, sap test results and corresponding fertiliser recommendations support responsible use of fertilisers. It is also a good idea to have documentation to support the credentials of the person providing the fertiliser recommendation.

Testing run-off and drainage water for nutrient content gives a good indication of any losses being experienced. Water can be monitored for nitrates using systems such as the CSIRO FullSTOP™.



### **References and further resources**

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

Further references and resources can be located at the end of this chapter.



### 4b Nutrient application

Ø

Objective – to ensure nutrient application methods and timing maximise benefits to the crop and minimise potential negative environmental impacts

Applying fertilisers correctly is as important as using the correct amount and type of fertiliser. Effective fertiliser application involves:

- The right rate and frequency, the right time; and
- The right placement.

To achieve this, application equipment must be set up correctly.

#### A note on organic farming:

Organic farming avoids using synthetic chemicals, artificial fertilisers or genetically modified (GM) organisms. Organic farming practices are described by an organic standard, Australia Certified Organic Standard (ACOS 2013), which describes parameters for compliance, noting that these requirements do not override legislative requirements but differentiate organic agriculture practices from traditional farming practices. The standard considers all other primary production or food preparation that does not conform as 'conventional' production. These guidelines would therefore be seen to cover 'conventional' horticulture production.

For more information on organic farming methods visit the Australian Organics website http://austorganic.com.



Horticulture Australia

Further references and

the end of this chapter.

resources can be located at

# 4b Nutrient application

## HIGH RISK

Review the risk of nutrient deficiencies & environmental pollution from poor nutrient management.

## HIGH RISK

Review fertiliser application practices. Review the Suggested Practices in this topic.

LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

**HIGH RISK** – You need to take some action. Read the Suggested Practices for that chapter. Further references and resources can be located at the end of this chapter.





### **Review checklist**

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

When applying fertilisers, some general rules should be followed:

- Avoid applying fertilisers to saturated soil or when heavy rain is forecast;
- Avoid applying fertiliser during extended drought;
- Avoid fertiliser application to steeply sloping ground. Use contour drains to minimise run-off;
- Monitor soil moisture to avoid leaching of nutrients past the plant root zone;
- Maintain vegetation cover during typically rainy periods to minimise run-off and leaching.

#### Type, timing and rates of application

Fertilisers need to be applied when they will do the most good for the crop. As a general rule, applying small amounts regularly is less likely to cause off-site losses from leaching and run-off. Schedule fertiliser applications according to seasonal conditions, cropping cycle and periods of greatest use by the crop. For instance, young vegetable crops require small amounts of nutrients until they begin to grow rapidly.

Nitrogen, in particular, should not be applied in large amounts while crops are young and nitrogen demand is low. Large applications of pre-plant fertilisers are vulnerable to loss in the slow growth period. Select the most suitable fertiliser type depending on the speed of availability of nutrients in relation to crop demand, acidity, alkalinity or salinity of fertiliser.

Pre-plant fertilisers should be incorporated. If there is a likelihood of heavy rain, minimise the amount of pre-plant fertiliser applied to reduce risks of fertiliser losses through leaching and soil wash from paddocks into nearby waterways. It may then be necessary to increase top-dressed fertiliser applications later in the season. However, light rain or irrigation is beneficial to incorporation and to reduce volatilisation of some fertilisers such as urea.

An auto steering system for application equipment will allow precise applications without overlaps and may reduce fertiliser costs.

#### Fertiliser placement

Accurate placement of fertilisers helps plants to access the nutrients required. Choose the right equipment and adjust it correctly to make sure fertiliser is applied to the area where it will do the most good but have the least impact on the environment.

Apply small amounts of fertiliser near the root zone of plants. Application methods suited to achieving this include:

- Fertigation using micro/trickle irrigation systems, banding to the side or below seeds or transplants, banding or drilling fertiliser in beside plant rows;
- Broadcasting fertiliser along the drip line of tree crops, and broadcasting fertiliser before the crop canopy closes.

Foliar application through spray equipment is a useful method for applying targeted amounts of micronutrients.

Be careful not to apply fertiliser to non-crop areas or adjacent to waterways. Take steps to prevent contamination of water sources from pump backflow during fertigation.

#### Care and calibration of equipment

Brand new spreaders can have poor spread patterns, and with use and 'wear and tear' even a well set up spreader can become inaccurate. Therefore fertiliser application equipment needs to be carefully calibrated and maintained to make sure it is capable of spreading fertiliser evenly at the correct rate.

Refer to the manufacturer's specifications when carrying out calibration.

Accu-Spread® is a testing and accreditation program that ensures your spreader can apply fertiliser evenly across the paddock, maximising productive response and minimising environmental risk.

#### Storage of fertilisers

All fertilisers including animal manures should be stored in such a way that prevents nutrients leaching into surface waterways and groundwater. Inorganic fertilisers should be stored in a covered area away from waterways. Manure heaps should also be covered to reduce leaching through rain.

Storage areas should be:

- Protected from direct sunlight and rain;
- Well ventilated with fresh air to keep fertilisers dry; • Designed to minimise the chance of pest infestation, mould growth and
- damage; and
- Designed to confine any spillage and allow easy clean up.

Store fertiliser in a way that lowers the risk of seepage into groundwater. With the exception of fertilisers applied with pesticides, fertilisers should be stored separately from other agricultural chemicals.

All liquid fertiliser storage should be bunded to eliminate the chance of run-off into waterways. In the absence of any national or state legislation, the bund should be 125% of the largest container, plus 25% of total volume stored.

In addition to regulations regarding storage of dangerous goods, there are new security regulations in place regarding ammonium nitrate that cover how these products are supplied, handled and stored. Advice from the appropriate local authority should be sought.

### Disposal of packaging

Used fertiliser packaging should be stored in a manner that prevents contamination and environmental harm and meets local government waste disposal regulations.

Further references and resources can be located at the end of this chapter.



4.10 Chapter 4 Nutrient management

© 2014 – Horticulture Australia Limited

# 4b Nutrient application

#### Further references and resources can be located at the end of this chapter





### Monitoring and recording

#### Fertiliser application records

It is recommended that an accurate record be maintained of all fertiliser applications, including foliar applications and fertigations. This applies both to organic (e.g. sheep, cattle, chicken manure) and inorganic fertilisers (e.g. superphosphate). Fertiliser application records are essential for nutrient budgeting.

Suggested headings in fertiliser records are:

- The location of the treated areas (block or paddock identification);
- Application dates;
- The type of fertiliser used including the trade name, type of fertiliser or concentration of nutrients;
- Amount of fertiliser applied per hectare (weight or volume);
- Method of application and machinery used e.g. fertigation, spreader; and
- Name of the operator applying the fertiliser.

Soil test results for the paddock and sap and leaf tests for the crop support these fertiliser records.

#### Machinery calibration and maintenance records

It is suggested that maintenance and calibration records for fertiliser application equipment be kept. This should include:

- Equipment/machinery name;
- Date on which calibration/maintenance was performed;
- Work undertaken; and
- Signature or initials of the person who performed the calibration/maintenance, or an appropriate invoice.



#### **References and further resources**

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter



Horticulture Australia

## Nutrient management - references and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www.horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

Accu-Spread - independent testing and accreditation of fertiliser spreading equipment for accuracy and evenness of spreading. http://www.afsa.net.au/index.php?action=content&page=12

Armour, J. & Daniells, J. 2002. Banana nutrition in North Queensland. HAL project report FR95013. Department of Natural Resources & Mines, QLD.

Australian Certified Organic (2013) Australian Certified Organic Standard, Australian Organic http://austorganic.com/australian-certified-organic-standard1/

Australian Soil and Plant Analysis Council http://www.aspac-australasia.com An independent international organisation consisting of individuals, laboratories, research and commercial organisations involved in soil and plant tissue analysis.

CSIRO FullStop www.fullstop.com.au The FullStop Wetting Front Detector helps you to "see" what is happening down in the root zone when you irrigate the soil.

FertCare http://www.fertilizer.org.au/files/pdf/fertcare/Fertcare%20Booklet.pdf FertCare is a national product stewardship initiative of the Fertilizer Industry Federation of Australia (FIFA) and the Australian Fertiliser Services Association (AFSA), supported by Australian Government. It provides training to assist industry participants understand the food safety and environmental risks involved in handling, transporting, storing and spreading fertilizers.

Fertilizer Industry Federation of Australia, (2001), Cracking the nutrient code: Guidelines for developing a Nutrient Management Code of Practice for your industry, region or farm, FIFA, Canberra, ACT. http://www.fertilizer.org.au/files/pdf/publications/cracking%20the%20nutrient%20code%20optimized.pdf

Food Standards Australia New Zealand (FSANZ) www.foodstandards.gov.au FSANZ is responsible for the development and administration of the Australia New Zealand Food Standards Code, which lists requirements for foods such as additives, food safety, labelling and GM foods.

Incitec Pivot Fertilizers http://www.incitecpivotfertilisers.com.au/Soil%20Plant%20Tests/FieldFacts This website provides link to fact sheets, such as Horticulture soil sampling procedure.

National Association of Testing Authorities, Australia http://www.nata.asn.au NATA provides assessment, accreditation and training services to laboratories and technical facilities.

National Land & Water Resources Audit (2008) Signposts for Australian Agriculture — The Australian Horticulture Industry. NLWRA, Canberra. http://lwa.gov.au/files/products/national-land-and-water-resourcesaudit/pn21405/pn21405.pdf

NSW Department of Primary Industries - Crop nutrient replacement in tropical horticulture - Links to Excel spreadsheets are provided for the following fruit crops so that you can calculate crop nutrient replacement – Avocados, Custard apples, Low-chill peaches, Mangoes, and Passionfruit. http://www.dpi.nsw.gov.au/agriculture/horticulture/tropical/nutrients/replacement

Phillips, D. (2010) Developing guidelines for environmentally sustainable use of mineral fertilisers (HAL Project Number: VG07036). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Potassium in Agriculture http://www.pda.org.uk/technical-potash-notes.php This website provides technical information on soil fertility, plant nutrition and fertiliser use with particular emphasis on potash (based in UK).

# Nutrient management - references and further



Chapter 4 Nutrient management 4.13



# Chapter 5 Biodiversity



Objective – native vegetation, wildlife and ecosystems are appropriately maintained and managed and, where possible and practical, contribute to regional biodiversity priorities

Biodiversity is the variety of all life forms: the different plants, animals, fish, birds, insects and micro-organisms, their genes and the ecosystems of which they are a part.

Biodiversity is increasingly being recognised for its contribution to farm sustainability and productivity.

Biodiversity does not just apply to native organisms, but the focus of this chapter is largely on managing native biodiversity. This does not imply that biodiversity is not important within the production area.

Native biodiversity refers to the biodiversity found in a particular locality. It is restricted to the local ecosystems and their components, be they native plants, animals or micro-organisms.

Native biodiversity provides many benefits that are essential to sustaining and fulfilling human life and maintaining productive agriculture. These benefits are called 'ecosystem services' and include:

- Fungi, worms and bacteria transforming sunlight, carbon and nitrogen into fertile soil;
- Pollination from insects, including native bees, such as in the photograph above;
- Regulation of climate;
- Provision of shade and shelter from native vegetation; and
- Waste absorption and breakdown.

Further references and resources can be located at the end of this chapter.

Horticulture and biodiversity have complex interactions. If native biodiversity and ecosystems deteriorate, then the quality of the soil, water, air and ultimately productivity declines. The initial clearing of land may have negative impacts but horticultural properties, once established, become part of the local landscape. Crops may provide shelter and food for wildlife. Some wildlife may help control pests in crops, while other species may become pests themselves.

Ecosystem resilience is the capacity of an ecosystem to respond to changes and disturbances, yet retain its basic functions and structures. The resilience of ecosystems in Australia is currently being reduced by a number of threats, including climate change.

There is a need to reorientate management objectives from preserving all species in their natural habitat and current locations to ensuring space and opportunities are available for ecosystems to adapt and reorganise. This will increase the chances that they can maintain the provision of ecosystem services through a diversity of well functioning ecosystems.

To manage biodiversity on your property you need to know what native vegetation exists and, if it is of special importance, consider the impacts of farming operations on the environment and develop practical ways to manage any native vegetation as well as problem plants and animals. Management practices such as using native plants in buffer plantations or as habitat for beneficial (pest controlling) birds and invertebrates can add to efforts to conserve remnant on-farm vegetation. There may also be areas where revegetation can occur and be coordinated to contribute to regional initiatives. The careful application of pesticides is another way to protect local biodiversity from harm.



-IAI

# **5** Biodiversity

Further references and resources can be located at the end of this chapter





### **Review checklist**

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

By adopting farming practices that are compatible with biodiversity protection, land managers can work towards sustainable agriculture and integrating agricultural production with biodiversity protection.

In some instances, rather than identify and assess all the native biodiversity on your property (i.e. all native animals, plants and insects etc.) it is more practical to use native vegetation as an indicator to identify if native biodiversity is present.

#### Identify native vegetation on your property

An initial assessment should try to identify any local native vegetation still left on the farm (exclude plantations and vegetation established for commercial purposes). This could include naturally occurring trees, shrubs, herbs and grasses, areas with understorey, or corridors connecting larger areas of natural habitat, native pasture, large old trees, wetlands, rocky outcrops, and areas of fallen timber. Dead trees should be included as important components as they provide habitat for native animals and insects. Create an inventory/file of this information.

Contact your State conservation, natural resources or sustainability department and ask them for any maps and/or lists of the native vegetation that is likely to be present and to provide advice regarding management of native plants and animals that may be on your property. Other people to help with identification include:

- Field naturalists, Greening Australia;
- Local catchment management authorities, Bushcare;
- Landcare groups;
- Regional Natural Resource Management (NRM) groups; and
- Local/State herbarium.

Universities and other research bodies are sometimes looking for farms on which to conduct studies of native plants and animals. This can be a good opportunity to learn more about the wildlife on your farm.

Attending field days in your catchment is also a good way to learn about local plant and animal species.

You may like to undertake a more complete and detailed assessment of all the species (native plants and animals) located, or likely to be present, on your property. You can ask a State biodiversity or environmental officer for advice. Biologists or ecologists are qualified to carry out this more detailed survey.

A full native biodiversity assessment could include:

- A map of the vegetation likely to be present prior to development;
- Maps of the location and type of native plants and animals species currently known to be on the farm;
- A list of local plants and animals in danger of dying out (i.e. threatened species);
- A list of current farm activities that is helping/harming native plants and animals;
- Future actions to improve biodiversity as well as maintaining farm productivity; and
- The expected improvement if these actions are followed.

The Environmental Management in Agriculture: Native Biodiversity Resource Kit provides guidance on these steps. See http://live.greeningaustralia.org.au/nativevegetation/pages/ pdf/Authors%20S/12\_Straker\_Platt.pdf.

#### Consider surrounding properties

No farm works in isolation of its neighbours. Just because you haven't found any native vegetation on your property doesn't necessarily mean there is no native biodiversity, or that you can ignore the impacts your farm operations may have on surrounding properties. Look for native birds and listen for frogs - chances are both are present, indicating that suitable habitat is located in surrounding areas.

Any basic knowledge of the surrounding area, aerial photos, satellite images or maps of vegetation will give you an idea of what native vegetation is located near your property.

#### Assess special importance

The Government has developed lists of plants that are considered important because of their rarity, they are particularly subject to threats, or may support other significant features (e.g. as a drought refuge for native animals).

Some native vegetation stands are more important than others. This includes:

- Threatened species protected under legislation;
- Remnant vegetation (land that has never been cleared, or if previously cleared, regrowth is now mature);
- Larger areas of vegetation; and/or
- Areas serving as wildlife corridors.

More information can be found on the Australian Government Department of Environment website: http://www.environment.gov.au/topics/about-us/legislation/ environment-protection-and-biodiversity-conservation-act-1999.

Contact your local government, State conservation department or the regional catchment management authority/Natural Resource Management committee for information about any important or significant vegetation that may be in your region. Financial assistance may be available to assist with management of important vegetation. Check out conservation incentives on www.environment.gov.au or contact your local catchment authority.

The following categories may have been used to describe threatened plant species:

- Extinct no reasonable doubt that the last member of the species has died;
- Critically endangered extremely high risk of extinction in the immediate future;
- Endangered very high risk of extinction in the near future; and
- Vulnerable high risk of extinction in the medium term.

Each of these categories may carry specific legislative responsibilities you are required to undertake.



Further references and

the end of this chapter.

resources can be located at

5.4 Chapter 5 Biodiversity

© 2014 – Horticulture Australia Limited

# **5** Biodiversity

#### Further references and resources can be located at the end of this chapte



#### Assess off-farm impacts and threats

Site development or redevelopment works need to be assessed for their potential impacts on the existing environment.

Management actions for biodiversity need to address the cause of decline or what threatens the continued existence of the native biodiversity. In many cases these threats have their origin in inappropriate land and natural resource management activities.

To assist you to assess your impacts on native biodiversity, potential management actions addressing threats should be organised according to farm activities. It is expected that each farm will be unique, therefore these actions should be considered as prompts that a land manager should use to broaden the search for solutions.

In the Environmental impacts section of these guidelines, you will note that most activities represent some form of impact both on and off-farm. However, the five key horticultural activities you may need to draw up specific actions for are:

- Spray drift;
- Fertiliser/water use and leakage;
- Topsoil loss and erosion;
- Control of pests (including native animal species); and
- Disposal of waste.

An integrated property management plan such as a Whole Farm Plan (WFP), Property Management Plan (PMP) or similar, can help you to work through the range of potentially conflicting needs and allocate resources according to priority.

#### Living with native birds and animals

There are many benefits to having native animals on or near agricultural land. For example, many native birds eat insect pests, pollinate plants and disperse native seed. However, occasionally problems may arise when native birds and animals eat or damage crops. Where growers are faced with 'problem' native animals, specialised advice must always be sought from your State conservation department to avoid exposing yourself to the risk of prosecution if any illegal response is adopted.

Avoid looking at symptoms of the problem, such as 'there are too many animals' – the question is why? Options for responding to problem animal management will include how to mitigate the problem and live with native animals. As standard practice, always consider non-lethal management options. These may include:

- Netting;
- Fencing;
- Sound or light based systems (sirens, gas cannons);
- Encouraging predators (e.g. hawks, although this may end up being lethal!); and/or
- Providing alternative habitat.

# See http://www.wildlifefriendlyfencing.com/WFF/Friendly\_Fencing.html and http://www.wildlifefriendlyfencing.com/WFF/Netting.html for examples.

If you have sought advice and trialed applicable non-lethal management options without success, as a last resort consider lethal management options (such as shooting). It is important to check whether wildlife is protected and be aware of any licensing requirements before undertaking lethal management options. These must only be implemented in a humane manner.

Consider registering your farm under the 'Land for Wildlife' program if it operates in your area. This program is delivered by different organisations in each state. See Greening Australia or Landcare Australia for relevant information.

Keep pets away from wildlife. Unrestrained cats and dogs can wreak havoc on local wildlife populations. If pets are kept in at night, this will help safeguard them as well as the wildlife. Ideally, cats should be confined to indoor areas at all times.

Look after wetlands and watercourses. The natural cycles of flood and drought are important in maintaining ecosystems and habitats. In wetland areas, floods rejuvenate the soil and create temporary habitat and breeding sites for waterbirds, tadpoles and fish. Waterbirds eat many crop damaging insects, and the common backswimmer and the nymphs of dragonflies and damselflies eat mosquito larvae. Adult dragonflies and damselflies also prey on mosquitoes and crop damaging insects such as aphids.

#### Well maintained waterways are also attractive and an asset to any property

Make a farm dam more wildlife friendly. A dam with a large surface area, variety of depths and gently sloping bank also provides different habitats for many vertebrate and insect species. Logs, rocks and boulders in and around the dam will provide homes many animals.

Leave river snags, submerged logs, branches and litter in place. If it looks messy, chances are it is great habitat! Snags, logs and litter provide habitat for fish, frogs and invertebrates. Half submerged logs also provide perching spots for birds and turtles. Hollow logs provide homes for many species. Logs from other areas can be used to resnag watercourses and improve habitat values.

#### Controlling feral animals and invasive species

Invasive species can be both native and exotic. They can have an impact on the environment by competing with native plants and animals and potentially can cause soil degradation and water quality issues. If pests, such as feral pigs, deer and weeds, are a problem, contact your local natural resource department or council, to get advice or assistance to control them.

Exotic plants may require specific management. Problem plants can be escapees or volunteers from commercial operations, such as escaped olive trees. In the right place these plants are not a problem, but once they start encroaching on native vegetation specific action is required. The landowner must take all reasonable measures to prevent the land being infested with a declared weed and take all reasonable measures to prevent a declared weed or potential weed spreading to other land.

Get rid of weeds to reduce potential for pest harbour and give native vegetation a better chance of survival (e.g. Bridal Creeper Asparagus asparagoides – a Weed of National Significance not only causes losses by shading citrus and avocado trees and interfering with fruit picking, it is also considered the most important weed threat to biodiversity).

If you are planning to use herbicides for weed control rather than physical removal, take special care not to damage the area you are trying to improve or to contaminate waterways. Removing dense areas of weeds should coincide with revegetation or regeneration activities, otherwise further weed infestation may result. For more specific information on threats and control methods for individual weeds refer to http://www.weeds.org.au.

#### Environmental weeds and nursery propagation

Weeds have a significant impact on the environment because they compete with native plants and reduce habitat and food sources for wildlife. Weeds Australia publish 'Weeds of National Significance' (WoNS), which currently lists 32 weeds that, by law, must be controlled if they are present on your land. See www.weeds.org.au/WoNS.

'Noxious' weeds must also be controlled under legislation at a state or territory level. Of the 2700 species of introduced plants now established, 429 have been declared noxious or are under some form of legislative control in Australia. Weeds Australia also publish a searchable list of noxious weeds at www.weeds.org.au/noxious.htm.

5.6 Chapter 5 Biodiversity

Horticulture Australia



# **5 Biodiversity**

#### Further references and resources can be located at the end of this chapter.



Identify and manage weeds on your property according to the relevant legislation. Weeds around riparian areas and watercourses are also the landowner's responsibility. General information relating to identification and control measures for weeds can be found on the Weeds Australia website www.weeds.gov.au.

Soils amendments, mulch and fill may harbor weeds, so only source these products from a trusted supplier and manage these inputs as well as possible. Inspect the places where these products are applied regularly to identify any potential weed incursions.

#### Role of the nursery and garden industry in reducing the spread of invasive garden plants

With the increasing focus on the impact that invasive garden species have on the environment and agriculture, the nursery and garden industry and its customers are frequently targeted for education and awareness strategies. The difficulty is that 'customers' include not only gardeners, but also a diverse range of people with differing needs, knowledge and skills.

The Nursery & Garden Industry NSW & ACT Limited's 'Grow Me Instead' programme has been very successful in highlighting the importance of industry and local government partnerships in identifying and targeting key invasive garden species on a regional scale. Through daily contact with customers and stakeholders who are making decisions and seeking advice about their gardens, the industry is in a unique position to educate the community about responsible plant choice and to promote appropriate care and maintenance to reduce the spread of invasive garden species and to better manage plants already in gardens.

#### Practical management of native vegetation

Once you have found out which native plants are on your property (including their significance) you will have some idea about how to prioritise your actions to protect them. These actions may include:

- Fencing off areas to exclude vehicles, people and stock. Select fence types that enable native animals to have access to natural drinking water sources and to move between habitats;
- Animals to have access to natural drinking water sources and to move between habitats;
- Leaving dead trees standing, logs, branches, twigs and rocks on the ground as homes for birds, insects and other animals;
- Not cleaning up places with native vegetation. By not tidying up understorey grasses, shrubs and fallen trees, birds and beneficial native animals will have places to hide from introduced predators or competitors or as a food source; and
- Protecting native vegetation from fertiliser. Avoid the use of fertiliser on native vegetation, ensure native vegetation is protected from accidental application such as drift. Fertilisers raise nutrient levels, encouraging invasion by exotic species and also reducing the amount of native ground cover species, including valuable perennial native grasses.

You can also ask a State biodiversity or environmental officer for advice or assistance on priorities for management.

Depending on the jurisdiction, it may be possible to enter into a voluntary conservation agreement or similar agreement with the relevant State agency to formalise protection of wildlife and significant habitat on part of your property. 'Land for Wildlife' is one such scheme and conservation incentives also exist (http://www.environment.nsw.gov.au/cpp/ landforwildlife.htm). Sometimes these agreements may contribute to providing compliance against international environmental standards.

Many pest animals not only prey on native animals but they also compete for the same space, food and shelter. Control animals such as rabbits, foxes and feral animals. Check local regulations and laws about the control of pest animals before you begin any control program.

Integrated Pest Management (IPM) is an excellent management practice to maximise biodiversity on-farm. IPM strategies are designed to reduce reliance on pesticides while still managing pests. Reduced reliance on pesticides can provide an environment where a greater diversity of flora and fauna exists. This in turn can assist in building the numbers of pest predators to improve the control of pests.

Whole farm planning is a structured approach to making informed decisions for the use and management of all aspects of your property. Your State biodiversity or environmental officer, catchment management authority or natural resource management group can provide advice.

#### Fire management

Management of vegetation areas needs to also consider fire control and the role of fire in maintaining the diversity of plants that make up the bush. Much of Australia's flora has evolved in an environment where fires regularly occurred, and many plants require fire to assist regeneration.

Considerable care is required to manage fires and local authorities should be consulted and alerted before burning. Neighbours may also be affected by smoke and should also be consulted (see Chapter 7 – Air management).

The following points should be reviewed:

- Choose a fire regime to suit the desired outcome. If you are burning to reduce fuel loads then fires will need to be more frequent than if you were burning to promote tree regeneration;
- If you have threatened species, choose a fire regime that suits their needs;
- Time burning to suit the plants' lifecycles. These will vary depending on where you are in Australia, but generally autumn burns are best; and
- Fireproof buildings and ensure sufficient fire breaks around production areas, boundaries and other areas that must not be burnt.

If the property is affected by bushfire or a prescribed burn, the bush will recover. This can be a valuable opportunity for controlling weeds and kick-starting natural regeneration.

#### Consider options for increasing on-farm native vegetation

Plantings can greatly enhance a property's productivity and wildlife habitat value. Depending on their design and placement, they provide a range of services including:

- Windbreaks, or shelterbelts protecting crops, pastures, livestock and farm buildings;
- Additional habitat area potentially boosting numbers of native animals that pollinate plants, and control insect pests;
- Corridors to help increase connectivity between patches of habitat;
- Help in reducing water tables and control salinity;
- Erosion control:
- Supply timber and fodder; and
- · Improvement to water quality in wetlands, dams and watercourses.

5.8 Chapter 5 Biodiversity Horticulture Australia



Further references and

the end of this chapter.

resources can be located at

# **5** Biodiversity

#### Further references and resources can be located at the end of this chapte



Think about planting windbreaks and shelterbelts using local native species. Shelterbelts and windbreaks may be best placed on the property boundaries and developed with consideration of establishing interconnecting wildlife corridors.

The resulting wind protection can contribute to overall farm productivity, plantings may provide habitat for beneficial native insects (i.e. contribute to Integrated Pest Management) as well as providing habitat for native species. Ideally, plantings should be at least 20 m wide to provide greater benefits, however, if space is limited, narrower shelterbelts can still provide some protection to crops and provide some habitat. Individual trees can also provide suitable habitats.

Choose some areas where it would be possible to begin a restoration project. It is a good idea to choose areas not currently used for farming, such as steep slopes, stream/ dam sides, saline areas and wetlands. These areas are usually poorly utilised and don't make a major contribution to horticultural production.

Select a mix of native plants, including trees, shrubs and grasses, preferably native to your local area (known as provenance species). Plantings should copy nature and not be as regular as a crop.

Dams and waterways also significantly contribute to increasing biodiversity and providing habitat for native animals, birds, frogs, insects, fish, invertebrates and plants. Wetlands, bogs and marshy areas can be turned from unprofitable areas to rehabilitated areas of great ecological significance with fencing and revegetation.

Some local authorities and organisations will provide advice and support to landholders for revegetation activities. Flora, fauna and bush regeneration consultants are also available to assist in design of restoration projects.

#### Soil biodiversity

Soils contain many living organisms ranging from microscopic bacteria and fungi to burrowing animals. All play a part in maintaining natural soil processes, which are vital for maintaining the chemical and physical fertility of soil. Soil organisms rely on organic matter for food. By 'feeding' on organic matter, micro-organisms release organic nutrients in a mineral form available to plants for uptake. Increasing the organic matter content of your soil (such as by incorporating compost) can help boost the population size, diversity and activity of soil organisms. The organisms in the soil food web play a part in breaking larger pieces of litter into smaller fragments, mixing these throughout the soil profile, binding soil particles, providing channels for water access and maintaining the health of the plant. Bacteria and fungi are responsible for the rate of nutrient release to the soil and rhizosphere (the area around plant roots where the biology and chemistry of the soil are influenced by the root).

Biodiversity can be improved in production areas by strategies such as inter-cropping or alley cropping (growing two or more crops in the same area), rotations with a range of crops and cover crops, or by simply being more tolerant of weeds.

#### Work with others

Neighbouring land managers may have practical experience in addressing certain biodiversity issues that they can share. You can demonstrate your involvement in environmental issues and gain practical advice from fellow growers by joining local groups such as Landcare.

Contact these organisations to see if your on-farm activities can contribute to any local environmental projects and encourage your neighbours to work with you. It is a good idea to contact your regional NRM group to see how your property and activities may contribute to regional targets or strategies and to see if there is any financial assistance available to help you achieve your goals.

The Commonwealth has developed a Biodiversity Resource Guide as part of the National Environmental Management Systems (EMS) training kit and provides

#### state-by-state contact details. To find out more, visit http://www.daff.gov.au/naturalresources/soils/ems/biodiversity.

As animals travel between farms, it is a good idea to cooperate with your neighbours and connect your native vegetation with theirs.



### Monitoring and recording

Records demonstrating that you have achieved progress with your management of native vegetation, fauna and ecosystems serve not only to prove to yourself that your property is a healthy environment, but can also demonstrate your environmental stewardship 'credentials' to your neighbours, authorities, wholesalers, retail customers and consumers, which may go a long way when next you wish to expand your operations, change your focus, access new markets or even sell your property.

A vegetation assessment is a good way to understand current on-farm biodiversity and establish a benchmark for your property. When repeated over time, a reassessment can monitor and measure changes. Some guidance may be available from government environment departments and regional NRM groups. In the absence of better information, applying the general principle of trying to maintain the current condition of natural areas and taking some steps to improve them will benefit the environment and demonstrate your environmental stewardship.

Native vegetation and individual native plants have 'resilience' - the ability to regenerate after disturbance. It is the resilience of the plants, and the associated animals - from large mammals to microscopic soil bacteria - which governs the capacity of an ecosystem to recover when damaged.

Signs of strong resilience include:

- Older trees, which will shower the site with seed;
- At least some natives in the shrub layer native grasses and ground cover plants;
- Native regrowth and evidence of regeneration;
- Natural watercourses:
- Leaf litter, fallen timber, lichen and mosses; and
- Intact soil profiles and soil that has had minimal impacts from farming or urbanisation.

Signs of reduced resilience include:

- Heavy weed loads;
- Introduced pasture grasses;
- Land degradation and soil erosion;
- Salt scalding;
- Few living trees, and signs of dieback;
- High nutrient levels from fertiliser or farm effluent runoff; and
- Stormwater runoff from urban areas and roads.



Further references and

the end of this chapter.

resources can be located at

Horticulture Australia

# **5** Biodiversity

#### Further references and resources can be located at the end of this chapte



A farm map and photos can be used to demonstrate revegetation of locally native species and future plans. You may find it helpful to record sightings of rare or unusual animals along with your vegetation assessment maps and documentation.

Records of training, implementation and the management of Integrated Pest Management on farm can be used to demonstrate some of the production biodiversity enhancements in your business.

A simple measure of soil biodiversity is to count the number of earthworms in a spade full of soil. More sophisticated measures of soil biodiversity can be provided by some research laboratories set up to measure parameters such as microbial and fungal biomass, microbial activity and nematode community analysis.

Strategies for control of problem native animals can be documented and kept along with any licences required.



### **References and further resources**

For access to relevant references and further resources click here.

### **Biodiversity - references and further resources**

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www.horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

AgBiodiversity - portal for information on decisions about managing biodiversity and production. http://www.agbiodiversity.com

Australian Fauna Care - comprehensive list of wildlife care groups in Australia http://www.fauna.org.au

Australian Government - Caring for Our Country - Biodiversity and natural icons http://www.nrm.gov.au/about/caring/priorities/biodiversity.html

Australian Government Department of Agriculture - Environmental Management Systems - Biodiversity Resource Guide http://www.daff.gov.au/natural-resources/soils/ems/biodiversity

Australian Government Department of Environment - Conservation of biodiversity http://www.environment.gov.au/topics/biodiversity

Australian Museum on line Australia's Biodiversity http://australianmuseum.net.au/Biodiversity

Australian National Botanic Gardens http://www.anbg.gov.au/chabg/bg-dir/

Australian Weeds Strategy – A national strategy for weed management in Australia. Natural Resource Management Ministerial Council (2006), Australian Government Department of the Environment and Water Resources, Canberra ACT. http://www.environment.gov.au/biodiversity/invasive/weeds/publications/ strategies/pubs/weed-strategy.pdf

Bennett, A.F., Kimber, S.L. and Ryan, P.A. (2000) Revegetation and Wildlife – A guide to enhancing revegetated habitats for wildlife conservation in rural environments. Development Program Research Report 2/00 http://nrmonline.nrm.gov.au/downloads/mgl:787/PDF

Bureau of Rural Sciences (2010) Revegtation monitoring and reporting in Australia, Australian Government BRS. http://nrmonline.nrm.gov.au/downloads/mql:2718/PDF

Colloff MJ. (2011) The role of soil biodiversity in providing ecosystem services. Report prepared for the Australian Government Department of Sustainability, Environment, Water, Population and Communities on behalf of the State of the Environment 2011 Committee. DSEWPaC, Canberra, ACT. http://www.environment.gov.au/system/files/pages/ba3942af-f815-43d9-a0f3-dd26c19d83cd/files/soe2011supplementary-role-soil-biodiversity-providing-ecosystem-services.pdf

Department of Agriculture, Fisheries and Forestry - Landcare and ecosystem services information http://www.daff.gov.au/natural-resources/landcare http://www.daff.gov.au/natural-resources/ecosystemservices

Department of Environment and Primary Industries, Victoria – Biodiversity Atlas http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/victorian-biodiversity-atlas

Department of Primary Industries, Parks, Water and Environment - Land for Wildlife http://www.dpiw.tas.gov.au/inter.nsf/WebPages/DRAR-7T8VRQ?open

DEWHA (2009) Assessment of Australia's Terrestrial Biodiversity 2008, Australian Government Department of the Environment, Heritage and the Arts, Canberra, ACT. http://www.environment.gov.au/resource/ assessment-australias-terrestrial-biodiversity-2008

Dorrough, J., Stol, J. and McIntyre, S. (2008) Biodiversity in the Paddock: a Land Managers Guide. Future Farm Industries CRC. http://www.futurefarmonline.com.au/knowledge-base-1/biodiversity-in-the-paddock-a-landmanagers-guide?A=SearchResult&SearchID=6739907&ObjectID=842971&ObjectType=35

DSEWPaC (2011) Australia state of the environment 2011 - Independent report to the Australian Government Minister for Sustainability, Environment, Water, Population and Communities, Canberra, ACT. http://www. environment.gov.au/topics/science-and-research/state-environment-reporting/soe-2011

Further references and resources can be located at the end of this chapter



Horticulture Australia

# **Biodiversity - references and further resources**



Environmental Taxation Concessions - administrative arrangements relating to taxation concessions that seek to conserve and protect the natural environment. http://www.environment.gov.au/topics/about-us/ business-us/tax-concessions

#### Flora of Australia Online http://www.environment.gov.au/biodiversity/abrs/online-resources/flora/main/

Gazey C. & Andrew J. (2010) Getting the soil pH profile right helps with weed control and sustainability. In: Soil Solutions for a Changing World. 9th World Congress of Soil Science pp. 232–5, Brisbane, QLD. http://www.iuss.org/19th%20WCSS/Symposium/pdf/0700.pdf

#### Greening Australia www.greeningaustralia.org.au

Grow me instead - educational initiative of the Nursery and Garden Industry Australia for addressing invasive plants http://www.growmeinstead.com.au

Identification of regional priorities - Conservation advices and Recovery Plans by NRM region http://www.environment.gov.au/cgi-bin/sprat/public/conservationadvice.pl

Lindenmayer, D, Cunningham, S & Young, A (2012) 'Land use intensification: A challenge for humanity', in David Lindenmayer, Saul Cunningham and Andrew Young (ed.), Land Use Intensification: Effects on Agriculture, Biodiversity and Ecological Processes, CSIRO Publishing, Australia, pp. 1-4. http://www.publish. csiro.au/pid/6808.htm (purchase via this link)

Lindenmayer, D & Gibbons, P, eds, (2012) Biodiversity Monitoring in Australia, CSIRO Publishing, Australia. http://www.publish.csiro.au/pid/6770.htm (purchase via this link)

Lindenmayer, D.B., Archer, S., Barton, P.S., Bond, S., Crane, M., Gibbons, P., Kay, G., MacGregor, C., Manning, A.D., Michael, D., Montague-Drake, R., Munro, N., Muntz, R. & Stagoll, K. (2011) What makes a good farm for wildlife?, CSIRO Publishing. http://www.publish.csiro.au/pid/6450.htm (purchase via this link)

Lindenmayer DB et al. (2003) Wildlife on farms - how to conserve native animals, CSIRO Publishing. http://www.publish.csiro.au/pid/3547.htm (purchase via this link)

Macfadyen S, Cunningham SA, Costamagna AC, Schellhorn NA. (2012) Managing ecosystem services and biodiversity conservation in agricultural landscapes: are the solutions the same? Journal of Applied Ecology 49:690-694.

Natural Resource Management Ministerial Council (2010) Australia's Biodiversity Conservation Strategy 2010-2030, Australian Government, Department of Sustainability, Environment, Water, Population and Communities, Canberra, ACT. http://www.environment.gov.au/node/14488

The Strategy is a significant national policy document that will guide how governments, the community, industry and scientists manage and protect Australia's plants, animals and ecosystems over the next twenty years.

Natural Resource Management Ministerial Council (2009) Australia's Strategy for the National Reserve System 2009–2030, Department of Sustainability, Environment, Water, Population and Communities, Canberra. ACT. http://www.environment.gov.au/system/files/pages/643fb071-77c0-49e4-ab2f-220733beb30d/files/nrsstrat.pdf

Natural Resource Management Ministerial Council (2007) Australian Pest Animal Strategy – A national strategy for the management of vertebrate pest animals in Australia, Department of Sustainability, Environment, Water, Population and Communities, Canberra, ACT. http://www.environment.gov.au/system/files/resources/ a7cb5991-e5c1-4c57-9037-1fd053ac8f2b/files/pest-animal-strategy.pdf

NSW Department of Primary Industries, Soil biology basics page http://www.dpi.nsw.gov.au/agriculture/resources/soils/biology/soil-biology-basics

NSW Government Office of Environment and Heritage (2011) Conservation Management Notes - Managing bushland and wildlife habitat: Restoring native vegetation, Office of Environment and Heritage, Department of Premier and Cabinet NSW. http://www.environment.nsw.gov.au/resources/cpp/RestoringVeg.pdf

NSW Government Office of Environment and Heritage (2011) Conservation Management Notes - Wildlife on your property: Integrating wildlife conservation and farm management, Office of Environment and Heritage, Department of Premier and Cabinet NSW. http://www.environment.nsw.gov.au/resources/cpp/ WildlifeFarmMan.pdf

Queensland Government Department of Environment and Heritage Protection - Nature refuges initiative http://www.ehp.qld.gov.au/ecosystems/nature-refuges/

Parks and Wildlife Service, Tasmania – Living with wildlife http://www.parks.tas.gov.au/index.aspx?base=2958

Platt, S.J., (2002). How to Plan Wildlife Landscapes: a guide for community organisations. Department of Natural Resources and Environment, Melbourne. http://live.greeningaustralia.org.au/nativevegetation/pages/ pdf/Authors%20P/4\_Platt.pdf

Radcliffe, E.B; Hutchison, W.D; & Cancelado, R.E. [eds.] Radcliffe's IPM World Textbook, URL: http://ipmworld. umn.edu, University of Minnesota, St. Paul, MN.

SARDI - Revegetation by design - Revegetation at property scale-designing the 'right' biodiversity for sustainable production. http://www.sardi.sa.gov.au/pestsdiseases/research\_projects2/research\_projects/ revegetation\_by\_design

Straker, A. and Lowe, K.W. (2004) Native Biodiversity Resource Kit – Environmental management in agriculture, Department of Sustainability and Environment, Melbourne. http://www.depi.vic.gov.au/agriculture-andfood/farm-management/legal-information-for-victorian-landholders/biodiversity-management or http://live. greeningaustralia.org.au/nativevegetation/pages/pdf/Authors%20S/12\_Straker\_Platt.pdf

Straker, A. and Platt, S. (2002) Living Systems Resource Kit – Biodiversity in Property Management Planning. Department of Natural Resources and Environment, Melbourne. http://live.greeningaustralia.org.au/ nativevegetation/pages/pdf/Authors%20S/13\_Straker\_Platt.pdf

Taverner, P; Wood, G; Jevremov, D; and Doyle, B. (2006) Revegetation by design Handbook: A guide to using selected native plants to reduce pests and diseases in the horticulture region of the Northern Adelaide Plains, SARDI. http://www.sardi.sa.gov.au/\_data/assets/pdf\_file/0008/44945/reveg\_by\_design\_guidebook.pdf

Tracey, J., Bomford, M., Hart, Q., Saunders, G. and Sinclair, R. (2007) Managing Bird Damage to Fruit and Other Horticultural Crops. Bureau of Rural Sciences, Canberra. http://www.dpi.nsw.gov.au/\_\_data/assets/ pdf\_file/0005/193739/managing\_bird\_damage-full-version.pdf

Wardlaw, F. (2004) Vegetable Integrated Pest Management (IPM) in Tasmania: Manual, Department of Primary Industries, Water and Environment, Tasmania. http://trove.nla.gov.au/work/20112924

Weeds Australia http://www.weeds.org.au

Weeds of National Significance www.weeds.org.au/WoNS

Wildlife Friendly Fencing http://www.wildlifefriendlyfencing.com/WFF/Home.html organisations involved in soil and plant tissue analysis.

Wood, G. (2010) Revegetation at property scale - designing the 'right' biodiversity for sustainable vegetable production (HAL Project Reference VG06014). South Australia Research & Development Institute (SARDI). Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au



5.14 Chapter 5 Biodiversity

# **Biodiversity - references and further resources**





# Chapter 6 Waste management



Objective – waste products are avoided, minimised, reduced, reused or recycled wherever feasible or are disposed in a manner in line with community expectations and legislation

Global outlooks suggest that the expansion of agricultural production is likely to slow, at least in the medium term, with limited area expansion and slower productivity growth, but supply should keep pace with demand such that prices that are expected to remain relatively high. In this context, measures to reduce food loss and waste will be important in meeting rising demand and for increasing productivity.

Production processes create waste. This section covers waste that is created as part of horticultural operations. This waste is usually put into landfills/tips, which is not good use of valuable space and can result in other environmental impacts such as creation of greenhouse gases and pollution of groundwater (e.g. chemicals and nutrients).

Wastes should be managed in accordance with the following order of preference (source: waste hierarchy from Environment Protection Act 1970):

- a) Avoidance;
- b) Reuse;
- c) Recycling;
- d) Recovery of energy;
- e) Treatment;
- f) Containment;
- g) Disposal.

Further references and resources can be located at the end of this chapter.

One of the key benefits associated with recycling is the avoidance of landfill. When organic waste (food, garden clippings, paper, timber) is treated in landfill, gases are emitted that contribute to green house gases emission. As organic matter breaks down in landfill both biogenic carbon dioxide ( $CO_2$ ) and methane ( $CH_4$ ) are emitted. Methane is the most important of these gases from a green house gases perspective because it has a high global warming potential (21–25 times that of  $CO_2$ ). Biogenic  $CO_2$  is not considered a source of anthropogenic green house gases because it is derived from natural sources and would be produced as part of natural cycles.

This chapter does not address water or chemical waste. For more information on these topics see 2b – Water quality, 3 – Chemical management, and 4b – Nutrient application.



Horticulture Australia

Further references and

the end of this chapter.

HAI

resources can be located at

## 6 Waste management



Review potential sources of waste oil, metal, paper and packaging, timber, glass, plastics, produce.

## HIGH RISK

Take actions to minimis, reduce and recycle. Review the Suggested Practices in this chapter.

## HIGH RISK

Take actions to minimise, reduce, re-use and recycle. Review the Suggested Practices in this chatper.

LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

**HIGH RISK** – You need to take some action. Read the Suggested Practices for that chapter. Further references and resources can be located at the end of this chapter.





## **Review checklist**

To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

#### Identify and prioritise waste products

The first step to managing waste is to determine the types of waste produced by your operations. Examples of waste products include:

Inert materials:

- Metal (car bodies etc.);
- Rubble; and
- Glass (building materials, bottles).

Persistent materials:

- Timber (wooden bins, pallets, crates);
- Packaging (waxed or unwaxed cartons, polystyrene boxes, plastic film, net wrap);
- Plastic (seedling trays, fertiliser and seed bags, mulch, irrigation drip tape; irrigation pipes); and

Tyres.

Biodegradable materials:

- Paper and cardboard (office paper waste, packaging);
- Substrate (any growing medium used in place of soil; for example potting mix, peat);
- Spent hydroponic solutions;
- Reject plants and vegetative waste; and
- Reject (unmarketable) produce.

Toxic materials:

- Waste oil;
- Batteries;
- Waste pesticide/chemical liquids (dip solution, rinsates, etc.); and
- Treated timber.

Are the wastes a danger to humans or the environment? Are the wastes subject to government regulation?

Once you have identified wastes, it is then useful to prioritise them. This can be done by considering the amount of that waste generated by the operation and the potential impact of the waste on the environment. Further references and resources can be located at the end of this chapter.



Waste avoidance or minimisation

After identifying and prioritising wastes from your operation, some sort of waste disposal plan will help you determine how you are going to deal with these wastes. Can they be eliminated, reduced, substituted for another less wasteful product, recycled or is the only option to send to conventional landfill? Sometimes you might not currently have too many options for dealing with wastes, particularly in remote areas where, for example, recycling options may be limited.

One option to reduce packaging is to opt for bulk supplies of inputs where appropriate. The exception is pesticides and fertilisers, where it is good practice to keep stored supplies to a minimum.

Minimising waste can have a positive financial impact, and is a matter of looking closely at what gets thrown out and how things are done to find opportunities to minimise the creation of waste in the first place. Take waste disposal into account when choosing products. Wherever possible choose methods and equipment that offer extended life and produce relatively low amounts of waste for disposal. Consider using materials that biodegrade after they have been used.

#### Reuse or recycle

Materials can be reused within the operation or sent for recycling. For instance, wooden bins can be repaired rather than sent to waste. Storage areas can be established for materials such as timber and steel. Materials being sent for recycling (e.g. paper, oil, glass, timber, steel) need to be collected and separated into dedicated recycling containers or areas for pickup. The local council may have recycling facilities in conjunction with the rubbish tip, or a local charity may collect materials for recycling. When donating waste materials to outside groups or organisations, ensure they are safe before releasing them. Consider distributing out-of-specification produce that is safe to be consumed to charity organisations, such as Fareshare, FoodBank, OzHarvest and SecondBite. Some of these organisations will collect from the farm.

Consider recycling substrates, particularly peat-based products, as not all sources of peat are environmentally sustainable.

Consider composting waste vegetation and produce. The composted product can be returned to production areas as a soil ameliorant. Waste produce can also be returned uncomposted to fallow areas. If recycling waste produce as feed for livestock, ensure it does not contain unacceptable chemical residues.

Ensure waste produce composting or dumping areas are well away from packing and handling facilities to avoid re-contamination of harvested produce with disease, and to avoid attracting vermin to the packing facility.

Also consider environmental impacts of compost sites, such as nutrient rich run-off and the potential for contamination of waterways (surface and groundwater).

Take advantage of returnable packaging systems, for example returnable bulk fertiliser bags.

Consider reusing plastic materials. If an item can be used several times before it becomes unserviceable, the quantity of material that needs to be disposed of will be greatly reduced. To maximise recycling, take care when handling and using plastics.

If plastic items such as plant trays are reused, choosing more durable products can increase their life.

Waste oil from farming activities may be contaminated with substances such as metal particles from engine wear, fuel from incomplete combustion, rust, dirt, carbon, heavy metals and water. If not dealt with effectively, waste oil can lead to pollution of the environment and potential risk to public health and safety. Wherever practicable, waste oil should be recovered for reuse and recycling. It should be stored in a leak-proof

## 6 Waste management

Some States have websites that assist in finding recycling options, See References and Further Resources.

Controlled/ prescribed wastes are any wastes that are hazardous to human health or the environment either directly or indirectly. These can be, for example, flammable, corrosive, toxic or give arise to gases that have these properties. If in doubt as to whether a waste is controlled or not, contact your local environment agency.

Further references and resources can be located at the end of this chapter.



container in a bunded area prior to collection by a reputable recycling business or delivery to a recognised disposal facility such as a local government collection depot or service station. Waste oil must never be applied to roadways as a dust mitigation strategy.

Contacting the manufacturer to see if a recycling system is in place is also a good option, for instance 'Netafim' has a cost-neutral\* recycling solution available to their growers whereby they supply a recoiling machine to the grower and a freight service for the coils to the processor. See www.netafim.com.au or contact Adam Roberts, Netafim Melbourne Office for more information on this service.

\*Note: there is no charge for freight only when sufficient quantity is at one location.

#### Disposal

Disposal of materials should be the last resort and can include burning or burial in landfill (onsite or council disposal facility). There may be regulations related to burning of certain types of waste, particularly wastes that are defined as "controlled" or "prescribed" wastes. Controlled wastes include items such as agricultural chemicals and chemical containers, tyres and oil. These wastes need to be carefully managed and are closely regulated because of their potential adverse impacts on human health and the environment. Some controlled wastes, such as tyres, are not strictly hazardous but they may also need special management.

Consideration needs to be given to the other potential environmental impacts associated with disposal, such as creation of dark smoke and pollution of groundwater.

If disposing of waste materials on site, do not bury or dump them close to waterways or in a way that run-off or leachates from the waste material can contaminate waterways or groundwater.



Monitoring and recording

Records that can (and in some cases must) be kept include:

- Waste management plans (can be as brief as a couple of sentences indicating major sources of waste and strategies taken to address them);
- Official receipts, offered to participants in the drumMUSTER program when they bring drums in for disposal. This is a signed document distributed through authorised inspectors at official drumMUSTER collection sites, listing the number of drums brought in for disposal. The receipt provides proof of participation in drumMUSTER and therefore proof of responsible disposal;
- ChemClear<sup>®</sup> documentation, issued to prove chemicals have been booked in for collection and also when chemicals are collected; and
- Receipts and invoices from recycling or commercial disposal businesses.

Disposal of surplus agricultural chemicals from the spray vat can be recorded on spray records.

The effectiveness of waste management can be assessed through water and soil tests.



#### **References and further resources**

For access to relevant references and further resources click here.

#### Waste management - references and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www.horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

Australian Government Department of Environment (2009) National Waste Policy http://www.environment.gov.au/topics/environment-protection/national-waste-policy

Australian Government Department of Environment - Used Oil Recycling http://www.environment.gov.au/ node/21308

Charities that recycle/redistribute surplus fresh produce in Australia: Fareshare (VIC only) http://fareshare.net.au

- Foodbank http://www.foodbank.org.au
- OzHarvest Food Rescue www.ozharvest.org
- SecondBite food for people in need http://secondbite.org

ChemClear® - Collection of unwanted chemical or out of date chemical, handling and storage tips www.chemclear.com.au

DrumMUSTER® - Collection of empty chemical containers www.drummuster.com.au

DECCW (2010) Environmental benefits of recycling, Sydney South: Department of Environment, Climate Change and Water, NSW. http://www.epa.nsw.gov.au/resources/warr/1058BenefitsOfRecycling.pdf

DEPI (2013) Getting Full Value, Melbourne: Department of Environment and Primary Industries. http://www.depi.vic.gov.au/environment-and-wildlife/sustainability/waste-management-and-resource-recovery

DSEWPaC (2012) Waste and Recycling in Australia, Canberra: Department of Sustainability, Environment, Water, Population and Communities. Report prepared by Hyder Consulting. http://www.environment.gov.au/resource/ waste-and-recycling-australia-2011-incorporating-revised-method-compiling-waste-and

FoodWise http://foodwise.com.au/foodwaste/food-waste-fast-facts/

Stanley, R. (2011) Commercial feasibility of banana waste utilization in the processed food industry (HAL Project Reference BA09025). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

#### State-specific information

- ACT http://www.environment.act.gov.au/waste
- NSW http://www.epa.nsw.gov.au/waste/index.htm
- NT http://www.ntepa.nt.gov.au/waste-pollution QLD http://www.ehp.qld.gov.au/waste/index.html
- SA http://www.epa.sa.gov.au/environmental\_info/waste
- TAS http://epa.tas.gov.au/epa/resource-recovery-and-waste
- VIC http://www.epa.vic.gov.au/your-environment/waste
- WA http://www.epa.wa.gov.au/Pages/default.aspx

Sustainability. Victoria - Integrated waste management http://www.sustainability.vic.gov.au/en/Our-Priorities/ Integrated-Waste-Management

National Technical Committee for Organics Recycling (2004) Best Practice Guidelines Series Composting, Edition 1: February 2004, Waste Management Association of Australia www.wmaa.asn.au (membership required to access publications).

NSW DPI (2003) How to compost on farm NSW DPI Agnote DPI-448. New South Wales Department of Primary Industries, NSW. http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0003/166476/compost-on-farm.pdf

OECD/FAO (2013) OECD-FAO Agricultural Outlook 2013-2022. OECD Publishing and FAO, Paris. http://www.oecd.org/site/oecd-faoagriculturaloutlook/

Wildman, H. (2008) Horticultural waste reduction and conversion through microbial bioremediation (HAL Project Reference HG06024). Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

ZeroWaste SA http://www.zerowaste.sa.gov.au/at-home/fact-sheets

Horticulture Australia

Further references and

the end of this chapter.

resources can be located at

# Waste management - references and further resources




# Chapter 7 Air management



Objective – air pollution from odours, dust, smoke, noise and greenhouse gases is minimised

Air pollution issues, particularly odours, dust, smoke and noise, can often be of most significance to your immediate neighbours. Disputes involving environmental nuisance can arise as a result of the breakdown of good neighbourly relations. Considering impacts of farming activities on neighbours and, where appropriate, discussing aspects of farming with neighbours can help in their understanding of primary production. Similarly, primary producers need to recognise that some activities can negatively impact on their neighbours and that at times it may be appropriate to adjust activities as far as reasonable to minimise the impact.

Note that greenhouse gases are covered in Chapter 8 – Energy & greenhouse gas management.

Further references and resources can be located at the end of this chapter.

## 7a Odour management



# Objective – odours from horticultural enterprises are managed to minimise potential conflicts

Odours can be caused by animal manures, fertilisers and chemicals, waste disposal sites for produce, composting sites and activities, mulches and waste management equipment.





### Review checklist

To go straight to the worksheet for this chapter click here.



### Relevant legislation and regulation

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### Suggested practices

#### Manure

For information on how to compost refer to NSW DPI Agnote on How to compost on farm and other documents in the References and other resources section.

Animal manure is often stored and used in horticulture. However, most people do not like the smell of raw manure. Growers must make sure manure is stored and used in a way to minimise the nuisance to neighbours.

#### Replacing raw manure

Growers can reduce the nuisance to neighbours by replacing raw animal manure with other less odorous products such as composted or dry manure. Although more expensive, using already composted manure greatly reduces any offensive odour. If fresh manure is used, growers must store and use it in the correct way.

#### Manure, fertiliser and chemical storage

Manure storage areas should be as far from neighbours as possible. If possible, storage areas should be located to prevent prevailing wind causing an odour issue for neighbours.

Visually screening the storage area can reduce the perception of odour problems. Providing a natural or artificial barrier between the storage area and the public eye can be very beneficial. Good ways of making a visual screen include planting a thick row of trees or putting up high solid fences.

Protection of stored manure (and compost) from rain, and containment of run-off effluent can avoid contamination of adjacent soils, work areas and waterways.

#### Manure application

Unfortunately, you cannot always keep a large distance between the area where manure is spread and the neighbours. The first thing that you can do is to always contact neighbours before spreading manure. This gesture will probably go a long way in maintaining good relationships with the public.

Whenever possible, you should schedule times to spread manure when it will have the least impact on neighbours. In general, manure spreading should be done on weekdays during office hours. At these times, neighbours are more likely to be away from home. If possible avoid spreading manure on weekends, holidays or when social events are taking place.

Horticulture Australia

Further references and

the end of this chapter.

resources can be located at

## 7a Odour management

Further references and resources can be located at the end of this chapter.



Carefully consider the weather before spreading manure. Manure should not be spread when the wind direction is going to take the smell to a neighbour. Even under appropriate weather conditions, the more manure used, the more likelihood that the smell will be excessive. Extra manure will not provide additional benefits to the crop and will only make the smell worse. Odours are reduced and drying is quicker when less manure is used and the manure is well spread out.

Another good practice is to dig the manure into the soil as quickly as possible. This practice also has positive implications for food safety. Ideally, manure should be incorporated as soon as spread. The best method is to incorporate it as it is put on, or to inject it. Adequate soil moisture to allow rapid initial breakdown of the applied manure helps to reduce odour. This is not possible or desirable in some permanent crop situations. Managers need to assess the risk of odour problems against the risk of soil compaction and loss of groundcover exposing the soil to erosion from incorporating manure.

As with the storage area, natural and man-made barriers between production areas and neighbours can greatly reduce the likelihood of complaints.

#### Produce waste disposal sites

Packing sheds often have produce waste disposal areas, which can produce odours. The same principles apply as for storage of animal manures:

- Disposal areas should be as far away as possible, and if possible, downwind of neighbours;
- Visual screening of the disposal area reduces the perception of odour problems.

Ideally, waste disposal sites should be regularly covered with soil to minimise odours and the risk of disease transmission, pest build-up and vermin.

Composting waste is also an option. Done correctly, aerated and balanced this is a low odour process. Adding gypsum to materials to be composted can cause strong sulphurous odours as the compost matures



### Monitoring and recording

A farm map can be used to document sites of any manure storage areas, produce waste sites or other odour-producing activities and proximity of these areas to neighbours. Evidence of adequate recording and/or planning is beneficial.

Odours can be monitored by visiting these areas and smelling for yourself!

A complaints register can record the nature of complaints and how they were resolved.

Records of application date and cultivation date can be useful to substantiate prompt incorporation of



### **References and further resources**

For access to relevant references and further resources click here.



Objective - to manage dust to minimise on and off site impacts

Excessive dust can cause annoyance and in some cases health problems to neighbours and staff. Dust created around packing sheds can also settle on packed produce, affecting visual quality and potentially having food safety implications.

The combination of soil type, farming system and weather patterns contributes to the risk of soil erosion by wind.



Risk assessment

### Do any of the following apply to the site:

- soil is light/prone to erosion?
- cropping/harvesting activity will leave soil exposed during windy weather?
- site is particularly exposed?



Further references and resources can be located at the end of this chapter.



Horticulture Australia



## 7b Dust management



LOW RISK – You probably don't have a significant problem in this area. You may like to read the Suggested Practices to check your understanding

HIGH RISK - You need to take some action. Read the Suggested Practices for that chapter.

Further references and resources can be located at the end of this chapter





To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

Control measures may include planting shelterbelts and windbreaks where practical, modifying cultural practices or reconsidering the appropriateness of particular cropping activities on exposed sites.

Constructing or planting a shelterbelt/windbreak will slow the velocity of wind across a site (shelterbelts/windbreaks should be designed to allow 30-50% of the wind to pass through). The protective effects from a shelterbelt/windbreak reduce with distance away from it (protection extends no more than 20 times the height of the vegetation).

Vegetation shelterbelts/windbreaks also provide wildlife habitat, assist in minimising spray drift, reduce the visual and noise impacts of site activity and can improve the quality of crops.

Choose cultivation practices carefully:

- Working soil to fine tilth in dry windy weather should be avoided if possible. Pre-irrigation to wet dry soil before cultivation will help to reduce dust;
- Use slower cultivation speeds when there is a risk of dust. Uncultivated crop stubble provides protection against wind erosion. Minimise the amount of time soil is left without vegetation or a cover crop. Minimum tillage techniques should be used where practical; and
- Inter-row spacings and headlands should have groundcover whenever possible.

Applying mulches to the surface of seedbeds after drilling on sandy soils is an effective control measure. Use of plastic mulch along plant rows will also contribute to dust control.

Wetting down, sealing and use of 'minimal dust materials' (for example blue metal or hardstand) for the surfaces of frequently used traffic ways (transport delivery and pickup areas, harvested produce delivery points and forklift routes at the packing shed) will dramatically reduce the dust problem. Do not apply oil to trafficways due to the potential for it to end up in waterways.



Monitoring and recording

See Topic 1a - Land and soil management, for details of ways to assess soil erosion by wind.

If dust is a major issue to neighbouring areas, cultivation records may be kept (as part of paddock records) to detail soil conditions when soils are worked and the equipment used.

Farm maps should show roads, sensitive areas and soil types and can be used to demonstrate the placement of current and planned shelterbelts and the direction of prevailing winds.

The effectiveness of these practices can be assessed by observing dust levels on windy days.



**References and further resources** 

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter



Horticulture Australia

## 7b Dust management

Further references and resources can be located at the end of this chapter



7.7

Chapter 7 Air management

# 7c Smoke management



Objective - to manage smoke to minimise on and off-site impacts

The on and off-site impacts of smoke need to be minimised through fire management techniques.



Risk assessment





#### **Review checklist**

To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

If burning in the open is the only practical method of disposal of materials, and it is legal in your area, precautions should be taken to prevent producing dark smoke and causing a nuisance. Burning of waste is banned, or requires a permit, in many local government areas. Before burning check local bushfire restrictions/permits and local government restrictions and requirements.

Do not burn plastics, rubber, tyres or other materials known to produce dark smoke. Where possible recycle, reuse or dispose of these waste items at local authority waste depots. Do not burn plastics or chemicals under any circumstances as they release toxic fumes and residues.

Avoid burning if it will cause a nuisance to nearby residential areas. Check wind direction before burning - only burn when wind direction is away from neighbours.

As a courtesy, inform immediate neighbours before burning.

Be aware of localised landscapes that can induce smoke problems, such as valleys.

Materials should be dry and have low moisture content. Do not burn green vegetation. Keep fires small and continually add combustible material, minimising the depth of the combustion area. Minimise the quantity of incombustible material added to the fire. Wherever possible keep incombustible materials separate from materials to be burnt. For better combustion, agitate the base of the fire to improve air supply.

If fire produces dark smoke, don't add any more material that burns slowly.



### Monitoring and recording

Further references and resources can be located at the end of this chapter.



The quantity of dark smoke produced during burning operations can only be assessed visually. Records of other disposal options for materials that are likely to produce dark smoke can also be retained.



### **References and further resources**

For access to relevant references and further resources click here.

# 7c Smoke management

Further references and resources can be located at the end of this chapter





Objective - to manage noise to minimise on and off-site impacts

Noise is a form of pollution that may result in environmental harm. Assessments of environmental harm include consideration of:

- The degree and scale of impact;
- The health and safety of people;
- Property damage;
- Unreasonable interference with the amenity of an area.

The nature of the pollution and the sensitivity of the receiving environment is also a consideration. Noise pollution could be associated with excessive noise from farm machinery (such as pumps, harvesters and vehicles), bird scaring devices, and/or frost protectors. This could impact on people's enjoyment of their property, may affect their mental or physical health, and may have adverse impacts on local communities.

Noise many not seem like an environmental management issue for growers, however most State legislation for environmental protection includes noise as part of the definition of the environment. For this reason, noise management is included in the environmental assurance process for horticultural businesses.



HA

# 7d Noise management

Further references and resources can be located at the end of this chapter





To go straight to the worksheet for this chapter click here.



#### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



#### Suggested practices

Identify and consider local government regulations.

Buffer zones are useful to reduce noise and are also helpful to mitigate impacts of off-target spray application.

Where pumps are located close to residential areas, consider changing from diesel to electric pumps or creating a sound barrier around the pump. Electric pumps will most likely be run at night time, when electricity tariffs are lower.

Consider muffling equipment where daytime intermittent noise levels are excessive. Where normal methods are not sufficient to reduce noise to acceptable levels, equipment that is continuously operated may require soundproofing or artificial mounds to help absorb and deflect the noise.

Some forms of seasonal activity, or current and accepted industry practice like harvesting, may require the use of machinery at night. Where sensitive places are close to noise and night-time activities occur, consider starting work closer to the sensitive area and moving away as night falls. The converse applies for early morning activities.

Where noise may be an issue, keeping records of machinery use may be beneficial.

Use of bird deterrents, such as gas guns, and frost protectors in a considerate manner and in accordance with local bylaws, such as meeting maximum accumulated peak level (APL) for impulsive noise devices and guidelines on hours of operation.

Transport operators picking up packed produce should be reminded not to use exhaust brakes where this noise would create a nuisance to neighbours.

Given the potential for disturbance and misunderstanding associated with noise pollution, it is sound practice for operators to consult with neighbours about their planned operation, to explain the reason for their use, and to discuss options for their operation. In some cases this may not be feasible (e.g., the number of people involved, uncertainty about who may be able to hear the devices in different conditions, or historic bad relations) but, if possible, it could help in their understanding of the activity and potentially prevent some problems from arising.



Monitoring and recording

Records of machinery maintenance should be maintained. In situations where a more definitive measure of noise is required, organisations such as state OH&S agencies can assist with testing.



#### References and further resources

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

## 7d Noise management

Further references and resources can be located at the end of this chapter.



#### Air management - references and further resources

(web links accurate as at 11 February 2014)

reference or final r www.ho	number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as sees within this document. This is by no means representative of all the research & development (R&D) reports available in this area. For full list of HAL final reports visit the HAL website rticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on a & development outcomes specific to your industry.			
	urces Board, California, Smoke Management Program <mark>ca.gov/smp/smp.htm</mark>			
	an Centre for Agricultural Health and Safety – Noise Injury Prevention ww.aghealth.org.au/index.php?id=5037			
	an Government Department of Environment - Air Quality ww.environment.gov.au/topics/environment-protection/air-quality			
	an Government Department of Environment – Smoke from Biomass Burning ww.environment.gov.au/resource/smoke-biomass-burning			
	of Meterology ww.bom.gov.au			
	th Australia (2007) Draft Environmental Noise Guidelines for Audible Bird Scaring Devices, Adelaide, SA. ww.epa.sa.gov.au/xstd_files/Noise/Guideline/guide_bird.pdf			
	2011) Protecting our water, soil and air, Department of Environment Food and Rural Affairs, UK. www.gov.uk/government/publications/protecting-our-water-soil-and-air			
	DEHNSW (2012) DustWatch, Department of Enviornment and Heritage, Sydney, NSW. http://www.environment.nsw.gov.au/dustwatch			
	MODIS – Land Atmosphere Near Real-Time Data – NASA https://earthdata.nasa.gov/data/near-real-time-data/rapid-response/modis-subsets			
	NSW Department of Primary Industries (2003) How to compost on farm, Agnote DPI - 448 http://www.dpi.nsw.gov.au/data/assets/pdf_file/0003/166476/compost-on-farm.pdf			
	overnment – Air Quality Monitoring ww.environment.nsw.gov.au/aqms/index.htm			
	d Organics ycledorganics.com			
	rk Australia – Agricultural Resources (noise management) ww.safeworkaustralia.gov.au/sites/swa/whs-information/agriculture/pages/agriculture			
State-sp	pecific information			
NSW	http://www.epa.nsw.gov.au/air/			
NT	http://www.ntepa.nt.gov.au/waste-pollution/air			
QLD	http://www.ehp.qld.gov.au/air/index.html			
SA	http://www.epa.sa.gov.au/environmental_info/air_quality			
	http://epa.tas.gov.au/epa/air			
TAS				
TAS VIC WA	http://www.epa.vic.gov.au/your-environment/air/air-pollution http://www.epa.wa.gov.au/Pages/default.aspx and			





# Chapter 8 Energy & greenhouse gas management



#### Objectives

- to identify and reduce energy inputs wherever feasible in the production system
- to ensure sources of greenhouse gases are identified and emissions are reduced wherever feasible

Although energy from sunlight is essential for plant growth, the energy balance of agricultural systems depends on additional energy, from non-renewable sources, to power machinery. Sustainable practices can improve the balance of energy and contribute to efficient energy use.

Most of our energy (including more than 90% of our electricity) comes from fossil fuels such as oil, coal and gas. Burning of fossil fuels releases carbon dioxide (CO<sub>2</sub>), nitrous oxide (N<sub>2</sub>O) and methane (NH<sub>4</sub>) into the atmosphere. This has resulted in the greenhouse effect known as 'global warming'.

All the major greenhouse gases have natural sources, but human activity increases the amount released. It is now generally agreed that this pollution has added to the natural greenhouse effect and will cause the temperature of the earth's surface to rise. Because some of the greenhouse gases last for a long time in the atmosphere, it is important that any action to reduce emissions of greenhouse gases is taken as early as possible.

Greenhouse gases of greatest concern in agriculture are methane and nitrous oxide. Methane arises from livestock, mainly through digestion (enteric fermentation). Nitrous oxide is released from soils through application, oxidation and natural transport of fertilisers and soil disturbance. Carbon dioxide emissions are released from transport and energy sources. Chlorofluorocarbons or CFCs (used for refrigeration and aerosol propellants) were also significant greenhouse gases but are now generally prohibited from use.

Further references and resources can be located at the end of this chapter.

An important issue is the high global warming potential of methane and nitrous oxide, expressed as 'carbon dioxide equivalents' or CO2-e. The CO22-e of CH4 is 21 while the  $CO_2$ -e of N<sub>2</sub>O is 310. This means that methane has a global warming effect that is 21 times greater than carbon dioxide while that of nitrous oxide is 310 times greater! The implication is that we should be especially careful to avoid methane and, particularly, nitrous oxide emissions.

The Australian agricultural industry as a whole accounts for 16-18% of greenhouse emissions. Although horticulture accounts for only 1% of Australia's total greenhouse gas emissions (not including fossil fuel use), it tends to have greater emissions intensity per hectare of land than other agricultural production enterprises.

Sources of greenhouse gases in horticulture include: fuel and electricity use (70%), especially for irrigation, nitrogen fertilisers and animal manures (20%), and waste and refrigerant loss to the atmosphere (10%). Methane may be produced from composting processes and stagnant pools of contaminated water, it appears not to be a major concern.

Australia's mechanism to reduce emissions, the carbon price, came into effect in July 2012. While agricultural industries are not directly involved in the carbon price mechanism, the carbon price will impact on the cost of key agricultural inputs. These price increases have been and will continue to be most significant for energy (i.e. electricity and fuel) and energy-intensive inputs such as fertiliser and chemicals.

Note: As at 11 February 2014 the carbon price mechanism was still in place, but legislation to repeal the carbon price was also being considered. Regardless, some mechanism for reducing Australia's emissions will be in place in order to meet the bipartisan agreed emissions target of 5% from 2000 levels by 2020 and 60% by 2050. Consequently, practices to reduce energy use will continue to be important. See http://www.climatechange.gov.au/reducing-carbon/reducingaustralias-emissions for more up to date information.

Improvements in the efficiency of energy use, more effective fertiliser management and minimizing soil disturbance are key actions for the industry to reduce greenhouse gas emissions and in reducing the operating costs of horticulture operations.



Horticulture Australia

## 8 Energy & greenhouse gas management

Further references and resources can be located at the end of this chapter







- undertake regular maintenance of all equipment, particulary that
- requiring fossil fuels and CFCs?
- regularly check insulation?
- strategically apply nitrogenous fertislisers?
- minimise unneccessarry journeys and cultivation passess?







To go straight to the worksheet for this chapter click here.



### **Relevant legislation and regulation**

Legal requirements are subject to change. Regularly check with Federal, State and Local authorities for updated requirements. See here for links.



### **Suggested practices**

By understanding your emissions profile, you can identify opportunities for reducing energy use and emissions.

In assessing the total energy used in agricultural production 'from field to fork', synthetic fertiliser and pesticide manufacture are important indirect energy inputs, while irrigation, cultivation, harvesting and transport are major direct energy inputs. By managing the amount of fertiliser and pesticide applied as well as the more tangible direct agricultural activities, greenhouse gas emissions can be reduced.

A carbon footprint is a measure of the greenhouse gas emissions produced by a particular product or activity. Carbon footprinting is a method of carbon accounting.

The carbon footprint of a horticulture enterprise would account for all on-farm emissions of carbon dioxide ( $CO_2$ ), nitrous oxide ( $N_2O$ ), methane ( $CH_4$ ) and hydroflourocarbon, ie. all emissions occurring within the farm boundary from activities such as energy and fertiliser use.

There are currently several examples of carbon footprinting tools available for use within horticulture industry, including:

- Vegie carbon tool http://vegiecarbontool.com.au/
- Cool farms tool http://www.coolfarmtool.org/CoolFarmTool
  - Nursery industry carbon calculator http://www.ngia.com.au/ Category?Action=View&Category\_id=466
- Nursery industry renewable energy calculator www.energycalc.ngi.org.au
- FarmGAS calculator (includes ability to select horticulture crops within the model, but was not specifically developed for horticulture) http://calculator.farminstitute. org.au/login
- Hort Carbon Info http://www.horticulture.com.au/areas\_of\_Investment/ Environment/Climate/climate tools.asp
- Orchard carbon tool http://www.orchardcarbontool.com.au/login. php?from=calculator

The primary reason for measuring carbon footprints is to report or account for the level of greenhouse gas emissions to a third party. At present (March 2014) there is no requirement for horticultural growers to undertake carbon accounting on their farms or to comply with carbon labeling requirements. However, consumer preferences and government policy may create the need for carbon accounting and reporting by growers in the future.

As an alternative to undertaking a full carbon footprint, consider an energy audit. Energy companies can give expert advice on how to save electricity and most can conduct an energy audit to identify areas for improvement.

Management actions in the horticulture industry to reduce greenhouse gas emissions include improving the efficiency of energy use and more effective fertiliser management.

#### Improve energy efficient to reduce carbon dioxide emissions

The most effective way of reducing carbon dioxide emissions is to use energy more efficiently and to exploit non-fossil fuels as alternative sources of energy.

#### Irrigation

Investigate options for replacing existing pumps or motors with more energy efficient models. Optimise the irrigation system to improve water use efficiency and use less energy.

Pumping water for irrigation is one of the main ways energy is used in horticultural production. Growers can use less energy and save costs by carefully choosing the type of irrigation equipment they use. Keeping irrigation equipment in good condition can also save energy. Irrigation pump engines should be serviced and well tuned. Make sure motors, switches and control panels are clean. Check connections to make sure they are tight, and lubricate moving parts. Variable speed electric motors are generally more efficient than fuel-driven pumps. Pumps will wear more quickly if water contains a lot of dirt or organic matter, leading to leaks and reduced energy efficiency. Regularly check and maintain inlet sieves.

The amount of energy used by your irrigation system depends on:

- System flow rate (larger-diameter pipes reduce flow rate losses due to friction, thus reducing pumping costs);
- Operating pressure (aim for low pressure systems, the less friction in the system the less pressure you need to start with);
- Hours of operation (utilise as much off peak power as possible in many cases off peak power is cheaper because it is generated whether it is used or not [e.g. coal and wind generation]); and
- The combined efficiency of all the components.

Horticulture Australia

Further references and

the end of this chapter.

resources can be located at

## 8 Energy & greenhouse gas management

Further references and resources can be located at the end of this chapter



Consider energy efficiency when new equipment is purchased.

Choose pumps that match the pressure flow you need. Pumps need to be used in their best operating range for best efficiency. Pumps should run at or above 65% efficiency. Some agricultural companies may be able to test your pump for efficiency. Pumps wear out over time - they may still pump water, but it is costing significantly more because of impeller wear. This test can help you know if a pumping plant is wasting energy.

#### Vehicles and equipment

Maintain and service vehicles and equipment regularly to ensure efficient operation. Well-maintained equipment operates better and costs less to run. This is good both for your business and the environment. Maintain engines by following the manufacturers' recommendations. A reduction of 5-15% in fuel consumption can be obtained by servicing air cleaners and fuel injectors regularly.

It is a good idea to have a regular maintenance program for all the equipment, machinery and vehicles used on your farm. Maintenance intervals will vary to suit levels and conditions of use for each vehicle and piece of equipment.

Choose tractors and machinery that are suitable for the tasks they will be performing. Use the lowest-powered tractor capable of doing the required job or adjust machinery (i.e. increase the width of implements) to match the tractor power.

Minimise unnecessary journeys and cultivation passes. Install a GPS on tractors to prevent overlap or missed coverage.

Install or turn on the power-save function on office equipment such as computers and photocopiers. Turn them off when you finish work.

#### Fuel

Try to save fuel. Every litre of petrol or diesel saved lowers greenhouse emissions and reduces production costs. Keep track of fuel use and set targets for saving fuel. Another good idea is to switch from diesel/petrol to LPG or compressed natural gas in cars, trucks and motorbikes. This can cut greenhouse gas emissions by 10 to 15%. Use a percentage of biofuels, which come from renewable sources.

Economise on fuel by precise control of the correct temperature regimes, using thermal screens and correct maintenance and insulation of boilers and burners.

Use electric fork lifts instead of internal combustion engine forklifts.

#### Lighting

By using energy-efficient lighting you can save money and help the environment at the same time. Compact fluorescent lighting is far more energy efficient that incandescent bulbs for use in greenhouses and work areas. For example, energy efficient compact fluorescent bulbs have about one-quarter lower wattage and about eight times the life of standard incandescent bulbs. This saves energy and lowers maintenance costs. Replacing mercury vapour yard lights with energy-efficient, high-pressure sodium lights can sometimes greatly cut electricity usage and costs.

Place movement sensors and timers on lighting systems so they are only on when required.

#### Coolrooms

Coolrooms should be properly designed and built to make sure that energy is not wasted. This includes fitting the right temperature control devices and keeping them properly calibrated. Incorrect calibration by only 1°C can greatly increase running costs and may affect the quality of stored produce.

Reduce loss of heating/cooling through effective insulation and preventing unintentional ventilation. Heated glasshouses, mushroom houses and polythenecovered structures are major users of energy. Don't overlook the huge thermal losses that occur through an uninsulated floor. A little more spent at the outset is quickly recovered. Polystyrene is the floor insulant of choice.

Once coolrooms have been built they need to be maintained and serviced. Regularly check things such as door seals, hinges and catches. This will minimise leaking of warm air into the coolrooms. The building should be checked for damage to insulation panels, roof and walls. If possible, build coolrooms with a shade roof covering or within a shed to reduce thermal loads. Windbreaks also reduce airflow onto the exterior surfaces and associated heat transfer.

Always try to remove field heat from produce (pre-cool) before storing it in the coolroom as this can greatly reduce the amount of energy used. You should also minimise the time coolroom doors are open. Don't open doors often or hold them open for a long time. If doors need to be kept open during some daily operations, use plastic door strips, automatic coolroom doors or rapid-rise curtains. These devices can help to keep warm air out of the coolroom. When forklifts are in coolrooms for extended periods, consider using electric forklifts (as opposed to gas) for heat and OH&S gains. Gas forklifts can also release gases that affect produce quality (maturity and taste).

#### **Renewable resources**

The efficient use of renewable energy resources such as hydro-electricity, solar or wind power should be targeted since the use of non-renewable sources, such as fossil fuel, is not sustainable in the long term. Switch to more environmentally friendly renewable energy sources, or what is known as 'green power', where this option is available. This electricity may cost a little more than conventional power from coal-fired power stations. However, costs can be maintained to the same total cost by improving energyuse efficiency. Reverse auctions and collective bargaining may help businesses achieve better deal from electricity suppliers. See www.greenpower.gov.au for more information.

Minimise use of fossil fuel for power generation, for example:

- Optimise field operations, including transportation from field to packhouse, carefully select equipment; and
- Ensure proper and timely maintenance of equipment.

Minimise the input of synthetic fertilisers and consider alternative organic and renewable fertiliser technologies taking into account crop needs, fertiliser cost and comparative costs (including fuel use) of delivery and spreading.

Review practicality of best current waste re-use, recycling and disposal technologies available.

Some growers and processors have large amounts of organic waste (or by-product), such as cores, skins, peels, outer leaves, tops, seeds, stems, shells, husks and other plant parts, and are looking at ways that the waste can be composted to produce methane that can be used as a fuel.

#### Other sources of carbon dioxide

Large amounts of fossil fuel are needed to manufacture nitrogen fertilisers. Only use fertilisers at the rates suitable for the cropping situation. Ensure fertiliser spreaders are properly maintained and use suitable settings for different types of fertiliser.

Carbon dioxide is also created from the breakdown of liming materials in the soil.

Using less packaging, or packaging made from recycled/ recyclable materials.



Further references and

the end of this chapter.

resources can be located at

Further references and resources can be located at the end of this chapter



Source products from companies making an effort to cut their emissions and lower the impact of the carbon price on their products.

Consider ways to minimise the time produce stays on farm post-harvest, in order to minimise the refrigeration requirements. Investigate options to reduce on-farm processing.

#### Nitrous oxide

Nitrous oxide from agriculture is released from nitrogen compounds in manures, fertilisers, crops, soils and watercourses. It is likely to be produced in oxygen-free conditions. The most effective way to reduce the release of this gas is to use nitrogen fertilisers and manures efficiently so that crop requirement is met while losses of nitrogen are minimised.

Maintaining good soil structure (see Topic 1c - Soil structure) will help to keep the soil well aerated and reduce the potential for an oxygen free environment.

#### Chloroflurocarbons (CFCs)

Apart from their effect as greenhouse gases, CFCs also damage the ozone layer. CFCs were used in refrigeration equipment but are now generally prohibited. If your equipment uses CFCs, it is important to keep refrigeration equipment properly maintained to minimise the risks of leaks of refrigerant. Whenever such equipment is serviced, make sure that no refrigerant is lost. Do not allow unused equipment to deteriorate on site. Specialist contractors can safely remove refrigerant so that it can be recycled or destroyed. CFC's have not been used in refrigeration equipment since 1995.

#### **Offset opportunities – Carbon Farming Initiative**

The rural sector is a major source of emissions, but also has a potential opportunity to play a role in Australia's mitigation (emissions reduction) effort.

However, uncertainties about the domestic emissions reduction market, the level of demand for credits, and realistic abatement and sequestration potentials make it difficult to estimate the actual economic potential for horticulture at this time. There may be opportunities for horticulture to access financial incentives for reduction of emissions or the increase of carbon sinks on farm (known as carbon farming) in the future. However, all of the available information suggests that the income potential for horticultural producers will be limited or at the very least lower than other agricultural industries.

Current on-farm offset opportunities in horticulture relate to the creation of carbon sinks (also known as carbon sequestration) by:

- Better soil management e.g. controlled traffic technologies;
- Increasing soil carbon;
- Growing native vegetation; and
- Incorporating biochar and organic matter into soil.

The most promising emissions reduction option seems to be mitigation of nitrous oxide emissions from improved fertiliser management. However, other opportunities could include reforestation, revegetation and carbon soil sequestration. Environmental plantings could be used as windbreaks or visual screens and thereby provide additional benefits to the farm.

For up to date information on the domestic policy situation visit the Clean Energy Regulator website www.cleanenergyregulator.gov.au. Specific information on the existing Carbon Farming Initiative (CFI) can be found at http://www.cleanenergyregulator.gov.au/ Carbon-Farming-Initiative/Pages/default.aspx.

Current on-farm offset opportunities relate to the creation of carbon sinks (also known as carbon sequestration) by:

- Better soil management e.g. controlled traffic technologies;
- Increasing soil carbon;
- Growing native vegetation; and
- Incorporating biochar and organic matter into soil.

#### Biochar

Considerable uncertainty still surrounds the use of biochar in farming systems given the range of production processes, the types of biochar, and the variety of soil types in horticulture. There is also uncertainty surrounding its value within a domestic carbon market as well. Work is being done to address these uncertainties and quantify the benefits.

For use of biochar see Topic 1c - Soil structure.

For more information on biochar see HAL/NSW DPI publication on biochar in horticulture (2012) http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0008/447857/DPI-BioChar-in-Horticulture.



Monitoring and recording

Energy usage can be monitored by checking electricity and fuel bills. Energy efficiency over time can be assessed against production yields or throughput.

Remember the strong relationship between energy saved and your chequebook! Other records that can be useful include:

- Coolroom and machinery maintenance records;
- Thermometer or temperature monitoring device calibration records;
- Irrigation records;
- · Fertiliser records detailing use of nitrogenous fertilisers; and
- Internal energy audits.



### References and further resources

For access to relevant references and further resources click here.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

## 8 Energy & greenhouse gas management

Further references and resources can be located at the end of this chapter



#### Energy and greenhouse gas management - references and further resources

(web links accurate as at 11 February 2014)

Note: A number of Horticulture Australia Limited (HAL)-funded project final reports have been identified as references within this document. This is by no means representative of all the research & development (R&D) or final reports available in this area. For full list of HAL final reports visit the HAL website www.horticulture.com.au. Alternatively, contact HAL or your peak industry body for more information on research & development outcomes specific to your industry.

AS/NZS (1998) AS/NZS ISO 14040:1998 (ISO 14041:1997) Environmental Management: Life cycle assessment - Principles and Framework. Homebush, Standards Australia and Standards New Zealand. www.standards.org.au

Australian Government Department of the Environment – Atmosphere page http://www.environment.gov.au/node/22791

Australian Government Department of the Environment – Carbon Farming Initiative http://www.cleanenergyregulator.gov.au/Carbon-Farming-Initiative/Pages/default.aspx

Climate Change Research Strategy for Primary Industries (CCRSPI) www.ccrspi.org.au

Climate Council - The Climate Council is an independent non-profit organisation funded by donations by the public that provides expert advice to the Australian public on climate change. http://www.climatecouncil.org.au

Carbon footprinting tools (developed for or relevant to horticulture):

- Vegie carbon tool http://vegiecarbontool.com.au/
- Cool farms tool http://www.coolfarmtool.org/CoolFarmTool
- Nursery industry carbon calculator http://www.ngia.com.au/Category?Action=View&Category\_id=466
- Nursery industry renewable energy calculator www.energycalc.ngi.org.au
- FarmGAS calculator (includes ability to select horticulture crops within the model, but was not specifically developed for horticulture) http://calculator.farminstitute.org.au/login
- Hort Carbon Info [link not currently available]
- Orchard carbon tool [link not currently available]

Climate Kelpie - round up of climate tools for Australian farmers http://www.climatekelpie.com.au

#### Climate Institute http://www.climateinstitute.org.au

Cox, J; Downie, A; Jenkins, A; Hickey, M; Lines-Kelly, R; McClintock, A; Powell, J; Singh, BP; Van Zwieten, L. (2012) Biochar in horticulture: Prospects for the use of biochar in Australian horticulture (HAL Project Reference AH11006), Horticulture Australia Ltd, Sydney, NSW. http://www.dpi.nsw.gov.au/\_\_data/assets/pdf\_file/0008/447857/DPI-BioChar-in-Horticulture.pdf

CSIRO (2011) Climate Change: Science and Solutions for Australia, CSIRO http://www.csiro.au/Outcomes/Climate/Climate-Change-Book.aspx

EcoBiz program, includes sections on energy efficiency, waste and water use efficiency

Foran, B. (1998) Looking for opportunity and avoiding obvious potholes: Some future influences on agriculture to 2050, CSIRO. http://www.regional.org.au/au/asa/1998/plenary/environmental/foran.htm#TopOfPage or www.cse. csiro.au/publications/1998/lookingforops-98-13.pdf

Garnaut, R. (2011) Update Paper 4: Transforming rural land use. http://www.garnautreview.org.au/update-2011/ update-papers.html

Grace, P. (2007) Soil Carbon – Carbon study shifts focus to nitrogen. GRDC. http://www.grdc.com.au/Media-Centre/Ground-Cover/Ground-Cover-Issue-70-September-October-2007/Soil-Carbon-Carbon-study-shifts-focusto-nitrogen

GreenPower – Accredited Renewable Energy www.greenpower.gov.au

Intergovermental Panel on Climate Change http://www.ipcc.ch/

HAL (2011) Climate Change: Managing Variability and Carbon, Horticulture Australia Ltd, Sydney, NSW. Project final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au. Or directly via http://ausveg.businesscatalyst.com/rnd/fact%20sheets/Climate%20Change%20and%20Managing%20Carbon.pdf

HAL - Horticulture climate change investment page http://www.horticulture.com.au/areas\_of\_Investment/ Environment/Climate/climate\_tools.asp

Horticulture Industry Network - Cooling Down About Electricity Costs - Apple & Pear http://www.hin.com.au/resources/cooling-down-about-electricity-costs

Klein, C; Novoa, R; Ogle, S; Smith, K; Rochette, P; Wirth, T; McConkey, B; Mosier, A; Rypdal, K; Walsh, M; and Williams, S. (2006) Chapter 11: Chapter 11: N2O Emissions from Managed Soils, and CO2 Emissions from Lime and Urea Application, IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4: Agriculture, Forestry and Other Land Use. http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4\_Volume4/V4\_11\_Ch11\_N2O&CO2.pdf

O'Connell D, Braid A, Raison J, Hatfield Dodds S, Wiedmann T, Cowie A, Littleboy L, Clark M (2013) Navigating sustainability: measurement, evaluation and action. CSIRO, Australia. http://www.csiro.au/Organisation-Structure/ Flagships/Energy-Flagship/Measuring-sustainability.aspx

Putland, D. (2012) Opportunities for the Australian horticulture in the Carbon Farming Initiative (Final Report for HAL Project Reference AH11020), Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Putland, D. (2012) The impacts of the carbon price on Australia Horticulture (Final Report for HAL Project Reference AH11019), Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au



Horticulture Australia

## Energy & greenhouse gas management - references & further resources





### Key messages

Australia's increasingly variable climate poses challenges for horticulture, given the sector's dependency on natural resources, especially water for irrigation, and temperature sensitivity. This makes horticulture inherently vulnerable to the impacts of both short-term climate variability and long-term climate change.

Climate change will lead to changes in horticulture:

- Some production regions will become more productive in the short term;
- Crops and/or varieties may need to change; and
- Some production regions will become more marginal or less productive.

The impact of climate change on horticulture will differ depending on location, industry, production season, supply chain and timeframe. However, this uncertainty of impact is not a reason for complacency or delay. Producers can respond to these challenges, especially if they focus on climate change as being another business issue requiring risk management. Growers need to be considering future resilience now and developing adaptation pathways to ensure a productive and secure future for their activities.

Many climate adaptation options for horticulture are similar to existing 'best practice' and good natural resource management, and do not require farmers to make radical changes to their operations and industries in the near term. These options can, and should, be prioritised as part of a 'no regrets' or win–win strategy for agriculture because they will provide immediate and ongoing benefits as well as preparing the sector for climate change.

Off farm, the sector needs to consider whether the infrastructure and services on which it relies on (e.g. port and wharf facilities and road networks) are being prepared for a different future climate.

Further research, development and extension is needed for the provision of appropriate data and decision support tools to allow the rapid (and cheap) assessment of climate risk and the identification of possible adaptation measures, and estimates of associated costs.

### Climate change and uncertainty

**Climate change** is the term used to refer to changes in long-term trends of environmental factors such as temperature and rainfall. These changes can be due to natural variability or as a result of human activity, or a combination of both.

Some things about Australia's future climate are fairly certain: there will be more hot days and those days will often be hotter; sea levels will rise; and extremes such as heatwaves, droughts and storm surges are likely to become more frequent and intense.

Other things are less certain: the rate and magnitude of change depends on how sensitive the climate system is to greenhouse gases. Beyond mid-century, it also depends on future emissions, making it sensitive to our success in reducing greenhouse gas emissions now and into the future. Rainfall projections are less certain. While it seems likely that the southern part of Australia will become drier, there is less confidence about northern Australia and parts of the eastern seaboard. We have limited information about how climate change will affect some extreme events such as cyclones and hail storms.

Furthermore, climate change will not proceed smoothly for a given change in heat radiation from changing greenhouse gas levels. There is a risk of abrupt changes as the climate shifts from one state to another as a result of feedbacks and thresholds in the climate system. This will mean increased risk for growers and changes in the way crops are grown.

Further references and resources can be located at the end of this chapter.



While some of this uncertainty can be reduced by better science, some residual uncertainty will always remain. Understanding the likely future changes to climate makes it possible to start drawing up action plans at national, state, regional and local levels to adapt to the most likely changes.

For further information on GLOBAL scenarios visit http://www.ipcc.ch/

In Australia, detailed climate change information is available via the Ozclim website (www. csiro.au/ozclim). Ozclim provides an interface that can produce maps and data for Australia that show the predicted values for a range of climate variables. This data can be used as inputs to crop models or to derive an estimate of the change of an important temperature variable for a key event in crop production (e.g. the timing of flowering or fruit set).

#### Climate change resources:

Intergovernmental Panel on Climate Change (IPCC) website: www.ipcc.ch United Nations Framework Convention on Climate Change (UNFCCC) website: www.unfccc.int

Australian Government Department of the Environment: www.climatechange.gov.au CSIRO http://www.csiro.au/Outcomes/Climate/Climate-Change-Book.aspx

Climate Change Research Strategy for Primary Industries (CCRSPI): www.ccrspi.org.au

Food Climate Research Network: www.fcrn.org.uk

Tyndall Centre for Climate Change Research: www.tyndall.ac.uk

UK Climate Impacts Programme: www.ukcip.org.uk

Stockholm Environment Institute: www.sei.se

### Climate change adaptation

**Adaptation** is the ability to adapt to unavoidable climate change. Adaptation can occur in a planned manner in response to known changes, or in an autonomous manner as farming systems change gradually over time.

For horticulture, adaptation is a response to the impacts of changing environmental factors such as temperature and rainfall on production systems. It should include both strategic preparation and tactical responses to suit the range of likely climates and variables.

As global climate change progresses, historical conditions may become increasingly less pertinent as a guide to industry activities or industry adjustment. Industry and farm managers will need to distinguish between 'old climate expectations' and 'new climate realities' in determining and implementing new adaptation strategies or options.

Successful adaptation incorporates the development of flexible, risk-based approaches that deal with future uncertainty and provide strategies that are robust enough to cope with a range of possible local climate outcomes and variations.

#### Climate change adaptation resources:

UNFCC on adaptation unfccc.int/adaptation/items/4159.php CSIRO Climate Adaptation Flagship: http://www.csiro.au/Organisation-Structure/ Flagships/Climate-Adaptation-Flagship.aspx

National Climate Adaptation Research Facility http://www.nccarf.edu.au/ and specifically the Primary Industries Adaptation Research Network www.piarn.org.au Australian Government Department of Agriculture http://www.daff.gov.au/climatechange/ climate/adaptation-strategies

WeADAPT: www.weadapt.org

#### Some state based policy initiatives can be found at:

NSW	http://www.environment.nsw.gov.au/climatechange/a
SA	http://www.sa.gov.au/topics/water-energy-and-enviro change/adapting-to-climate-change
QLD	http://www.ehp.qld.gov.au/climatechange/centre/inf
TAS	http://www.climatechange.tas.gov.au
VIC	http://www.climatechange.vic.gov.au/adapting-to-cli Victorian-Climate-Change-Adaptation-Plan
WA	http://www.walgaclimatechange.com.au/policy.htm

Increasingly adaptation plans are being developed at a local or regional level. The NCCARF local government portal is an entry point for more information on adaptation strategies at a local level http://www.nccarf.edu.au/localgov/ Industries are also developing climate resources. See http://www.vegetableclimate.com

### Climate risk management framework

The first step for adapting horticulture to climate change involves helping producers to understand why adaptation is needed now – as decisions made now will either increase or decrease vulnerability to future climate change.

A risk management framework is a structured approach to identifying impacts, risks, opportunities and desirable actions. This approach has been widely used in many contexts (such as workplace health and safety) but can be applied in assessing risk of climate change impacts.

There are examples of climate change risk management frameworks in other agricultural sectors, such as the Climate Change Risk Management Matrix developed for the beef industry. Because of the diversity and complexity of horticulture sector it has been too difficult to develop a generic climate risk management framework.

However, the following diagram describes the process required to identify risk and quantify adaptive capacity in order to prioritise options that can be implemented to reduce vulnerability to climate change.





Horticulture Australia

# **Climate adaptation**

/adaptation.htm ronment/climate-

formationdata.html

limate-change/

Further references and resources can be located at the end of this chapter.



## **Climate adaptation**



In general terms, the risk management framework provides an approach to:

- Identify potential risks and hazards for each element of an activity;
- Assess the possible consequences of these risks;
- Estimate the likelihood of each risk;
- Determine a level of exposure or vulnerability (determined from a combination of likelihood and consequence); and
- Identify options to manage or adapt to the risks and consequences.

There is evidence whereby such a risk framework has worked well with horticultural producers. For example, the beef risk management matrix was recently translated for use with tomato growers in the Granite Belt, QLD. This project demonstrated how successful the matrix was in identifying how vulnerable the industry was, what growers could do about it, and the limitations of adapting to climate change. Growers concluded that best practice/good agronomy was going to be their greatest defence against many potential impacts of climate change.

Similar approaches and conclusions have been derived specifically for the apple and pear industry, cherry industry (for both 'Kordia' variety and major cherry production regions), the Queensland cut flower industry and the vegetable industry. In addition, a project was undertaken to document critical temperature thresholds for a number of major horticultural commodities.

The following sections provide further detail on the components of the risk management framework, i.e., impacts, adaptive capacity and adaptation strategies.

#### Potential impacts

Some of the potential impacts on horticultural production resulting from climate change include:

- Changes in the optimum growing period or season (e.g. frost frequency);
- Shifts in optimum production locations;
- Changes in the distribution and/or abundance of pests, diseases and weeds;
- Reduced availability and increased cost of irrigation water in most locations and in some seasons;
- Increased incidence of temperature-related disorders;
- Increased risk of impacts on product guality and/or nutrition;
- Changes in the suitability, availability and adaptability of cultivars; and
- Damage from extreme events (rain, hail, wind and heat stress).

Figure 1: Generic risk management framework Source: Adapted from Stokes & Howden (2011)

It should be noted that, for horticulture, it's often discrete events or 'secondary' conditions that have the potential for greater impact. For example, it is not just increasing mean temperature, but rather the change in the number of chill hours, timing of frosts, heat events, implications for crop thresholds, duration/timing of these events etc., that have the potential for highest impact. The impacts of these discrete events are highly crop-specific.

In addition, the current policy uncertainty has implications for the sector. The carbon price and associated Carbon Farming Initiative will impact on energy and water prices, influencing irrigation/cooling costs etc.

The extent that these impacts affect horticultural products, and businesses will be further shaped by:

- The magnitude of climate change locally and how strongly each amount of change affects farm productivity;
- Growing global demand for food and productivity growth;
- Continuing uncertainty/development of both global and domestic climate change policy;
- Increasing competition for natural resources; and
- Requirements for ever more efficient and sustainable production practices.

#### Adaptive capacity

Growers are the primary agents of adaptation - they will need to assess the risk, identify options, make a decision, fund and then implement the action. Governments, researchers and industry people will need to continue to provide the information and tools, but all the rest is up to individual growers.

Adaptive capacity can be influenced by the following:

- Access to technologies that enable adaptation, such as irrigation technologies;
- Individuals 'health, age and life stage' that will influence their capacity and motivation to make decisions, achieve output, adapt to new situations and adopt new practices;
- Industry services, such as group relations, partnerships, norms and networks, which facilitate the diffusion of knowledge and innovation, provide support during structural adjustment and promote cooperative behavior; and
- Supporting policies, such as the policies adopted by government at various levels and the signals they send to producers and others in the food chain.

Successful adaptation is likely to be helped by considering the system-wide consequences of proposed adaptation measures at all social levels, at all points in the supply chain, and in relation to other concurrent challenges (such as meeting the growing demand for food and fibre). This includes reducing barriers to adaptation, while at the same time building adaptive capacity.

Further references and

the end of this chapter

resources can be located at

Horticulture Australia

# **Climate adaptation**

Further references and resources can be located at the end of this chapter



A.5

### Adaptation strategies

#### Short-term adaptation strategies - implementation of 'best practice'.

The simplest adaptation options are closely associated with existing 'best practice' and good natural resource management, and do not require farmers to make radical changes to their operations and industries in the near term. These are the use of more adaptable cultivars and a range of cultural practices that enable growers to maintain current production in current locations – i.e. adapt to the 'new' climate in the current location.

Adaptation strategies will have different challenges, opportunities, costs and risks for different crops, but could include (and are not limited to):

- Variety selection and diversification to take advantage of changes in climate conditions or manage risk of extreme events - consider rootstock selection and use of drought tolerant or 'climate ready' varieties. For many perennial crops, plan for earlier harvest times and address marketing issues such as access and timing;
- Irrigation efficiency consider identifying less water-intensive production options, developing better water delivery technologies, implementing water markets and watersharing and increase water storage capacity to better meet irrigation requirements;
- Soil and water conservation methods become even more important as climates fluctuate more and extreme events become more frequent - consider careful management of the soil moisture profile, irrigation scheduling based on soil moisture monitoring devices and landscaping to reduce erosion during heavy falls;
- Biosecurity, quarantine, monitoring and control measures can be strengthened to control the spread of pests, weeds and diseases under a warming climate - consider providing habitat for increasing beneficial insects;
- Physical/structural measures consider canopy protection using netting in fruit orchards to increase protection from heat stress, frosts and hail; evaporative cooling as a technique for reducing sunburn; and/or shade nets and some commercial kaolin-based coatings for repeling pests (shade nets have the added benefit of preventing hail damage);
- Chemical options consider chemical dormancy breakers to help counteract the lack of suitable chilling hours;
- Property design consider modifying the management of the inter-row environment, to take into account potential frost risk and the impacts of high rainfall events such as erosion and salinity, as well as integration of biocontrol techniques (e.g. host-free zones) to reduce increased use of pesticides; and
- Forecasting and planning consider tools for improving medium and long-term forecasting of seasonal conditions. Undertake risk assessment to assess sustainability in more marginal areas (e.g. chilling requirements, increased frost risk, increased quality problems).

These options can, and should, be prioritised as part of a 'no regrets' or win-win strategy for agriculture because they will provide immediate and ongoing benefits, such as productivity benefits and market access, as well as preparing the sector for climate change.

In addition, monitoring and evaluation systems are needed to continue to track changes in climate, further impacts and the effectiveness/refinement of adaptation measures. Closer collaboration between policy makers, managers, researchers, extension agencies and growers will ensure timely inclusion of climate information as it becomes available.

Further references and resources can be located at the end of this chapter.



Horticulture Australia

#### Medium term adaptation strategies - understanding and managing climate variability

In the short to medium term, existing strategies for dealing with climate variability, as above, will help to plan for, and deal with, extreme events (droughts, floods, fire, hail, etc.).

None of the existing tools used in managing climate variability have been designed specifically with any horticultural industry or application in mind. Most of these have been developed for the management of rainfall variability only and do not address temperature variability.

Currently, climate science does not adequately addresses the lead-time and season length requirements of horticultural crops, nor crop-specific data, such as thresholds for chilling hours, heat stress days, etc. For example, there are no forecast systems based on the Southern Oscillation Index (SOI) and Sea Surface temperatures (SST) which have been extensively tested for longer lead-times and shorter seasons.

The combination of long season (3 months) and short lead-time (zero), which are appropriate for other agricultural industries, is a significant constraint to the use of forecasting tools in horticulture, where a much shorter season length (several weeks to one month) and a much longer lead-time (3 to 4 months), would be much more useful. Given a sound forecast system that meets the requirements of the industry the appropriate tools can then be produced.

Existing tools for managing climate variability in agriculture include:

- POAMA (Predictive Ocean Atmosphere Model for Australia) http://poama.bom.gov.au/operational\_products.shtml - Registered Users can access additional information on (click on the 'Request Login' to gain access). The Predictive Ocean Atmosphere Model for Australia (POAMA) is a seasonal to inter-annual seasonal forecast system based on a coupled ocean/atmosphere model.
- Climate Kelpie round up of climate tools for Australian farmers http://www. climatekelpie.com.au
- Bureau of Meteorology (BOM) Seasonal Temperature Outlook http://www.bom.gov. au/climate/
- LongPaddock http://www.longpaddock.gld.gov.au/
- Madden Julian Oscillation (MJO) http://www.bom.gov.au/climate/tropnote/tropnote. shtml
- Climate Dogs http://www.depi.vic.gov.au/agriculture-and-food/farm-management/ weather-and-climate/understanding-weather-and-climate/the-climatedogs-the-fourdrivers-that-influence-victoriaas-climate
- Rainman StreamFlow version 4.3 http://www.daff.gld.gov.au/plants/field-crops-andpastures/broadacre-field-crops/cropping-efficiency/rainman
- Southern Oscillation Index (SOI) http://www.bom.gov.au/climate/glossary/soi.shtml or for latest values see http://www.longpaddock.qld.gov.au/seasonalclimateoutlook/ southernoscillationindex/index.php or seasonal outlook see http://www.longpaddock. gld.gov.au/SeasonalClimateOutlook/RainfallProbability/index.html
- Sea Surface Temperatures (SST) http://www.longpaddock.qld.gov.au/ seasonalclimateoutlook/seasurfacetemperature/index.php
- SILO climate data (available through paid licensing agreement) http://www.longpaddock.gld.gov.au/silo/

# **Climate adaptation**

Further references and resources can be located at the end of this chapter





## **Climate adaptation**

Climate Dogs are a series of short, clever and humorous animations that capture key climate science messages and package them in a way that is relevant to farmers. Through stories about four 'climate dogs', the animations discuss key drivers of Victoria's weather patterns and seasonal variability: El Niño Southern Oscillation, Indian Ocean Dipole, Subtropical Ridge and Southern Annular Mode. The four sheep dogs that round up Victoria's rainfall are ENSO, INDY, Ridgy and Sam. Some of the Climate Dogs have application to other regions of Australia as well.

Website: http://www.depi.vic.gov.au/agriculture-and-food/farm-management/weatherand-climate/understanding-weather-and-climate/the-climatedogs-the-four-drivers-thatinfluence-victoriaas-climate

The Managing Climate Variability (MCV) program is a national climate variability research initiative. To date there have been two horticulture specific projects established through collaboration with MCV:

- Australian horticulture's response to climate change and climate variability (AH06019); and
- Critical thresholds ('tipping points') and climate change impacts/adaptation in horticulture (HG08037).

See also http://www.managingclimate.gov.au

#### Long-term adaptation strategies - transformational change

As more extreme climate change is observed, the benefits obtained from each major type of adaptation are likely to plateau, as Figure 2 indicates. This means that there are likely to be limits to the effectiveness of incremental adaptations and that, at some point, a major change or transformational adaptation will be called for.

Examples of transformational adaptation could include restructuring of a business (e.g. into protected cropping or adoption of substantial off-farm work to increase income), shifting location (e.g. large financial investment into new land in a more favourable regions), and/or change to production system (e.g. crop type or production system from conventional to organic).

Common to all change, transformational adaptation to climate change can involve some significant costs (although this should not be considered alongside opportunity costs, i.e. that which emerges when action is not taken) and requires a higher level of adaptive capacity than incremental or systems-scale adaptation due to the greater risks and complexity arising from the change.

MCREASING COMPLEXITY. **BENEFIT FROM ADAPTATION** Transformation from use or distribution ch New products such a ecosystem services Climate-change ready germplasm Climate-sensitive precision agricultu Diversification and risk managemen Varieties, planting times, spacing Stubble, water, nutrient and canopy management, etc.

### **CLIMATE CHANGE**

#### A note on mitigation

It is vital that adaptation responses are integrated with mitigation as the two are intimately linked. Mitigation is a process whereby the sources of greenhouse gas emissions are reduced, and/or carbon sinks are created or enhanced. Options for mitigation in horticultural production have been addressed in Chapter 8 - Energy & greenhouse gas management.

The horticulture industry is extremely susceptible to the impacts of climate change/variability and, as a result, needs to be supporting the need for mitigation and reduction of emissions to reduce the impacts of climate change at the same time as implementing adaptation approaches (where appropriate/needed).

This involves ensuring that adaptations do not increase greenhouse gas emissions, so making the underlying cause worse - and, similarly, that mitigation options do not undermine adaptation efforts. For example, adopting management practices that improve the efficiency of nitrogen use can generate benefits for farm profitability, while also reducing greenhouse impacts and improving environmental sustainability.

Continual improvement of horticulture's response to the challenges of climate change and climate variability.

Flexibility has been the key to adaptation in horticulture to date, and is likely to continue to be an important component of adaptation strategies as climates continue to change. Growers have been able to manage climate variability reasonably well, although major improvements could be made if tools to assist with the management of climate variability, both temperature and rainfall, and extreme events were designed specifically with the needs of horticultural growers and industries in mind. This includes the development of capacity along with the provision of appropriate data and decision support tools to allow the rapid (and cheap) assessment of climate risk and the identification of possible adaptation measures, and estimates of associated costs.

It will also be important to identify those countries/regions that currently export product to Australia, which will be significantly impacted by rising temperatures along with those that will become more competitive on the Australian market because of favourable impacts as a result of further changes to the world's climate.

This will ensure both the long-term viability and sustainability of the sector, and continued availability to consumers of fresh and health-giving horticultural outputs.

Further references and resources can be located at the end of this chapter



Horticulture Australia

# **Climate adaptation**

land nange
S
ire
t

Figure 2: The potentia benefit from different levels of adaptation with increasing climate change from incremental, withinsystem responses, to system-level changes to transformational changes. Source: Stokes, & Howden, M. (2011)

> Further references and resources can be located at the end of this chapter





### **References and further reading**

ABS (2012) Completing the Picture - Environmental Accounting in Practice, Australian Bureau of Statistics, Canberra, ACT.

http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4628.0.55.001main+features30May+2012#anchor2

Amati, M. (2013) Understanding the carbon and pollution mitigation potential of Australia's urban forest (HAL Project Number: NY11002). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Brown, G. (2011) Investigating and overcoming negative effects of global warming on cherry dormancy (HAL Project Number: CY09012). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Brunt, C (2012) Cherry cultivar selection: chill hours and climate change (HAL Project Number: CY11010). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Carey, D.; White, N.; & Deuter, P. (2012) Horticulture and Climate Q - improving on our understanding of climate change, vulnerability, adaptation and risk management responses, in major horticultural industries and regions of Queensland. Horticulture and Forestry Science Department of Fisheries and Forestry Queensland.

Cobon, D.; Stone, G.; Carter, J.; Scanlan, J.; Toombs, N.; Zhang, X.; Willcocks, J.; & McKeon, G. (2009) The climate change risk management matrix for the grazing industry of northern Australia. CSIRO Publishing.

Chen, D. (2013) Greening City - Mitigate Heat Stress with Urban Vegetation (HAL Project Number: NY11013). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Corner, D. (2011) Positioning the Queensland Cut Flower Industry to Strategically Respond to the Impact of Climate Change (HAL Project Number: FL09006). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

CSIRO (2011) Climate Change: Science and Solutions for Australia, CSIRO. http://www.csiro.au/Outcomes/Climate/Climate-Change-Book.aspx

DIICCSRTE (2013) Climate Adaptation Outlook: A proposed national adaptation assessment framework. Australian Government Department of Industry, Innovation, Climate Change, Science, Research and Tertiary Education. http://www.climatechange.gov.au/climate-change/adapting-climate-change/climate-adaptation-outlook)

DIICCSRTE (2013) Climate Change Adaptation Good Practice - Case Study: South Australian Integrated Climate Change Adaptation. Australian Government Department of Industry, Innovation, Climate Change, Science, Research and Tertiary Education

Deuter, P. pers comms (2014) Senior Principal Horticulturist, Horticulture and Forestry Science, Department of Agriculture, Fisheries and Forestry (Queensland).

Deuter, P. (2011) Horticulture contribution to Managing Climate Variability Program - critical thresholds project (HAL Project Number: HG08037).

http://www.horticulture.com.au/areas\_of\_Investment/Environment/Climate/Climate\_critical\_thresholds.asp and http://www.managingclimate.gov.au/research/completed-projects/critical-thresholds-tipping-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-and-climate-points-anchange-impacts-adaptation-in-horticulture/

Deuter, P. (2008) Defining the impacts of climate change on horticulture in Australia. Chapter of the Garnaut Climate Change Review. Department of Primary Industries and Fisheries, Queensland. http://www.garnautreview.org.au/CA25734E0016A131/WebObj/01-GHorticulture/\$File/01-G%20Horticulture.pdf

Dwyer, E.; Gunner, E. & Sheperd, K. (2009) The Changing Climate: Impacts and adaptation options for South Australian primary producers. Primary Industries and Resources, SA.

Gaunaut Climate Change Review - multiple reports (2011). www.garnautreview.org.au

HAL. (2010) Climate Change: Managing Variability and Carbon. Horticulture Australia Ltd. http://ausveg.businesscatalyst.com/rnd/fact%20sheets/Climate%20Change%20and%20Managing%20Carbon.pdf

Howe, T. (2012) Carbon footprinting and labelling - understanding the language, trends and tools (HAL Project Number: HG10045). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

IPCC (2013) Summary for Policymakers. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. http://www.ipcc.ch

McKenzie, R. (2011) First steps in a response to climate change and climate variability: desktop study (HAL Project Number: AP09019). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Pearson, L. and Langridge, J. (2008) 'Climate change vulnerability assessment: Review of agricultural productivity' CSIRO Climate Adaptation Flagship Working paper No.1. Web-ref http://www.csiro.au/resources/CAF-Working-Papers

Putland, D. & Deuter, P. (2011) The effects of high temperatures on vegetable production and the rapid assessment of climate risk in agriculture. Proceedings of the 2011 APEC Workshop on Collaboration on the Promotion of Indigenous Vegetables for Coping with Climate Change and Food Security, pg 45-62. APEC Agricultural Technical Cooperation Working Group, 21-24 November 2011, Chinese Taipei.

Rickards, L. & Howden, M. (2012) Transformational adaptation: agriculture and climate change. CSIRO Publishing. Crop and Pasture Science 63(3) 240-250. http://www.publish.csiro.au/paper/CP11172.htm

Roche, M. (2013) Adaptation of warm-season turf grasses for tropical Australia (HAL Project Number: TU09001). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Rogers, G. (2013) Understanding and managing impacts of climate change and variability on vegetable industry productivity and profits (HAL Project Number: VG12041). Final report published by Horticulture Australia Ltd, Sydney, NSW. www.horticulture.com.au

Stokes, CJ and Howden, SM. (2011) Adapting agriculture to climate change, Chapter 7 of Climate Change: Science and Solutions for Australia, CSIRO. http://www.publish.csiro.au/?act=view\_file&file\_id=CSIRO\_CC\_Chapter%207.pdf

Stokes CJ and Howden SM (eds) (2010) Adapting Agriculture to Climate Change: Preparing Australian Agriculture, Forestry and Fisheries.



Horticulture Australia

# **Climate adaptation**



### Check environmental laws and regulations

Various laws and regulation will apply to your property. These laws could be local, state or Commonwealth.

As laws and regulations can vary considerably between states and regions, you should contact your State/Local conservation department/council and find out what laws apply to your property.

Refer to the following websites to access electronic copies of the Principal Acts and Statutory Rules in operation and advice on relevant laws.

#### Australasian Legal Information Institute www.austlii.edu.au

Australian Commonwealth, State, Territory, and Local Government Environmental Information – useful link via Environmental Defenders Office http://www.edo.org.au/ links/govlinks.html

**ComLaw** - has the most complete and up-to-date collection of Commonwealth legislation and includes notices from the Commonwealth Government Notices Gazette from 1 October 2012 www.comlaw.gov.au

Environmental Defenders Office - a network of independent community environmental law centres www.edo.org.au

#### **Environment Protection Authorities**

- ACT www.environment.act.gov.au
- NSW www.environment.nsw.gov.au or www.planning.nsw.gov.au
- NT www.ntepa.nt.gov.au
- QLD www.nprsr.qld.gov.au
- SA www.epa.sa.gov.au
- TAS www.epa.tas.gov.au VIC
- www.epa.vic.gov.au WA
- www.epa.wa.gov.au

National Farmers Federation – NFF has three Liaison Officers on secondment from Australian Government departments to provide services direct to the agriculture sector in the areas of environmental law, biosecurity, and immigration and labour http://www.nff.org.au/policy/liaison-officers.html

#### National Library of Australia - Australian Law websites

For assistance with international environmental laws visit http://edo.org.au/links/iel.html

For instance, The Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) is the Australian Government's environmental legislation. It covers environmental assessment and approvals, protects significant biodiversity and integrates the management of important natural and cultural places. The EPBC Act protects matters that are of national environmental significance. Those most relevant to farmers are:

- Nationally threatened and migratory species;
- Nationally threatened ecological communities;
- Wetlands of international importance;
- World and national heritage properties; and
- The Great Barrier Reef.

New farm activities, such as land clearing, may require approval from the Federal Environment Minister under national environment law.



Relevant national legislation/regulation:

- Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) http://www.environment.gov.au/topics/about-us/legislation/environmentprotection-and-biodiversity-conservation-act-1999/about-epbc
- Hazardous Waste Act 1989 http://www.environment.gov.au/topics/environmentprotection/hazardous-waste/about-hazardous-waste-act
- Ozone Protection and Synthetic Greenhouse Gas Management Act 1989 http://www.environment.gov.au/topics/environment-protection/ozone-andsynthetic-greenhouse-gases/legislation
- Water Act 2007 [e.g. Basin Salinity Management Strategy is implemented under Schedule B of the Murray-Darling Basin Agreement (Schedule 1 of the Water Act 2007 Cwlth)]. http://www.environment.gov.au/topics/water/australiangovernment-water-leadership/water-legislation
- Water Quality National Water Quality Management Strategy (NWQMS) (2000) Australian and New Zealand guidelines for fresh and marine water quality: Volume 3 - primary industries http://www.environment.gov.au/resource/australian-andnew-zealand-guidelines-fresh-and-marine-water-quality-volume-1-guidelines (note that the guidelines are currently under review and next edition due to be released in June 2014).

### **Regional/local information**

It is difficult to be specific on the local legislative requirements for horticultural enterprises. However, information regarding the local applicability of legislative requirements can often be sourced through the relevant catchment management authority.

To find your relevant catchment management authority visit the national Caring for Our Country website: http://www.nrm.gov.au/about/nrm/regions/index.html

Alternatively, see below for your relevant state:

- http://www.actnrmcouncil.org.au ACT
- NSW http://www.nrm.gov.au/about/nrm/regions/nsw.html
- NT http://www.territorynrm.org.au
- QLD http://www.nrm.gov.au/about/nrm/regions/qld.html
- SA http://www.nrm.gov.au/about/nrm/regions/sa.html
- TAS http://www.nrm.gov.au/about/nrm/regions/tas.html
- VIC http://www.nrm.gov.au/about/nrm/regions/vic.html
- WA http://www.nrm.gov.au/about/nrm/regions/wa.html

### By specific chapter/topic

### **SOIL & GROUNDWATER**

#### **EPHC/NEPC** Assessment of Site Contamination National Environment Protection Measure - NEPM 2001

National Environment Protection Measures (NEPMs) outline agreed national objectives for protecting or managing particular aspects of the environment. The NEPM for assessment of contaminated sites contains two schedules, Schedule A and Schedule B. Schedule A describes the site assessment process that indicates which general guidelines within the NEPM are applied to each level of site investigation. Schedule B identifies 10 individual guidelines for the assessment of contaminated sites, including soil and groundwater. The objective of the site assessment is to determine whether

site contamination poses an actual or potential risk to human health and/or the environment of sufficient magnitude to warrant remediation. An online version of this document can be found at the following address: http://www.scew.gov.au/system/ files/resources/93ae0e77-e697-e494-656f-afaaf9fb4277/files/schedule-b1-guidelineinvestigation-levels-soil-and-groundwater-sep10.pdf

#### WATER (Including Sediment)

#### Australian and New Zealand Guidelines for Fresh and Marine Water Quality 2000 http://www.deh.gov.au/water/quality/nwqms/index.html#quality

These guidelines are designed to help users assess whether the water quality of a water resource is good enough to allow it to be used for humans, food production or aquatic ecosystems (these uses are termed environmental values). If the water quality does not meet the water quality guidelines, the water may not be safe for those environmental values and management action could be triggered to either more accurately determine whether the water is safe for a particular use or, if not, to remedy the problem. These Guidelines include chemicals within sediments, and water considered for primary industry, human health (recreational and aesthetic) and water within aquatic ecosystems. They are not meant to be applied directly to recycled water quality, contaminant levels in discharges from industry, mixing zones, or stormwater quality, unless stormwater systems are regarded as having conservation value. These Guidelines should not be used as mandatory standards.

#### NHMRC Australian Drinking Water Guidelines 2001

https://www.nhmrc.gov.au/guidelines/publications/eh52

The National Health and Medical Research Council (NHMRC) Australian Drinking Water Guidelines (ADWG) are the primary reference on drinking water quality in Australia and provide necessary guidance for the provision of a safe and high quality drinking water supply that protects public health and meets the needs and expectations of consumers. They are concerned with the safety of water from a health point of view and with its aesthetic quality. The guidelines are applicable to any water intended for drinking (except bottled or packaged water, and ice) irrespective of its source (municipal supplies, rainwater tanks, bores, point-of-use treatment devices etc.) or where it is used (the home, restaurants, camping areas, shops etc.). Exceeding a guideline value should be a signal to investigate the cause and, if appropriate, to take remedial action. If the characteristic is health-related, the relevant health authority should be consulted.

Note: Other documents exist outlining guidelines and standards for chemicals in the environment, including a wide range of documents produced by international agencies. It is recommended that guidelines presented above should be consulted first when assessing chemical concentrations in the Australian environment.

### BIOTA

#### Australian New Zealand Food Standards Code (ANZFSC) http://www.comlaw.gov.au/Details/F2013C00651/Download

The Australian New Zealand Food Standards Code (ANZFSC) is a collection of individual food standards, and deals with standards that apply to all foods, standards affecting particular classes of foods and food hygiene issues in Australia. ANZFSC Standards for contaminants and natural toxicants sets out the maximum levels (MLs) of specified metal and non-metal contaminants and natural toxicants in nominated foods. Unless expressly provided elsewhere in the ANZFSC, the provisions of the Code apply to food products sold or prepared for sale in Australia and/or New Zealand, and/or imported into Australia and/or New Zealand.

Horticulture Australia

## Legislation & regulation resources



#### National Residue Survey

http://www.daff.gov.au/agriculture-food/nrs http://www.apvma.gov.au

The National Residue Survey (NRS) was established for the purpose of monitoring and reporting the level of contaminants in food, inputs to production and/or the environment. Monitoring for residues, as undertaken by the NRS, helps audit the use of currently registered agricultural and veterinary chemicals in Australia. At present the chemicals that may be detected include: antimicrobial agents (disease control); anthelmintics (parasite control); hormonal growth promotants; fungicides; insecticides and acaricides; fumigants (e.g. grain protectants); and herbicides used to control weeks in crops. The NRS monitors residue levels against maximum residue limits (MRLs) listed in the ANZFSC. However, where the Australian Pesticides & Veterinary Medicines Authority (APVMA) has established an MRL that has not yet been adopted into the ANZFSC, this fact is taken into consideration by the NRS when interpreting the significance of any results that do not have limits expressed by the ANZFSC.

### AIR

#### National Environment Protection Measure (Ambient Air NEPM and Air Toxics NEPM) http://www.comlaw.gov.au/Details/C2004H03935/Download

National Environment Protection Measures (NEPMs) outline agreed national objectives for protecting or managing particular aspects of the environment. The Ambient Air Quality NEPM sets uniform standards for a set of agreed criteria air pollutants in ambient air (ambient air does not include indoor air) that affect human health. The Air Toxics NEPM establishes a national framework for monitoring and reporting ambient air toxics. The Measure is primarily concerned with the collection of data on ambient levels of formaldehyde, toluene, xylene, benzene and polycyclic aromatic hydrocarbons at locations where elevated levels are expected to occur and there is a likelihood that significant population exposure could occur.

#### Methods for the sampling and analysis of air pollutants in New South Wales http://www.epa.nsw.gov.au/air/faqamsampling.htm

This document lists the methods to be used and provides guidance for the modelling and assessment of air pollutants from stationary sources in New South Wales for statutory purposes. The document covers:

- Impact assessment criteria for criteria pollutants, hydrogen fluoride, deposited dust and total suspended particulate (TSP) matter;
- Ground-level concentration (GLC) criteria for individual odorous and toxic air pollutants;
- GLC criteria for hydrogen sulphide;
- Odour performance criteria for complex mixtures of odours;
- Impact assessment methodology based on dispersion modelling;
- The procedure for developing site-specific emission limits.

This document contains guidance for estimating the air quality impacts of a much larger set of potential air pollutants than the Air NEPMs.

Note: This is a State government document and has not been through the national approval process and is not applicable in all jurisdictions.



## Crop production – process flow

This diagram shows a flow of steps that may occur while growing crops in the ground, the inputs that could introduce an environmental impact and the areas in which processes may have an impact. In practice the steps do not follow a set order after planting and some steps are not required for all crops. For hydroponics, the nutrient solution and the root support medium are extra inputs. For nursery container production, pots and growing media are extra inputs.



**Process steps and inputs** 

HAL

**C.1** 

## Field packing – process flow

This diagram shows a flow of steps that may occur during field packing of fresh produce and the inputs for each step that could introduce an environmental impact. Some steps are not required for all crops. For example, some crops are not washed before packing and others are not pre-cooled before dispatch.



## Shed packing – process flow

This diagram shows a flow of steps that may occur during shed packing of fresh produce and the inputs for each step that could introduce a food safety hazard. The order and presence of steps varies with crops.



Activity can have impact on
Land and soil, water, biodiversity
Air, waste, energy
Land and soil, water, air, waste, energy
Land and soil, water, biodiversity, waste
Land and soil, water, biodiversity, waste
Land and soil, water, biodiversity, waste
Land and soil, water, biodiversity
Biodiversity, air, waste, energy
Land and soil, water, biodiversity, air, waste, energy
Air, waste
Air, waste
Land and soil, water, biodiversity, air, waste, energy
Land and soil, water, biodiversity, air, waste, energy

HAL

## Nursery container production - process flow







## Growing the crop – process flow

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Select site	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Removal or destruction of rare/ endangered flora or fauna	Loss of biodiversity	Reduction of wildlife corridors
	Catchment management plan or other statutory requirements not considered/ adhered to	Legislative penalties	Compromises catchment aims and objectives
	Horticultural production (monoculture)		Loss of biodiversity, reduction of wildlife corridors
	Unaware of existence of rare/endangered flora or fauna		Loss of biodiversity, reduction of wildlife corridors
	Detrimental impact on riparian zones, wetlands or other sensitive vegetation types	Soil erosion Reduction of water quality	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways Loss of biodiversity, reduction of wildlife corridors
	Detrimental impact on water resource/ environmental flows	Reduction of water availability and quality	Sedimentation of rivers/waterways Loss of aquatic habitat
	Inappropriate land use or clearing	Salinity Reduction of arable land Sodicity Reduction of arable land Soil erosion Waterlogging	Reduction of arable land Spread of saline land and water Effects on buildings Effects on native vegetation Reduction of arable land Sedimentation of rivers/waterways Nutrients and agricultural chemicals Effects on native vegetation
Prepare site	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off
	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Nutrient leaching (inappropriate rate or type of fertiliser)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Atmospheric pollution by greenhouse gases Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Pollution – inappropriate application of fertiliser to non- cropping areas (throwing into waterways, dams, remnant vegetation)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal bloom Adverse impact on flora and fauna

HAL

# Environmental impact identification table - Growing the crop

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Prepare site (cont)	Pollution – inappropriate storage of fertilisers	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal bloom Adverse impact on flora and fauna Discomfort or Inconveneince of local residents (odour)
	Pollution – inappropriate storage of agricultural chemicals	Adverse impact on flora and fauna Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate disposal of agricultural chemicals, surplus agricultural chemicals and rinsates	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate mixing site for agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Adverse impact on flora and fauna
Prepare nursery potting media	Pollution – leaching of nutrients and other soluble pollutants	Contamination of surface and/or groundwater	Contamination of surface and/or groundwater
	Pollution- litter and other floating contaminants	Contamination of surface and/or groundwater	Contamination of surface and/or groundwater
	Pollution – dust, odour	Inconvenience to staff	Inconvenience to local residents
Planting	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off
	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Nutrient leaching (inappropriate rate or type of fertiliser)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Atmospheric pollution by greenhouse gases Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Pollution - inappropriate application of fertiliser to non cropping areas (throwing into waterways, dams, remnant vegetation)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna

# Environmental impact identification table - Growing the crop

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Planting (cont)	Inefficient use of water resource	Rising water table Increasing soil salinity from rising water table Soil erosion (see above) Nutrient leaching (see above) Waterlogging Contamination of land and water from run-off of contaminated water (chemicals/ nutrients) Adverse impact on flora and fauna (environmental flows, reduction of habitat in dams/lakes) Depletion of water table (water taken from bores) Reduction of water quality	Rising water table Increasing soil salinity from rising water table Soil erosion (see above) Nutrient leaching (see above) Contamination of land and water from run-off of contaminated water (chemicals/ nutrients) Loss of aquatic habitat, adverse impact on flora and fauna (environmental flows, reduction of habitat in dams/lakes) Depletion of water table (water taken from bores) Reduction of water quality
	Energy inefficiency – irrigation equipment/timing of irrigation	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases
	Pollution – leaching of nutrients	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Salinity/sodicity – inappropriate rate of irrigation or quality of irrigation water	Reduction of arable land Soil erosion Waterlogging	Reduction of arable land Spread of saline land and water Effects on buildings
	Selecting environmental weeds for nursery propagation and sale		Effects on native flora and fauna Loss of biodiversity
Irrigating	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off
	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Inconvenience to local residents Adverse impact on flora and fauna
	Inefficient use of water resource	Rising water table Increasing soil salinity from rising water table Soil erosion (see above) Nutrient leaching (see above) Waterlogging Contamination of land and water from run-off of contaminated water (chemicals/ nutrients) Adverse impact on flora and fauna (environmental flows, reduction of habitat in dams/lakes) Depletion of water table (water taken from bores) or water from rivers and creeks Reduction of water quality	Rising water table Increasing soil salinity from rising water table Soil erosion (see above) Nutrient leaching (see above) Contamination of land and water from run-off of contaminated water (chemicals/ nutrients) Loss of aquatic habitat, adverse impact on flora and fauna (environmental flows, reduction of habitat in dams/lakes) Depletion of water table (water taken from bores) or water from rivers and creeks (environmental flows) Reduction of water quality
	Energy inefficiency – irrigation equipment/timing of irrigation	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases

HAL

# Environmental impact identification table - Growing the crop

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Irrigating (cont)	Pollution – leaching of nutrients	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Salinity / sodicity – inappropriate rate of irrigation or quality of irrigation water	Reduction of arable land Soil erosion Waterlogging	Reduction of arable land Spread of saline land and water Effects on buildings Effects on flora and fauna
Crop nutrition	Pollution – inappropriate storage of fertilisers	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and alga blooms Adverse impact on flora and fauna
	Pollution – inappropriate application of fertiliser to non cropping areas (throwing into waterways, dams, remnant vegetation)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
	Nutrient leaching (inappropriate rate or type of fertiliser)	Contamination of surface water and/or groundwater Eutrophication of surface waters and algal blooms Soil contamination Soil acidification Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Atmospheric pollution by greenhouse gases Eutrophication of surface waters and algal blooms Adverse impact on flora and fauna
Pest/ disease / weed management	Pollution – inappropriate storage of agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate disposal of used chemical containers	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate disposal of used chemical containers	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Impacts on landfill capacity
	Pollution – off target application of agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna Disruption of Integrated Pest Management strategies Adverse impact on other crops	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
Pruning / training	Escapes/volunteer growth	Adverse impact on flora and fauna, invasion of native habitat Requirement for control measures to be implemented (chemical, mechanical etc)	Adverse impact on flora and fauna, invasion of native habitat Requirement for control measures to be implemented (chemical, mechanical etc)
Crop / growth regulation	Pollution – inappropriate storage of agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination	Contamination of surface water and/or groundwater Adverse impact on flora and fauna

# Environmental impact identification table - Growing the crop

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Crop / growth regulation (cont)	Pollution – inappropriate disposal of used chemical containers	Adverse impact on flora and fauna Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna Impacts on landfill capacity
	Pollution – inappropriate disposal of agricultural chemicals, surplus agricultural chemicals and rinsates	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – off target application of agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
Weather	Trapping fauna (in netting)	Adverse impact on flora and fauna	Adverse impact on flora and fauna
/ vermin protection	Noise generation (scarers)	Disturbance to other fauna	Discomfort or inconvenience of local residents
Harvest	Change to visual landscape Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Disturbance to other fauna Contamination of surface water and/or groundwater inconvenience of local residents
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources Traffic – noise, dust, fumes Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Inconvenience to local residents Adverse impact on flora and fauna
	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off
	Escapes/volunteer growth	Adverse impact on flora and fauna, invasion of native habitat Requirement for control measures to be implemented (chemical, mechanical etc)	Adverse impact on flora and fauna, invasion of native habitat Requirement for control measures to be implemented (chemical, mechanical etc)
Transport to packing shed	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Inconvenience to local residents Adverse impact on flora and fauna
	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
Post-harvest site management	Use of livestock to remove crop residues (scavengers	Soil compaction Degradation of soil quality Increased run-off Escapes/volunteer growth	Sedimentation of rivers/waterways increased run-off Escapes/volunteer growth
	Weed transmission by scavengers	Requirement for control measures to be implemented (chemical, mechanical etc) Inefficient and polluting equipment	Requirement for control measures to be implemented (chemical, mechanical etc)

Based on 'Growing the crop' process flow – Guidelines for on-farm food safety for farm produce process flow (DAFF, 2004).

HAL

HAL

# Environmental impact identification table - Field packing

## Field packing

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Washing	Soil erosion – inappropriate disposal of wash water	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/ waterways
	Pollution – inappropriate disposal of wash water, particularly if containing sanitisers/agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Salinity / sodicity – inappropriate rate of irrigation or quality of irrigation water	Reduction of arable land Soil erosion Waterlogging	Reduction of arable land Spread of saline land and water Effects on buildings Effects on flora and fauna
Grading and packing	Pollution – inappropriate disposal of waste produce	Contamination of land, surface water and/or groundwater Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Discomfort or inconvenience of local residents
	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Pollution – inappropriate disposal of unsatisfactory packaging		Rubbish in waterways Inconvenience to local residents Atmospheric pollution by greenhouse gases
	Noise	Adverse impact on/disturbance of fauna	Adverse impact on fauna Inconvenience to local residents
Palletising	Pollution – inappropriate disposal of unsatisfactory pallet wrap		Rubbish in waterways Inconvenience to local residents Atmospheric pollution by greenhouse gases
Transport	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Inconvenience to local residents Adverse impact on flora and fauna
	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/ waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality increased run-off	Sedimentation of rivers/waterways increased run-off
Precooling / storage	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents Adverse impact on fauna
	Pollution – use of electricity and coolants	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases (CFC refrigerants)
	Pollution – coolant leakage	Contamination of land	Contamination of surface water and/or groundwater

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Dispatch	Oil or fuel spillage	Contamination of land	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna Movement of contaminants (chemical, weed seed) by machinery	Inconvenience to local residents Adverse impact on flora and fauna
	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/ waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off

Based on 'Field Packing' process flow – Guidelines for on-farm food safety for farm produce process flow (DAFF, 2004).

# Environmental impact identification table - Field packing



## Shed packing

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Receive/ unload	Soil erosion – inappropriate disposal of dump tank/wash water	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Pollution – inappropriate disposal of dump tank/wash water, particularly if containing sanitisers/ agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate disposal of waste produce	Contamination of surface water and/or groundwater Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Discomfort or inconvenience of local residents
Curing	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases
	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents Adverse impact on flora and fauna
Precooling	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents Adverse impact on flora and fauna
	Pollution – use of electricity and coolants	Atmospheric pollution by greenhouse gases Wasting resources	Inconvenience to local residents Adverse impact on flora and fauna
	Pollution – coolant leakage	Contamination of land Contamination of surface water and/or groundwater	Atmospheric pollution by greenhouse gases (CFC refrigerants)
Destemming / dehanding / defuzzing / trimming	Pollution – inappropriate disposal of dump tank/wash water, particularly if containing surfactant/ sanitisers/ agricultural chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwaterAdverse impact on flora and fauna
J	Pollution – inappropriate disposal of waste produce	Contamination of surface water and/or groundwater Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Discomfort or inconvenience of local residents
Washing	Soil erosion – inappropriate disposal of wash water	Loss of topsoil and nutrients	Sedimentation of rivers/waterways
	Pollution – inappropriate disposal of wash water, particularly if containing sanitisers/agricultural chemicals and/ or organic matter	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater with chemicals and organic matter Adverse impact on flora and fauna
	Pollution – coolant leakage	Contamination of land	Contamination of surface water and/or groundwater
Chemical treatment	Pollution – inappropriate storage of post harvest chemicals	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
	Pollution – inappropriate disposal of used chemical containers	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Impacts on landfill capacity
	Pollution – inappropriate disposal of post harvest chemicals, surplus post harvest chemicals and rinsates	Contamination of surface water and/or groundwater Soil contamination Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna
Waxing	Pollution – inappropriate disposal of waxes	Contamination of surface water and/or groundwater Soil contamination	

Step	Hazard (the thing that happened, the thing you do)	Environmental impact – on farm	Environmental impact – off farm
Drying / polishing	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases
	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents
Grading and packing	Pollution – inappropriate disposal of waste produce	Contamination of surface water and/or groundwater Adverse impact on flora and fauna	Contamination of surface water and/or groundwater Adverse impact on flora and fauna Discomfort or inconvenience of local residents
	Pollution – inappropriate disposal of unsatisfactory packaging		Rubbish in waterways Inconvenience to local residents Atmospheric pollution by greenhouse gases
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases
	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents
	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contaminataion of surface water and/or groundwater
Top-icing	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases
Palletising	Pollution – inappropriate disposal of unsatisfactory pallet/product wrap		Rubbish in waterways Inconvenience to local residents Atmospheric pollution by greenhouse gases
	Pollution – use of electricity and coolants	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases (CFC refrigerants
	Pollution - coolant leakage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
Dispatch	Oil or fuel spillage	Contamination of land Contamination of surface water and/or groundwater	Contamination of surface water and/or groundwater
	Inefficient or polluting equipment	Atmospheric pollution by greenhouse gases Contamination of land Contamination of surface water and/or groundwater Wasting resources	Atmospheric pollution by greenhouse gases Contamination of surface water and/or groundwater
	Traffic – noise, dust, fumes	Adverse impact on flora and fauna	Inconvenience to local residents Adverse impact on flora and fauna
	Soil erosion	Loss of topsoil and nutrients	Sedimentation of rivers/waterways Nutrients and agricultural chemicals entering rivers/waterways
	Detrimental impact on soil structure, soil compaction	Degradation of soil quality Increased run-off	Sedimentation of rivers/waterways Increased run-off
	Pollution – use of electricity	Atmospheric pollution by greenhouse gases Wasting resources	Atmospheric pollution by greenhouse gases Inconvenience to local residents (lighting at night)
	Noise	Adverse impact on/disturbance of fauna	Inconvenience to local residents Adverse impact on fauna

Based on 'Shed Packing' process flow. Guidelines for on-farm food safety for farm produce process flow (DAFF, 2004)

HAL

# Environmental impact identification table - Shed packing

HAL



#### **Further resources**

(web links accurate as at 11 February 2014)

#### THIRD PARTY SPECIFICATIONS PROGRAMS

#### Freshcare Code of Practice Environmental

www.freshcare.com.au

The Freshcare Code of Practice Environmental (2nd Edition 2011) is an industry owned standard, describing the practices required on farm to provide assurance that produce has been grown and packed with care for the environment.

Freshcare endorsed certification bodies include (as of 30 September 2013):

Australian Certified Organic www.aco.net.au

Aus-Qual www.ausqual.com.au

NCS International Pty Ltd www.ncsifood.com

SCI-Qual International www.scoqual.com.au

SGS Systems & Services Certification www.au.sgs.com

Silliker Global Certification Services Pty Ltd www.certification.silliker.com

#### Globalgap

www.globalgap.org

The GLOBALG.A.P. Fruit & Vegetables Standard covers all stages of production, from pre-harvest activities such as soil management and plant protection product application to post-harvest produce handling, packing and storing. Certified in Australia through approved bodies (as of 30 September 2013):

Aus-Qual www.ausqual.com.au

NCS International Pty Ltd www.ncsifood.com

SAI Global Certification Services Pty Ltd www.saiglobal.com

SCI-Qual International www.scoqual.com.au

SGS Systems & Services Certification www.au.sgs.com

#### LEAF

http://www.leafuk.org/leaf/home.eb

LEAF (Linking Environment And Farming) Marque is a globally recognized, independently certified assurance scheme. It is based on Integrated Farm Management principles of sustainable farming. LEAF is a UK based organization working with farmers and organisations from around the world to promote sustainable food and farming. Certified in Australia through approved bodies (as of 30 September 2013 - SAI Global and Asure Quality Ltd).

Asure Quality Ltd http://www.asurequality.com

SAI Global Certification Services Pty Ltd www.saiglobal.com

#### OTHER LAND SECTOR ENVIRONMENTAL SYSTEMS

#### Certified Land Management

http://almg.org.au/index.htm

The Australian Land Management Group's Certified Land Management (CLM) system helps land managers, industry organisations and natural resource management agencies improve environmental and animal welfare management and provides an independent verification of performance. Supported by foundation partner, Elders, CLM operates across industries on a whole-of-farm and landscape-linked basis. Based on myEMS software system.

#### **Cotton Best Management Practice (BMP)**

https://www.mybmp.com.au/home.aspx and http://www.bmpcotton.com.au

BMP is the Australian cotton industry's guide for growing cotton in harmony with the natural environment. It is a grower driven, voluntary farm management system that provides self-assessment mechanisms, practical tools and auditing processes to ensure that cotton is produced with best practice across a range of focus areas.

#### Dairying for Tomorrow

http://www.dairyingfortomorrow.com.au

The dairy environmental self-assessment tool, DairySAT, is the entry point. Dairy farmers also have access to two other tools - the Dairy Greenhouse Abatement Strategies calculator and the Farm Nutrient Loss Index.

## Freshcare Environmental – Viticulture (2nd Edition, June 2011) and Winery (2nd Edition, 2013)

http://www.wfa.org.au/entwineaustralia/

The Freshcare Environmental Viticulture and Environmental Winery Codes of Practice are industry-owned codes that provide businesses with a tailor-made environmental management system specific to the processes involved with wine grape production and wine making. Links to 'Entwine Australia' the voluntary national environmental assurance program available to all Australian wine companies and grape growers.



#### HORTICULTURE INDUSTRY

#### Australian Organic

Supports Australian organic producers and maintains the Australian Certified Organic Standard 2013. http://austorganic.com

Australian Society of Horticultural Science Promotion and enhancement of Australian science and the horticulture industry. http://www.aushs.org.au

#### Growcom

Provides representation, leadership and solutions for horticultural businesses in Queensland. Hosts and implements the Farm Management System/Farmcare training package. Modules available include water use efficiency, soil nutrient management, water quality, energy efficiency and sediment runoff. www.growcom.com.au

Horticulture Australia Limited Horticulture's research, development and extension corporation. Links to industry contacts, research priorities and research reports: www.horticulture.com.au

#### Horticulture Industry Network

An initiative aimed at strengthening working partnerships between the Department of Environment and Primary Industries Victoria and the horticulture industry to deliver better services to farmers. http://www.hin.com.au

http://www.hin.com.au/resources/hort-service-finder - The Hort Service Finder App (iOS) was build as a free service to help Australian horticultural growers to find horticultural service providers close to them.

#### Northern Territory Horticulture Association

The NT Horticultural Association along with the NT Agricultural Association are undergoing a process of amalgamation to form the: Northern Territory Farmers Association. http://www.ntha.com.au

#### **Organic Federation of Australia**

Peak body for Australia's organic and biodynamic producers. www.ofa.org.au

#### Protected Cropping Australia

Peak industry body representing commercial Australian hydroponic and greenhouse growers. http://www.protectedcroppingaustralia.com

#### FARMER ORGANISATIONS

National Farmers Federation www.nff.org.au Northern Territory Farmers Federation www.ntfarmers.com NSW Farmers Association www.nswfarmers.org.au Queensland Farmers Federation www.qff.org.au South Australia Farmers Federation www.saff.org.au Tasmania Farmers and Graziers Association www.tfga.org.au Victorian Farmers Federation www.vff.org.au Western Australia Farmers Federation www.waff.org.au

#### OTHER RELATED ORGANISATIONS AND RESOURCES

#### ΔRC

News, events and information for horticulture sector. http://www.abc.net.au/news/rural/horticulture/

#### **Agricultural Links**

Integrates international agriculture and livestock websites help you find agritourism, poultry, garden, production, farmer, goat, fodder crops, horticultural crop and more. http://www.agriculturallink.com

#### Australian Food and Grocery Council

AFGC is the leading national organisation representing Australia's food, drink and grocery manufacturing industry. Links to sustainability commitments, reports and case studies: http://www.afgc.org.au/sustainability.html

#### Australian Farm Institute

Independent farm policy research institute. Link to FarmGAS scenario tool: http://www.farminstitute.org.au

#### Australian Government NRM knowledge repository

This repository is designed as an open access digital archive for publicly funded information derived from Australian Government investments in natural resource management activities. http://nrmonline.nrm.gov.au

Australian River Restoration Centre

A not-for-profit organisation that brings together people to work on river and knowledge sharing projects. http://arrc.com.au

#### Australian Society of Agronomy

The Development of Ecological Performance Indicators for Sustainable Systems. www.regional.org.au/au/asa/2001/4/c/kemp.htm

Bureau of Meteorology Australia's national weather, climate and water agency. www.bom.gov.au

#### CSIRO Plant Industry

Plant Industry promotes profitable and sustainable agrifood, fibre and horticultural industries, developing new plant products and improving how plants use natural resources through world-leading research. http://www.csiro.au/pi

#### Department of Agriculture, Forestry and Fisheries

Develops and implements policies and programs for Australia's agricultural, fisheries, food and forestry industries. Responsible for National Food Plan. www.daff.gov.au

#### Department of Environment

Portal for Departments of Sustainability, Environment, Water, Population and Communities. http://www.environment.gov.au For State and Territory links visit http://www.environment.gov.au/node/22763

Farmsafe Australia Farmsafe Australia Inc is an association of national agencies that share a common interest in Australian farm

#### safety. http://farmsafe.org.au

Federal Grants Link

Portal for information on available grants. For portal on agricultural grants: http://grantslink.gov.au/grants/grants.aspx?query=x:%22Agriculture%22&link-search=true

Get Farming Australia Free farming and agribusiness referral service. http://www.getfarming.com.au

#### Greening Australia

Community owned organization for Australian landscapes. This website provides access to resources for vegetation management. www.greeningaustralia.org.au/resources/resources

Landcare Australia

our country. www.landcareonline.com.au

Ollie's Island Ollie's Island is an interactive 'edu-tainment' program for middle school students that explores sustainable production and consumption.

http://www.olliesworld.com/island/aus/index.htm

QFVG (1998) Farmcare

## Code of Practice for Sustainable Fruit and Vegetable Production in Queensland

#### State of the Environment reports

State of the Environment reporting occurs at both the national and state/territory level. Some regional-scale reporting also occurs in many areas throughout Australia. Latest report was completed in 2011. http://www.environment.gov.au/topics/science-and-research/state-environment-reporting

#### INTERNATIONAL RESOURCES

#### International Organization for Standardisation (ISO)

http://www.iso.org/iso/home.html%20

The ISO 14000 family of environmental management standards that can be implemented in any type of organization in either public or private sectors. ISO14000 series should cooperate closely with the ISO 9000 quality management and quality assurance standards. ISO 14001 is a widely used and understood international standard for environmental management practices, providing a rigorous system of accountabilities and an annual auditing schedule.

Food and Agriculture Organisation (FAO) www.fao.org Sustainable Agriculture Initiative http://www.saiplatform.org The Earth Charter Initiative http://www.earthcharterinaction.org Global Reporting Initiative https://www.globalreporting.org/Pages/default.aspx

Horticulture Australia

## Further resources

A national network of thousands of locally-based community groups who care for the natural resources of





# **Guidelines for Environmental Assurance** in Australian Horticulture - Review Checklist

### Developed by:

# HORTICULTURE FOR TOMORROW

Managed by Horticulture Australia Limited

## Second Edition - 2014

Disclaimer

This guide has been produced by Horticulture Australia Ltd (HAL) to provide general information on environmental management in horticulture. The information has been provided in good faith, and the views expressed do not necessarily represent the views of HAL. HAL does not accept responsibility for the accuracy of the information provided and no responsibility or liability will be accepted for any use or reliance on the material contained in the document.

Laws governing environmental protection and management by horticultural producers can be complex, and may be include numerous Federal, State and Local Acts and Regulations. In addition, changes to these laws may be made from time to time. The user of this guide should seek expert advice from an appropriate professional or the relevant government agency to ensure the precise effect of current laws is fully understood before implementing any course of action referred to in this guide.

© 2014 – Horticulture Australia Limited. Parts of this document may be reproduced provided they are not sold for profit and the copyright owner is clearly identified on any reproduction.







## **Review checklist**

This review checklist identifies the range of environmental issues that should be addressed on-farm in horticulture. The checklist covers 20 topics divided into eight sections which match the Management Chapters of the Guidelines folder.

The checklist can be used to supplement existing audit checklists used by certification bodies or for internal audits carried out by individual businesses.

As with any generic checklist, you may not consider some items to be relevant and they can be excluded.

By working through the checklist you will get an idea of your priority areas and you can then read the relevant topic in the Guidelines in more detail.

The Checklist is divided into tables covering a range of topics. Select a topic and turn to the matching topic in the main working section of the Guidelines and look for the Risk Assessment diagram. By working though the Risk Assessment you can quickly determine if that particular topic of the guidelines is significant for your business.

If the topic is significant for your business, complete the relevant checklist table and record the answers.

Read the Suggested Practices for the topic to explore your management options.

Management chapters	
1 Land and soil management	F.3
2 Water management	F.6
3 Chemical management	F.8
4 Nutrient management	F.9
5 Biodiversity	F.10
6 Waste management	F.11
7 Air management	F.12
8 Energy and greenhouse gas management	F.14

#### Date:

Are you aware of any targets, priorities or objectives of your Natural Resource Management Authority?

Have you used the Risk Assessments within each management chapter to ass important for your property?

Have you considered the relevant legislation requirements for your property?

1 Land and soil management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
1a Soil erosion caused by water						
1 An assessment of the risk of soil erosion by water has been undertaken						
2 Based on the result of the risk assessment, are further actions required?						
3 If so, what actions are needed or have l	been und	lertaken	?			
Soil cover maximised						
• Water control measures in place						
Soil structure improved						
Sediment traps in place						
1b Soil erosion caused by wind		Î	•	·	• •	•
5 An assessment of the risk of soil erosion by wind has been undertaken						
6 Based on the result of the risk assessment, are further actions required?						
7 If so, what actions are needed or have l	Deen und	Jertaken	?		·	·
Soil cover maximised						
<ul> <li>Windbreaks in place to moderate wind speed</li> </ul>						
Soil structure improved						
<ul> <li>Other management strategies implemented</li> </ul>						
8 The effectiveness of the actions is being monitored and recorded						
1c Soil structure						
9 An assessment of the soil structure has been undertaken						
10 Based on the result of the risk assessment, are further actions required?						
11 If so, what actions are needed or have	been ur	ndertake	n?			
12 The effectiveness of the actions is being monitored and recorded						

# Review checklist

	YES	NO
es Management group/Catchment		
sist you in identifying issues that might be		
?		

HAL

1 Land and soil management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
1d Salinity						•
13 An assessment of the risk of groundwater and soil salinity has been undertaken						
14 An assessment of the risk of irrigating with saline water has been undertaken						
15 Based on the result of the risk assessment, are further actions required?						
16 If so, what actions are needed or ha	ve been	undert	aken?			
<ul> <li>Determination of the source of salinity (irrigation water, groundwater or soil salinity)</li> </ul>						
<ul> <li>Advice sought on irrigation strategies</li> </ul>						
• Actions taken to lower water table						
<ul> <li>Actions taken to minimise rise in water table (including review of irrigation requirements and improvement of subsurface drainage)</li> </ul>						
<ul> <li>Alternate source of irrigation water or shandying / diluting saline water</li> </ul>						
Leaching fraction applied						
Vegetation cover maintained						
<ul> <li>Specialist advice sought</li> </ul>						
17 The effectiveness of the actions is being monitored and recorded						
1e Soil acidity and alkalinity						
18 An assessment of the risk of soil acidity or alkalinity has been undertaken						
19 Based on the result of the risk assessment, are further actions required?						
20 If so, what actions are needed or ha	ve been	undert	aken?			1
• Soil pH is monitored regularly						
The relative acidifying impact of fertilisers is considered when deciding on fertiliser purchase/application						
• The potential for leaching is considered prior to applying fertilisers or irrigation post-fertiliser application						
• Lime or dolomite is applied as required						
21 The effectiveness of the actions is being monitored and recorded						

	Land and soil management	Yes	No	Not relevant to my property	
1f	Sodicity				
22	An assessment of the risk of sodicity has been undertaken				
23	Based on the result of the risk assessment, are further actions required?				
24	If so, what actions are needed or ha	ve been	undert	aken?	
•	Gypsum or lime applied (dependent on soil pH)				
•	Soils not deep ripped				
•	Surface drainage improved or raised beds used				
•	Irrigation water slightly saline				
•	Specialist advice sought				
25	The effectiveness of the actions is being monitored and recorded				

Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months



2 Water management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required In the next 12 months
2a Irrigation efficiency					·	•
26 An assessment of the irrigation efficiency has been undertaken						
27 Based on the result of the risk assessment, are further actions required?						
28 If so, what actions are needed or have	ve been	underta	aken?		•	^
<ul> <li>Property goal defined</li> </ul>						
• Soil survey undertaken						
<ul> <li>Irrigation system suitable for property goal and is efficient way of applying water to crop</li> </ul>						
Water budget developed						
<ul> <li>Crop water requirements and water availability understood</li> </ul>						
<ul> <li>Irrigation schedule developed</li> </ul>						
<ul> <li>Soil moisture measurement/ assessment carried out</li> </ul>						
<ul> <li>Strategies implemented to manage nutrient input and salinity</li> </ul>						
29 The effectiveness of the actions is being monitored and recorded						
2b Water quality	-					
30 An assessment of irrigation water quality has been undertaken						
31 An assessment of the risk to downstream water quality has been undertaken						
32 Based on the result of the risk assessments, are further actions required?						
33 If so, what actions are needed or ha	ve been	undert	aken?			
• Irrigation water quality tested						
<ul> <li>Quality of water leaving propertytested</li> </ul>						
<ul> <li>Irrigation water quality tested</li> </ul>						
Specialist advice sought						
<ul> <li>Quality of water leaving property tested</li> </ul>						
<ul> <li>Water quality tests take place at appropriate times</li> </ul>						
• Watercourses are protected						
• Soil erosion is controlled						

2 Water management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required In the next 12 months
2b Water quality cont.						
<ul> <li>Fertilisers are appropriately stored</li> </ul>						
<ul> <li>Nutrient application is appropriately managed</li> </ul>						
<ul> <li>Nutrient application is appropriately managed</li> </ul>						
<ul> <li>Environmental impact of fertiliser application and storage is minimised</li> </ul>						
<ul> <li>Agricultural chemicals are appropriately stored</li> </ul>						
<ul> <li>Agricultural chemicals are appropriately applied (spray drift and contamination of waterways are considered)</li> </ul>						
<ul> <li>Fuels and oils are appropriately stored and transported (mobile fuel tanks)</li> </ul>						
<ul> <li>Septic tanks and manure storage areas well away from watercourses</li> </ul>						
34 The effectiveness of the actions is being monitored and recorded						
2c Managing wastewater						
35 An assessment of the risk of mismanaging wastewater has been undertaken						
36 Based on the result of the risk assessment, are further actions required?						
37 If so, what actions are needed or hav	e been	underta	ken?			
• Drains and water storage facilities in place						
• Wastewater is recycled or reused						
• Wastewater is treated						
38 The effectiveness of the actions is being monitored and recorded						



3 Chemical management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
39 An assessment of the risk of misapplication, incorrect handling, storage, application and disposal of chemicals, surplus spray and chemical containers has been undertaken						
40 An assessment of the risk of spray drift has been undertaken						
41 Based on the result of the risk assessment, are further actions required?						
42 If so, what actions are needed or ha	ive been	undert	aken?		•	
<ul> <li>IPM strategies employed</li> </ul>						
Chemicals safely stored						
<ul> <li>Materials available to deal with chemical spillage</li> </ul>						
<ul> <li>Chemicals used responsibly and by trained personnel</li> </ul>						
<ul> <li>Mixing and washdown areas appropriate</li> </ul>						
<ul> <li>Spraying when weather conditions are least likely to cause spray drift</li> </ul>						
<ul> <li>Using appropriate drift minimisation strategies (e.g. large droplets, not spraying next to sensitive areas, boom kept low, modifying spray equipment)</li> </ul>						
• Establishing barriers and/or buffer zones to minimise spray drift						
Water supplies protected from chemical contamination						
<ul> <li>Appropriate disposal of empty chemical containers</li> </ul>						
<ul> <li>Appropriate disposal of surplus chemicals, dip tank contents and washings</li> </ul>						
<ul> <li>Appropriate disposal of old, de- registered or unwanted chemicals</li> </ul>						
<ul> <li>Approriate disposal of other chemical products (e.g. rodent baits) or contaminated goods</li> </ul>						
<ul> <li>Fuels are safely stored away from watercourses</li> </ul>						
43 The effectiveness of the actions is being monitored and recorded						

4 Nutrient management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
4a Nutrient requirements						
44 A risk assessment of appropriate nutrient application has been undertaken						
45 Based on the result of the risk assessment, are further actions required?						
46 If so, what actions are needed or have been undertaken?						
Nutrient requirements of crop understood						
• Soil, sap or leaf test undertaken						
<ul> <li>Test results assessed by a suitably qualified/competent person and appropriate recommendations made</li> </ul>						
Nutrient budget developed						
<ul> <li>Nutrients applied do not exceed crop needs (as indicated by soil, leaf and/or sap tests)</li> </ul>						
• Fertilisers low in contaminants are used						
47 The effectiveness of the actions is being monitored and recorded						
4b Nutrient application						
48 An assessment of potential nutrient losses to the environment (leaching, run-off, wind) has been undertaken						
49 Based on the result of the risk assessment, are further actions required?						
50 If so, what actions are needed or have be	en unde	rtaken?				
<ul> <li>Type, rate and timing of application of fertiliser are appropriate</li> </ul>						
<ul> <li>Nutrients that are most readily lost (N and K) are applied in small amounts often to match plant growth</li> </ul>						
- Fertiliser applications are timed to match crop growth/need						
• Fertiliser placement appropriate						
<ul> <li>Fertilisers applied where crop can most easily use them</li> </ul>						
<ul> <li>Fertiliser application equipment regularly calibrated and maintained</li> </ul>						
<ul> <li>Fertilisers are safely stored, protected from direct sunlight and rain and away from watercourses</li> </ul>						
51 The effectiveness of the actions is being monitored and recorded						

HAL



5 Biodiversity	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
52 An assessment of risk to biodiversity has been undertaken						
53 Based on the result of the risk assessment, are further actions required?						
54 If so, what actions are needed or ha	ve been	undert	aken?			
<ul> <li>Native vegetation on the property is identified</li> </ul>						
<ul> <li>Native vegetation on surrounding properties is identified</li> </ul>						
<ul> <li>Initial assessment of the importance of native flora and fauna on property and nearby undertaken</li> </ul>						
<ul> <li>Biodiversity laws and regulations checked</li> </ul>						
<ul> <li>Threats to native biodiversity identified</li> </ul>						
<ul> <li>Strategies developed for managing 'problem' plants and native animals</li> </ul>						
<ul> <li>Practical ways to manage native vegetation developed</li> </ul>						
- habitats protected						
- habitats improved						
- habitats created						
<ul> <li>Soil biodiversity encouraged</li> </ul>						
• Join forces with others						
55 The effectiveness of the actions is being monitored and recorded						

6 Waste management	Yes	No	Not relevant to my property	Needs attention	Actions taken in the past 12 months	Actions required in the next 12 months
56 An assessment of the opportunities to minimise waste has been undertaken						
57 Based on the result of the risk assessment, are further actions required?						
58 If so, what actions are needed or have been undertaken?						
<ul> <li>Sources of waste identified and prioritised</li> </ul>						
<ul> <li>Waste avoidance or minimisation strategies in place</li> </ul>						
<ul> <li>Reuse or recycling carried out wherever possible</li> </ul>						
<ul> <li>Disposal methods appropriate for particular types of waste</li> </ul>						
59 The effectiveness of the actions is being monitored and recorded						

HAL



7 Air management	Yes	No	Not relevant to my property	Needs attention	Actions taken In the past 12 months	Actions required In the next 12 months	
7a Odour management							
60 An assessment of the risk of odour has been undertaken							
61 Based on the result of the risk assessment, are further actions required?							
62 If so, what actions are needed or hav	ve been	undert	aken?				
<ul> <li>Composted manure used in preference to raw manure</li> </ul>							
<ul> <li>Animal manure stored to minimise concern to neighbours/community/ employees</li> </ul>							
<ul> <li>Manure applied in a manner that minimises concern to neighbours/ community/ employees</li> </ul>							
<ul> <li>Waste produce disposed of in a manner that minimises concern to neighbours/ community/employees</li> </ul>							
63 The effectiveness of the actions is being monitored and recorded							
7b Dust management							
64 An assessment of the risk of dust has been undertaken							
65 Based on the result of the risk assessment, are further actions required?							
66 If so, what actions are needed or have been undertaken?							
• Shelterbelts and windbreaks in place							
<ul> <li>Appropriate cultivation practices, minimising working soil to fine tilth in dry windy weather</li> </ul>							
<ul> <li>Mulches applied to seedbeds</li> </ul>							
• Traffic ways wetted down or sealed							
67 The effectiveness of the actions is being monitored and recorded							
7c Smoke management							
68 An assessment of the risk of smoke has been undertaken							
69 Based on the result of the risk assessment, are further actions required?							
70 If so, what actions are needed or have been undertaken?							
<ul> <li>Alternate disposal options explored for materials that create dark smoke</li> </ul>							

7 Air management	Yes	No	Not relevant to my property	
7c Smoke management cont				
<ul> <li>Materials burnt when smoke will cause minimal concern to neighbours/community</li> </ul>				
<ul> <li>Materials burnt in a manner that minimises smoke creation</li> </ul>				
71 The effectiveness of the actions is being monitored and recorded				
7d Noise management				
72 An assessment of the risk has been undertaken				
73 Based on the result of the risk assessment, are further actions required?				
74 If so, what actions are needed or h	ave beer	n under	taken?	
Buffer zones established				
<ul> <li>Reducing noise from equipment (e.g. use electric motors, muffle equipment, sound proofing)</li> </ul>				
<ul> <li>Minimising noise generation at times that causes concern to neighbours/community</li> </ul>				

75	The effectiveness of the actions is		
	being monitored and recorded		

HAL

Needs attention	Actions taken In the past 12 months	Actions required In the next 12 months



8 Energy and greenhouse gas management	Yes	No	Not relevant to my property	Needs attention	Actions taken In the past 12 months	Actions required In the next 12 months
80 An assessment of the opportunities to minimise energy usage has been undertaken (carbon footprint or energy audit)						
81 Based on the result of the risk assessment, are further actions required?						
82 If so, what actions are needed or ha	ve been	undert	aken?			
<ul> <li>Carbon footprint or Energy audit undertaken and action plan developed</li> </ul>						
<ul> <li>Irrigation equipment appropriate for flow required and regularly maintained</li> </ul>						
<ul> <li>Equipment and vehicles are maintained and efficienct</li> </ul>						
<ul> <li>Turn off any machinery or equipment that is not being used</li> </ul>						
<ul> <li>Fuel usage monitored, LPG and natural gas used where possible</li> </ul>						
• Energy-efficient lighting						
<ul> <li>Efficient coolroom design and equipment regularly maintained and calibrated</li> </ul>						
<ul> <li>Where possible renewable resources used or greenpower purchased</li> </ul>						
<ul> <li>Reducing nitrous oxide emissions through efficient use of nitrogenous fertilisers</li> </ul>						
<ul> <li>Reducing CFC emissions through maintenance of refrigeration equipment</li> </ul>						
83 The effectiveness of the actions is being monitored and recorded						

